

**Homework 0 – Probability Review**  
**Due Friday, Sept. 9th at 11:59pm**

This homework is to be completed *individually* and the score will be **based on the problems attempted**, not the problems that are correctly answered. *Note: Problem 12 is required for credit.*

1. Brownstown Central High School has 500 seniors (300 boys and 200 girls). The number who pass and fail the final exam in a particular course is given below:

Number of students

	Fail	Pass	Total
Boys	28	272	300
Girls	10	190	200

If one of the 500 seniors is chosen at random (i.e., if each senior has a  $1/500$  chance of being chosen), D and A denote the events that the senior fails or passes, respectively; while B and G denote the events that the senior is a boy or girl. Calculate each of the following probabilities:

- $P(D)$
  - $P(A \text{ or } B)$
  - $P(A \text{ and } B)$
  - $P(B|A)$
  - $P(D|G)$
  - $P(A | \text{not } G)$
  - Is the event denoted by D independent of the event denoted by G? Why, or why not?
  - Is the event denoted by A independent of the event denoted by G? Why, or why not?
2. A machine repair can be performed either on time or late, and either satisfactorily or unsatisfactorily. The probability of a repair being both on time and satisfactory is 0.26. The probability of a repair being on time is 0.76. The probability of a repair being satisfactory is 0.41.
- What is the probability of a repair being both late and unsatisfactory?
  - What is the probability that a repair is on time, given that the repair is satisfactory?


3. Seventy percent of the light aircraft that disappear while in flight in a certain country are subsequently discovered. Of the aircraft that are discovered, 60% have an emergency locator, whereas 90% of the aircraft not discovered do not have such a locator. Suppose a light aircraft has disappeared.
- If it has an emergency locator, what is the probability that it will not be discovered?
  - If it does not have an emergency locator, what is the probability that it will be discovered?
4. Determine whether the events in the following situations are independent:
- In a normal 52 card deck pick one card at random. Let E be the event that the selected card is an ace and let F be the event that it is a spade.
  - Two fair coins are flipped. Let E be the event that the first coin lands heads. Let F be the event that the second lands tails.
  - Suppose two fair dice are tossed. Let E denote the event that the sum of the dice is 7 and let F denote the event that the first die equals 4.
5. If  $P(A|B) = 1$ , must  $A = B$ ? Use a Venn diagram to explain your answer.
6. A batch of 500 containers of orange juice contains 5 that are past the expiration date. Two are selected at random, without replacement, from the batch.
- What is the probability that the second one selected is expired given that the first one was expired?
  - What is the probability that both are expired?
  - What is the probability that neither are past expiration?

7. Suppose that a table of part counts is as follows:

		Conforms?	
		Yes	No
Supplier	1	ka	kb
	2	a	b

Where a, b, and k are positive integers. Let A denote the event that a part is from supplier 1 and let B denote the event that a part conforms to specifications.

- What is  $P(A|B)$ ?
- What is  $P(\bar{B}|A)$ ?
- Are A and B independent? Explain mathematically.

8. Customers are used to evaluate preliminary product designs. In the past, 95% of highly successful products received good reviews, 60% of moderately successful products received good reviews, and 10% of poor products received good reviews. In addition, 40% of products have been highly successful, 35% have been moderately successful, and 25% have been poor products. 

- What is the probability that a product attains a good review?
- If a new design attains a good review, what is the probability that it will be a highly successful product?
- If a product does not attain a good review, what is the probability that it will be a highly successful product?

9. A card is drawn from a standard pack of 52 well-shuffled cards (which contains 4 kings). What is the probability that it is a king? Without replacing this first king card, a second card is drawn. What is the probability that the second card pulled is a king? What is the probability that the first four cards drawn from a standard deck of 52 well-shuffled cards are all kings? Once drawn, a card is not replaced in the deck.
10. A card is drawn at random from a standard deck of 52 well-shuffled cards. Let  $A$  be the event that the card drawn is a queen and let  $B$  the event that the card pulled is red. Find the probabilities of the following events and state in words what they represent.
- a.  $A \cap B$   
b.  $A \cup B$
11. A factory has three machines that manufacture widgets. The percentages of a total day's production manufactured by the machines are 10%, 35%, and 55%, respectively. Furthermore, it is known that 5%, 3%, and 1% of the outputs of the respective three machines are defective. What is the probability that a randomly selected widget at the end of the day's production runs will be defective?
12. **Required Problem:** Find one example of a stochastic process in practice and summarize the example on a single powerpoint slide. On the slide: (1) Briefly describe the example, (2) explain what is uncertain, (3) explain why you think it is a stochastic process. This example can be from industry, it can be from the newspaper, it can be something that you have personally experienced, etc.