

Homotopy Theory

Frank Kong, 2017012181

November 26, 2019

Preface

Introduction

This is the lecture note to the course Homotopy Theory. The course started in the autumn semester in 2018, and 46 lectures has been given so far. The statements given are unproved, the proofs can be found in references below.

References

- The main page of the course on Banana Space:
<https://www.bananaspace.net/page/51800441>.
- The main page of the course on GitHub:
<https://github.com/Frank990919/Homotopy-Theory>.
- Hovey: *Model Categories*.
- Lurie: *Higher Algebra*.
- Lurie: *Higher Topos Theory*.
- The nLab: <https://ncatlab.org/>.
- Lecture notes for the lecture *Homotopic Algebra and Homological Algebra*, by Bu Chenjing:
Banana Space version: <https://www.bananaspace.net/page/25842635>;
GitHub version: <https://github.com/abccsss/Homotopical-Algebra>.

Acknowledgements

The lecturer would like to thank Bu Chenjing for giving me support on the lecture, and discussions, ideas and thoughts that helps the lecturer to understand the subject better and the notes more readable.

Contents

1	Model Categories (Semester 1)	3
1.1	Basic Concepts	3
1.1.1	Model Categories and Homotopy Categories	3
1.1.2	Quillen Adjunctions and Derived Functors	7
1.1.3	2-Categories	9
1.2	Examples of Model Categories	12
1.2.1	Cofibrantly Generated Model Categories	12
1.2.2	The Category of Modules over a Frobenius Ring	14
1.2.3	The Category of Chain Complexes of Modules over a Ring	15
1.2.4	The Category of Topological Spaces	16
1.3	Simplicial Sets	18
1.3.1	The Definition of Simplicial Sets	18
1.3.2	The Standard Model Structure on Simplicial Sets	20
1.4	Monoidal Model Categories	24
1.4.1	Monoidal Categories	24
1.4.2	Monoidal Model Categories	28
1.5	Framings	33
1.5.1	Diagrams over Reedy Categories	33
1.5.2	Homotopy Categories as $\text{Ho } \underline{\text{SSet}}$ -Modules	36
1.6	Pointed Model Categories	42
1.6.1	The Suspension and Loop Functors	42
1.6.2	Cofiber and Fiber Sequences	45
1.6.3	Pre-Triangulated Categories	49
1.7	Triangulated Categories and Stable Homotopy Categories	53
1.7.1	Triangulated Categories	53
1.7.2	Stable Homotopy Categories	55
2	Infinity Categories (Semesters 2 & 3)	56
2.0	Preliminaries	56
2.0.1	Preliminaries for Ordinary Categories	56
2.0.2	Preliminaries for Model Categories	56
2.1	Introduction to Infinity Categories	56
2.1.1	Foundations of Infinity Categories	56
2.1.2	The Language of Infinity Categories	62

Chapter 1

Model Categories (Semester 1)

The following contents are given in the lectures in the autumn semester of 2018. 21 lectures with a total time of 70 hours were given to finish the book *Model Categories*.

1.1 Basic Concepts

1.1.1 Model Categories and Homotopy Categories

Definition 1.1.1.1. Suppose $f : A \rightarrow B, g : C \rightarrow D$ are morphisms in \mathcal{C} . We say f is a **retract** of g if there exists a commutative diagram

$$\begin{array}{ccccc} A & \xrightarrow{p} & C & \xrightarrow{q} & A \\ f \downarrow & & \downarrow g & & \downarrow f \\ B & \xrightarrow{r} & D & \xrightarrow{s} & B \end{array}$$

such that $sr = 1, qp = 1$.

Definition 1.1.1.2. A **functorial factorization** of a category \mathcal{C} is an ordered pair (α, β) of functors $\mathcal{C}^{[1]} \rightarrow \mathcal{C}^{[1]}$ such that $d\alpha = d, c\alpha = d\beta, c\beta = c$, where $d, c : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ are the domain functor and the codomain functor, respectively, and for any $f \in \mathcal{C}^{[1]}$, $f = \beta f \circ \alpha f$. Here $[1]$ denotes the category with 2 objects and 3 morphisms.

Definition 1.1.1.3. Suppose $i : A \rightarrow B$ and $p : X \rightarrow Y$ are morphisms in \mathcal{C} . We say i has **left lifting property (LLP)** with respect to p , and p has **right lifting property (RLP)** with respect to i , written $i \nearrow p$, if for any commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & X \\ i \downarrow & & \downarrow p \\ B & \xrightarrow{g} & Y, \end{array}$$

there exists some $h : B \rightarrow X$ such that $ph = g, hi = f$.

Definition 1.1.1.4. Suppose \mathcal{C} is a category. A **model structure** on \mathcal{C} consists of three classes of morphisms $\text{Cof}, \text{Fib}, \text{W}$ together with two functorial factorizations $(\alpha, \beta), (\gamma, \delta)$, such that:

- i) $\text{Cof}, \text{Fib}, \text{W}$ are closed under compositions and retracts.
- ii) For any morphisms f, g such that gf is defined, if two of the three morphisms f, g, gf are in W , so is the third. (This is called the **2-out-of-3 property**)

iii) $\text{Cof} \cap W \nearrow \text{Fib}, \text{Cof} \nearrow \text{Fib} \cap W$.

iv) For any morphism f , $\alpha f \in \text{Cof}, \beta f \in \text{Fib} \cap W, \gamma f \in \text{Cof} \cap W, \delta f \in \text{Fib}$.

Morphisms in $\text{Cof}, \text{Fib}, W, \text{Cof} \cap W, \text{Fib} \cap W$ are called **cofibrations**, **fibrations**, **weak equivalences**, **trivial cofibrations**, **trivial fibrations**, respectively.

Definition 1.1.1.5. A **model category** is a category \mathcal{C} with all small limits and colimits together with a model structure on \mathcal{C} .

Example 1.1.1.6. Suppose \mathcal{C} is an arbitrary category, then the following structures are all model structures on \mathcal{C} :

- i) $\text{Cof} = \text{Iso } \mathcal{C}, \text{Fib} = W = \text{Mor } \mathcal{C}, \alpha f = \gamma f = \mathbb{1}_{df}, \beta f = \delta f = f$;
- ii) $\text{Fib} = \text{Iso } \mathcal{C}, \text{Cof} = W = \text{Mor } \mathcal{C}, \alpha f = \gamma f = f, \beta f = \delta f = \mathbb{1}_{cf}$;
- iii) $W = \text{Iso } \mathcal{C}, \text{Cof} = \text{Fib} = \text{Mor } \mathcal{C}, \gamma f = \mathbb{1}_{df}, \alpha f = \delta f = f, \beta f = \mathbb{1}_{cf}$,

where $\text{Mor } \mathcal{C}$ is the class of all morphisms in \mathcal{C} , $\text{Iso } \mathcal{C}$ is the class of all isomorphisms in \mathcal{C} . These model structures are called the **trivial model structures**.

Example 1.1.1.7. If $\mathcal{C}_i (i \in I)$ are model categories and I is a set, then $\prod_{i \in I} \mathcal{C}_i$ is a model category, where $\text{Cof} = \prod_{i \in I} \text{Cof}_{\mathcal{C}_i}, \text{Fib} = \prod_{i \in I} \text{Fib}_{\mathcal{C}_i}, W = \prod_{i \in I} W_{\mathcal{C}_i}$. The model structure on $\prod_{i \in I} \mathcal{C}_i$ is called the **product model structure**.

Example 1.1.1.8. If \mathcal{C} is a model category, then \mathcal{C}^{op} is a model category, where $\text{Cof}_{\mathcal{C}^{\text{op}}} = \text{Fib}_{\mathcal{C}}^{\text{op}}, \text{Fib}_{\mathcal{C}^{\text{op}}} = \text{Cof}_{\mathcal{C}}^{\text{op}}, W_{\mathcal{C}^{\text{op}}} = W_{\mathcal{C}}^{\text{op}}$. This model structure called the **dual model structure**.

Definition 1.1.1.9. Suppose \mathcal{C} is a model category. An object A is called **cofibrant** if the morphism $\phi \rightarrow A$ is a cofibration, where ϕ is the initial object; it is called **fibrant** if the morphism $A \rightarrow *$ is a fibration, where $*$ is the terminal object.

Definition 1.1.1.10. A category \mathcal{C} is called **pointed** if the initial object and the terminal object are isomorphic.

Definition 1.1.1.11. Suppose \mathcal{C} is a category with terminal object $*$. Define the **pointed category** of \mathcal{C} , written \mathcal{C}_* , to be the category with objects $\{(X, v) | X \in \mathcal{C}, v : * \rightarrow X\}$, and $\text{Hom}_{\mathcal{C}_*}((X, v), (Y, w)) = \{f : X \rightarrow Y | fv = w\}$. If $(X, v) \in \mathcal{C}_*$, we denote v the **basepoint** of X . Define the functor $-_+ : \mathcal{C} \rightarrow \mathcal{C}_*$ to be the functor that takes X to $X \amalg *$ with basepoint $*$.

Lemma 1.1.1.12. For any category \mathcal{C} with terminal object $*$, \mathcal{C}_* is pointed. Moreover, \mathcal{C} is pointed if and only if $-_+$ is an isomorphism. Furthermore, $-_+$ is left adjoint to the forgetful functor $U : \mathcal{C}_* \rightarrow \mathcal{C}, (X, v) \mapsto X$.

Lemma 1.1.1.13. For any category \mathcal{C}, \mathcal{D} with terminal objects, there is a functor

$$-_* : \text{Adj}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Adj}(\mathcal{C}_*, \mathcal{D}_*), (F, G, \varphi) \mapsto (F_*, G_*, \varphi_*),$$

where $G_*(Y, w) = (GY, Gw)$ and $F_*(X, v)$ is defined by the pushout diagram:

$$\begin{array}{ccc} F* & \xrightarrow{Fv} & FX \\ \downarrow & & \downarrow \\ * & \longrightarrow & F_*(X, v) \end{array}$$

Furthermore, $F_*(-_+)$ and $(F-)_+$ are naturally isomorphic.

Proposition 1.1.1.14. *If \mathcal{C} is a model category, then \mathcal{C}_* is a model category, where*

$$\text{Cof}_{\mathcal{C}_*} = U^{-1}(\text{Cof}_{\mathcal{C}}), \text{Fib}_{\mathcal{C}_*} = U^{-1}(\text{Fib}_{\mathcal{C}}), \text{W}_{\mathcal{C}_*} = U^{-1}(\text{W}_{\mathcal{C}}).$$

Definition 1.1.1.15. Suppose \mathcal{C} is a model category. We define the functor $Q : \mathcal{C} \rightarrow \mathcal{C}, X \mapsto d\beta(\phi \rightarrow X)$ to be the **cofibrant replacement functor**, and the functor $R : \mathcal{C} \rightarrow \mathcal{C}, X \mapsto c\gamma(X \rightarrow *)$ to be the **fibrant replacement functor**.

Lemma 1.1.1.16. *Suppose \mathcal{C} is a model category. Then there is a natural transformation $q : Q \rightarrow \mathbb{1}$ with $q_X = \beta(\phi \rightarrow X)$ for any X , and a natural transformation $r : \mathbb{1} \rightarrow R$ with $r_X = \gamma(X \rightarrow *)$ for any X . Moreover q_X is a trivial fibration and r_X is a trivial cofibration. Furthermore, Q, R preserves and reflects weak equivalences.*

Lemma 1.1.1.17 (The Retract Argument). *Suppose \mathcal{C} is a category, and we have a factorization $f = pi$. If $f \nearrow p$ then f is a retract of i . Dually if $i \nearrow f$ then f is a retract of p .*

Lemma 1.1.1.18. *Suppose \mathcal{C} is a model category. Then $f \nearrow \text{Fib} \cap \text{W}$ if and only if $f \in \text{Cof}$, $f \nearrow \text{Fib}$ if and only if $f \in \text{Cof} \cap \text{W}$, $\text{Cof} \cap \text{W} \nearrow f$ if and only if $f \in \text{Fib}$, $\text{Cof} \nearrow f$ if and only if $f \in \text{Fib} \cap \text{W}$.*

Corollary 1.1.1.19. *Suppose \mathcal{C} is a model category. Then Cof , $\text{Cof} \cap \text{W}$ are closed under pushouts, Fib , $\text{Fib} \cap \text{W}$ are closed under pullbacks.*

Lemma 1.1.1.20 (Ken Brown's Lemma). *Suppose \mathcal{C} is a model category, \mathcal{D} is a **category with weak equivalences**, i.e. a category with a class of morphisms W called weak equivalences, such that W is closed under composition and satisfies 2-out-of-3 property. If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor that takes trivial cofibrations between cofibrant objects to weak equivalences, then F takes weak equivalences between cofibrant objects to weak equivalences. Dually if $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor that takes trivial fibrations between fibrant objects to weak equivalences, then F takes weak equivalences between fibrant objects to weak equivalences.*

Definition 1.1.1.21. Suppose \mathcal{C} is a model category. Define $\mathcal{C}_c, \mathcal{C}_f$ and \mathcal{C}_{cf} to be the full subcategory of cofibrant, fibrant, cofibrant and fibrant objects of \mathcal{C} , respectively.

Definition 1.1.1.22. Suppose \mathcal{C} is a model category, $A \in \mathcal{C}$. A' is called a **cylinder object** of A if the map $A \amalg A \rightarrow A$ factors through A' : $A \amalg A \xrightarrow{i_0 + i_1} A' \xrightarrow{\sigma} A$, such that $i_0 + i_1 \in \text{Cof}$ and $\sigma \in \text{W}$. A'' is called a **path object** of A if the map $A \rightarrow A \times A$ factors through A'' : $A \xrightarrow{s} A'' \xrightarrow{(p_0, p_1)} A \times A$, such that $s \in \text{W}$ and $(p_0, p_1) \in \text{Fib}$.

Lemma 1.1.1.23. *Suppose \mathcal{C} is a model category, $A \in \mathcal{C}$, if A' is a cylinder object or a path object of A in \mathcal{C} , then A' is a path object or a cylinder object of A in \mathcal{C}^{op} , respectively.*

Lemma 1.1.1.24. *Suppose \mathcal{C} is a model category. There is a functor $\text{Cyl} : \mathcal{C} \rightarrow \mathcal{C}, A \mapsto d \circ \beta(A \amalg A \rightarrow A)$, such that $\text{Cyl}(A)$ is a cylinder object, and the structure map σ is a trivial fibration. Dually there is a functor $\text{Path} : \mathcal{C} \rightarrow \mathcal{C}, A \mapsto c \circ \gamma(A \rightarrow A \times A)$, such that $\text{Path}(A)$ is a path object, and the structure map s is a trivial cofibration.*

Lemma 1.1.1.25. *Suppose \mathcal{C} is a model category, $A \in \mathcal{C}$. Then for any cylinder object A' of A , there exists a map $A' \rightarrow \text{Cyl}(A)$ which is compatible with i_0, i_1, σ . Dually for any path object A'' of A , there exists a map $\text{Path}(A) \rightarrow A''$ which is compatible with p_0, p_1, s .*

Definition 1.1.1.26. Suppose \mathcal{C} is a model category, $f, g : A \rightarrow X$ are morphisms in \mathcal{C} . A **left homotopy** from f to g is a map $H : A' \rightarrow X$, where A' is some cylinder object of A and $H(i_0 + i_1) = f + g$. If such H exists for some cylinder object A' we say f is **left homotopic** to g , denoted $f \sim^l g$. Dually a **right homotopy** from f to g is a map $K : A \rightarrow X'$, where X' is

some path object of X and $(p_0, p_1)K = (f, g)$. If such K exists for some path object K' we say f is **right homotopic** to g , denoted $f \stackrel{r}{\sim} g$. Finally we say f is **homotopic** to g , denoted $f \simeq g$, if $f \stackrel{l}{\sim} g$ and $f \stackrel{r}{\sim} g$. Moreover we call f a **homotopy equivalence** if there exists $h : X \rightarrow A$ such that $fh \simeq 1_X, hf \simeq 1_B$.

Proposition 1.1.1.27. *Suppose \mathcal{C} is a model category, $f, g : B \rightarrow X, h : A \rightarrow B, k : X \rightarrow Y$ are morphisms in \mathcal{C} . If $f \stackrel{l}{\sim} g$ then $kf \stackrel{l}{\sim} kg$, and if in addition X is fibrant then $fh \stackrel{l}{\sim} gh$. Dually if $f \stackrel{r}{\sim} g$ then $fh \stackrel{r}{\sim} gh$, and if in addition B is cofibrant then $kf \stackrel{r}{\sim} kg$.*

Proposition 1.1.1.28. *Suppose \mathcal{C} is a model category, $A, X \in \mathcal{C}$. If A is cofibrant then $\stackrel{l}{\sim}$ is an equivalence relation on $\text{Hom}_{\mathcal{C}}(A, X)$. Dually if X is fibrant then $\stackrel{r}{\sim}$ is an equivalence relation on $\text{Hom}_{\mathcal{C}}(A, X)$.*

Proposition 1.1.1.29. *Suppose \mathcal{C} is a model category, $h : A \rightarrow B, k : X \rightarrow Y$ are morphisms in \mathcal{C} . If B is cofibrant and k is a trivial fibration or a weak equivalence between fibrant objects, then $k_* : \text{Hom}_{\mathcal{C}}(B, X) / \stackrel{l}{\sim} \rightarrow \text{Hom}_{\mathcal{C}}(B, Y) / \stackrel{l}{\sim}$ is a bijection. Dually if X is fibrant and h is a trivial cofibration or a weak equivalence between cofibrant objects, then $h^* : \text{Hom}_{\mathcal{C}}(B, X) / \stackrel{r}{\sim} \rightarrow \text{Hom}_{\mathcal{C}}(A, X) / \stackrel{r}{\sim}$ is a bijection.*

Proposition 1.1.1.30. *Suppose \mathcal{C} is a model category, $f, g : A \rightarrow X$ are morphisms in \mathcal{C} . If A is cofibrant and $f \stackrel{l}{\sim} g$ then $f \stackrel{r}{\sim} g$, and the right homotopy may pass any path object of X . Dually if X is fibrant and $f \stackrel{r}{\sim} g$ then $f \stackrel{l}{\sim} g$, and the left homotopy may pass any cylinder object of A .*

Corollary 1.1.1.31. *If \mathcal{C} is a model category, then \simeq is an equivalence relation on morphisms of \mathcal{C}_{cf} and is compatible with composition.*

Proposition 1.1.1.32 (Whitehead). *If \mathcal{C} is a model category, then a map in \mathcal{C}_{cf} is a weak equivalence if and only if it is a homotopy equivalence.*

Definition 1.1.1.33. Suppose \mathcal{C} is a category with a class of morphisms W such that W is closed under composition. For any $X, Y \in \mathcal{C}$, define $[X, Y]$ to be the class $F(X, Y) / \sim$, where $F(X, Y)$ is the class of all strings (f_1, \dots, f_n) of morphisms in \mathcal{C} or reversals of morphisms in W of finite length, such that $df_1 = X, cf_n = Y, df_i = cf_{i-1}$ for any $1 < i \leq n$; \sim is the equivalence relation generated from $() \sim (1), (f, g) \sim (g \circ f), (w^{-1}, v^{-1}) \sim ((w \circ v)^{-1}), () \sim (w, w^{-1}), () \sim (w^{-1}, w)$ for any f, g morphisms in \mathcal{C} and $w \in W$, and $(W_1, W_2, W_4) \sim (W_1, W_3, W_4)$ whenever W_1, W_2, W_3, W_4 are strings and $W_2 \sim W_3$. If for any $X, Y \in \mathcal{C}$, $[X, Y]$ is a set, define the **homotopy category** $\text{Ho } \mathcal{C}$ to be the category with objects being objects in \mathcal{C} and $\text{Hom}_{\text{Ho } \mathcal{C}}(X, Y) = [X, Y]$. We denote the obvious functor $\mathcal{C} \rightarrow \text{Ho } \mathcal{C}$ by γ .

Lemma 1.1.1.34. *Suppose \mathcal{C} is a category with a class of morphisms W such that W is closed under composition. If $\text{Ho } \mathcal{C}$ exists and $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor which sends morphisms in W to isomorphisms, then there exists a unique functor $\text{Ho } F : \text{Ho } \mathcal{C} \rightarrow \mathcal{D}$ such that $(\text{Ho } F)\gamma = F$. Conversely if $\alpha : \mathcal{C} \rightarrow \mathcal{E}$ is a functor which sends morphisms in W to isomorphisms and satisfies the above universal property, then $\text{Ho } \mathcal{C}$ exists and there exists a unique isomorphism $\delta : \text{Ho } \mathcal{C} \rightarrow \mathcal{E}$ such that $\delta\gamma = \alpha$.*

Lemma 1.1.1.35. *Suppose \mathcal{C} is a category with a class of morphisms W such that W is closed under composition, and $\text{Ho } \mathcal{C}$ exists. If $\tau : F \rightarrow G$ is a natural transformation between functors which sends morphisms in W to isomorphisms, then $\text{Ho } \tau : \text{Ho } F \rightarrow \text{Ho } G$ with $\text{Ho } \tau_X = \tau_X$ is a natural transformation.*

Proposition 1.1.1.36. *Suppose \mathcal{C} is a model category, then $\text{Ho } \mathcal{C}_{cf}$ exists and there exists a unique isomorphism $\text{Ho } \mathcal{C}_{cf} \rightarrow \mathcal{C}_{cf} / \simeq$, such that the isomorphism is identity on objects.*

Proposition 1.1.1.37. Suppose \mathcal{C} is a model category. Then the inclusions induces isomorphisms $[X, Y]_{\mathcal{C}_{cf}} \cong [X, Y]_{\mathcal{C}_c} \cong [X, Y]_{\mathcal{C}}$ and $[X, Y]_{\mathcal{C}_{cf}} \cong [X, Y]_{\mathcal{C}_f} \cong [X, Y]_{\mathcal{C}}$ for any $X, Y \in \mathcal{C}_{cf}$.

Theorem 1.1.1.38. Suppose \mathcal{C} is a model category. Then:

i) $\text{Ho } \mathcal{C}$ exists, and we have equivalences between categories $\text{Ho } \mathcal{C}_{cf} \rightarrow \text{Ho } \mathcal{C}_c \rightarrow \text{Ho } \mathcal{C}$ and $\text{Ho } \mathcal{C}_{cf} \rightarrow \text{Ho } \mathcal{C}_f \rightarrow \text{Ho } \mathcal{C}$ induced by inclusions;

ii) We have for any $X, Y \in \mathcal{C}$,

$$\begin{aligned} [X, Y] &\cong (\text{Hom}_{\mathcal{C}}(QRX, QRY) / \simeq) \\ &\cong (\text{Hom}_{\mathcal{C}}(RQX, RQY) / \simeq) \\ &\cong (\text{Hom}_{\mathcal{C}}(QX, RY) / \simeq), \end{aligned}$$

and if moreover X is cofibrant and Y is fibrant, then $[X, Y] = \text{Hom}_{\mathcal{C}}(X, Y) / \simeq$;

iii) $\gamma : \mathcal{C} \rightarrow \text{Ho } \mathcal{C}$ identifies maps that are left homotopic or right homotopic;

iv) If f is a morphism in \mathcal{C} such that γf is an isomorphism, then f is a weak equivalence.

1.1.2 Quillen Adjunctions and Derived Functors

Definition 1.1.2.1. Suppose \mathcal{C} and \mathcal{D} are categories, $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is an adjunction. Define its **unit** to be the natural transformation $\eta : 1 \rightarrow GF, \eta_X = \varphi(FX \rightarrow FX)$, and its **counit** to be the natural transformation $\varepsilon : FG \rightarrow 1, \varepsilon_Y = \varphi^{-1}(GY \rightarrow GY)$.

Definition 1.1.2.2. Suppose \mathcal{C} and \mathcal{D} are model categories. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is called a **left Quillen functor** if it is a left adjoint and preserves cofibrations and trivial cofibrations. Dually a functor $G : \mathcal{D} \rightarrow \mathcal{C}$ is called a **right Quillen functor** if it is a right adjoint and preserves fibrations and trivial fibrations. An adjunction $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is called a **Quillen adjunction** if F is left Quillen.

Example 1.1.2.3. Suppose \mathcal{C} is a model category and I is a set, then the coproduct functor and the diagonal functor form a Quillen adjunction $\mathcal{C}^I \rightarrow \mathcal{C}$, and the diagonal functor and the product functor form a Quillen adjunction $\mathcal{C} \rightarrow \mathcal{C}^I$.

Example 1.1.2.4. Suppose \mathcal{C} is a model category, then $(-_+, U, \varphi) : \mathcal{C} \rightarrow \mathcal{C}_*$ is a Quillen adjunction.

Lemma 1.1.2.5. If \mathcal{C} and \mathcal{D} are model categories, then an adjunction $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction if and only if G is right Quillen.

Lemma 1.1.2.6. Quillen adjunctions are closed under compositions.

Lemma 1.1.2.7. If \mathcal{C} and \mathcal{D} are model categories and $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction, then $(G, F, \varphi^{-1}) : \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$ is a Quillen adjunction.

Proposition 1.1.2.8. If \mathcal{C} and \mathcal{D} are model categories and $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction, then $(F_*, G_*, \varphi_*) : \mathcal{C}_* \rightarrow \mathcal{D}_*$ is a Quillen adjunction.

Definition 1.1.2.9. Suppose \mathcal{C} and \mathcal{D} are model categories. The **left derived functor** of a left Quillen functor $F : \mathcal{C} \rightarrow \mathcal{D}$, denoted LF , is the composite $\text{Ho } \mathcal{C} \xrightarrow{\text{Ho } Q} \text{Ho } \mathcal{C}_c \xrightarrow{\text{Ho } F} \text{Ho } \mathcal{D}$. The **derived natural transformation** of a natural transformation $\tau : F \rightarrow F'$ between left Quillen functors, denoted $L\tau$, is the composite $\text{Ho } \tau \circ \text{Ho } Q$. Dually, the **right derived functor** of a right Quillen functor $G : \mathcal{D} \rightarrow \mathcal{C}$, denoted RG , is the composite $\text{Ho } \mathcal{D} \xrightarrow{\text{Ho } R} \text{Ho } \mathcal{D}_f \xrightarrow{\text{Ho } G} \text{Ho } \mathcal{C}$. The **derived natural transformation** of a natural transformation $\tau : G \rightarrow G'$ between right Quillen functors, denoted $R\tau$, is the composite $\text{Ho } \tau \circ \text{Ho } R$.

Lemma 1.1.2.10. If τ is a natural transformation between left Quillen functors, then $L\tau_X = \tau_{QX}$. Dually if τ is a natural transformation between right Quillen functors, then $R\tau_X = \tau_{RX}$.

Lemma 1.1.2.11. If $\tau : F \rightarrow F'$, $\tau' : F' \rightarrow F''$ are natural transformations between left Quillen functors, then $L(\tau' \circ \tau) = (L\tau') \circ (L\tau)$, and $L(1_F) = 1_{LF}$. Dually if $\tau : G \rightarrow G'$, $\tau' : G' \rightarrow G''$ are natural transformations between right Quillen functors, then $R(\tau' \circ \tau) = (R\tau') \circ (R\tau)$, and $R(1_G) = 1_{RG}$.

Theorem 1.1.2.12. For any model categories $\mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}$ and left Quillen functors $F : \mathcal{C} \rightarrow \mathcal{D}$, $F' : \mathcal{D} \rightarrow \mathcal{E}$, $F'' : \mathcal{E} \rightarrow \mathcal{F}$, there exists natural isomorphisms $\alpha_{\mathcal{C}} : L(1_{\mathcal{C}}) \rightarrow 1_{\text{Ho}\mathcal{C}}$ and $\mu_{F'F} : LF' \circ LF \rightarrow L(F' \circ F)$, given by $(\alpha_{\mathcal{C}})_X : QX \xrightarrow{q} X$ and $(\mu_{F'F})_X : F'QFQX \xrightarrow{F'qFQX} F'FQX$, such that the following three diagrams commute:

$$\begin{array}{ccc}
(LF'' \circ LF') \circ LF & \xrightarrow{\mu_{F''F'} \circ LF} & L(F'' \circ F') \circ LF & \xrightarrow{\mu_{(F'' \circ F')F}} & L((F'' \circ F') \circ F) \\
\parallel & & & & \parallel \\
LF'' \circ (LF' \circ LF) & \xrightarrow{LF'' \circ \mu_{F'F}} & LF'' \circ L(F' \circ F) & \xrightarrow{\mu_{F''(F' \circ F)}} & L(F'' \circ (F' \circ F))
\end{array}$$

$$\begin{array}{ccc}
L(1_{\mathcal{D}}) \circ LF & \xrightarrow{\mu_{1_{\mathcal{D}}F}} & L(1_{\mathcal{D}} \circ F) & LF \circ L(1_{\mathcal{C}}) & \xrightarrow{\mu_{F1_{\mathcal{C}}}} & L(F \circ 1_{\mathcal{C}}) \\
\alpha_{\mathcal{C}} \circ LF \downarrow & & \parallel & LF \circ \alpha_{\mathcal{C}} \downarrow & & \parallel \\
1_{\text{Ho}\mathcal{D}} \circ LF & \xlongequal{\quad} & LF & LF \circ 1_{\text{Ho}\mathcal{C}} & \xlongequal{\quad} & LF
\end{array}$$

Dually there is a statement for right derived functors.

Proposition 1.1.2.13. Suppose $\sigma : F \rightarrow G$ is a natural transformation between two left Quillen functors $\mathcal{C} \rightarrow \mathcal{D}$, $\tau : F' \rightarrow G'$ is a natural transformation between two left Quillen functors $\mathcal{D} \rightarrow \mathcal{E}$, then the following diagram is commutative:

$$\begin{array}{ccc}
LF' \circ LF & \xrightarrow{\mu_{F'F}} & L(F' \circ F) \\
L\tau * L\sigma \downarrow & & \downarrow L(\tau * \sigma) \\
LG' \circ LG & \xrightarrow{\mu_{G'G}} & L(G' \circ G)
\end{array}$$

Dually there is a statement for right derived functors.

Proposition 1.1.2.14. If \mathcal{C} and \mathcal{D} are model categories and $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction, then $(LF, RG, \text{Ho}\varphi) : \text{Ho}\mathcal{C} \rightarrow \text{Ho}\mathcal{D}$ is an adjunction. We denote $\text{Ho}(F, G, \varphi) = (LF, RG, \text{Ho}\varphi)$ and call it the **derived adjunction**.

Example 1.1.2.15. If \mathcal{C} is a model category, then the right derived functor of the product functor on \mathcal{C} is a product functor on $\text{Ho}\mathcal{C}$. Dually the left derived functor of the coproduct functor on \mathcal{C} is a coproduct functor on $\text{Ho}\mathcal{C}$.

Definition 1.1.2.16. Suppose \mathcal{C} and \mathcal{D} are model categories. A Quillen adjunction is called a **Quillen equivalence**, if for any X cofibrant in \mathcal{C} and Y fibrant in \mathcal{D} , a map $f : FX \rightarrow Y$ is a weak equivalence if and only if $\varphi(f) : X \rightarrow GY$ is a weak equivalence.

Proposition 1.1.2.17. Suppose \mathcal{C} and \mathcal{D} are model categories and $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction. Then the following statements are equivalent:

- i) (F, G, φ) is a Quillen equivalence.
- ii) For any X cofibrant in \mathcal{C} the map $X \xrightarrow{\eta} GFX \xrightarrow{Gr_{FX}} GRFX$ is a weak equivalence, for any Y fibrant in \mathcal{D} the map $FQGY \xrightarrow{FqGY} FGY \xrightarrow{\varepsilon} Y$ is a weak equivalence.

iii) $\text{Ho}(F, G, \varphi)$ is an adjoint equivalence of categories.

Corollary 1.1.2.18. Suppose $(F, G), (F', G), (F, G')$ are Quillen adjunctions between model categories \mathcal{C} and \mathcal{D} . If one of the three adjunctions is a Quillen equivalence, so are the other two.

Corollary 1.1.2.19. Suppose $F : \mathcal{C} \rightarrow \mathcal{D}, G : \mathcal{D} \rightarrow \mathcal{E}$ are left (resp. right) Quillen functors. If two of the functors F, G, GF are Quillen equivalences, so is the third.

Corollary 1.1.2.20. Suppose \mathcal{C} and \mathcal{D} are model categories and $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction. Then the following statements are equivalent:

- i) (F, G, φ) is a Quillen equivalence.
- ii) For any morphism f between cofibrant objects in \mathcal{C} with Ff a weak equivalence, f is a weak equivalence; for any Y fibrant in \mathcal{D} the map $FQGY \xrightarrow{Fq_{GY}} FGY \xrightarrow{\varepsilon} Y$ is a weak equivalence.
- iii) For any morphism g between fibrant objects in \mathcal{D} with Gg a weak equivalence, g is a weak equivalence; for any X cofibrant in \mathcal{C} the map $X \xrightarrow{\eta} GFX \xrightarrow{Gr_{FX}} GRFX$ is a weak equivalence.

Proposition 1.1.2.21. Suppose \mathcal{C} and \mathcal{D} are model categories and $F : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen equivalence. If the terminal object $*$ of \mathcal{C} is cofibrant, and F preserves terminal object, then $F_* : \mathcal{C}_* \rightarrow \mathcal{D}_*$ is a Quillen equivalence.

Proposition 1.1.2.22. Suppose $\tau : F \rightarrow G$ is a natural transformation between left (resp. right) Quillen functors. Then $L\tau$ (resp. $R\tau$) is a natural equivalence if and only if for any cofibrant (resp. fibrant) X , τ_X is a weak equivalence.

1.1.3 2-Categories

Definition 1.1.3.1. A **2-category** \mathcal{C} consists of three superclasses $\mathcal{C}_0, \mathcal{C}_1, \mathcal{C}_2$, each called the objects, the 1-morphisms (or morphisms) and the 2-morphisms, two maps $d : \mathcal{C}_1 \rightarrow \mathcal{C}_0, c : \mathcal{C}_2 \rightarrow \mathcal{C}_1$ called the domain maps, two maps $c : \mathcal{C}_1 \rightarrow \mathcal{C}_0, c : \mathcal{C}_2 \rightarrow \mathcal{C}_1$ called the codomain maps, two maps $1 : \mathcal{C}_0 \rightarrow \mathcal{C}_1, \mathcal{C}_1 \rightarrow \mathcal{C}_2$ called the identity maps, and three maps $\circ : \mathcal{C}_1 \times_{\mathcal{C}_0} \mathcal{C}_1 \rightarrow \mathcal{C}_1, \circ : \mathcal{C}_2 \times_{\mathcal{C}_1} \mathcal{C}_2 \rightarrow \mathcal{C}_2, * : \mathcal{C}_2 \times_{\mathcal{C}_0} \mathcal{C}_2 \rightarrow \mathcal{C}_2$, each called the composite, the vertical composite and the horizontal composite, where the pullbacks are taken along the maps c, d , the maps c, d and the maps c^2, d^2 , respectively, satisfying the following conditions:

- i) As maps $\mathcal{C}_2 \rightarrow \mathcal{C}_0$ we have $dc = dd, cd = cc$;
- ii) Either as a map $\mathcal{C}_0 \rightarrow \mathcal{C}_0$ or as a map $\mathcal{C}_1 \rightarrow \mathcal{C}_1$ we have $d1$ and $c1$ are the identity maps;
- iii) If f, g are 1-morphisms such that $dg = cf$ then $d(g \circ f) = df$ and $c(g \circ f) = cg$; if σ, τ are 2-morphisms such that $d\tau = c\sigma$ then $d(\tau \circ \sigma) = d\sigma$ and $c(\tau \circ \sigma) = c\tau$; if σ, τ are 2-morphisms such that $d^2\tau = c^2\sigma$ then $d(\tau * \sigma) = d\tau \circ d\sigma$ and $c(\tau \circ \sigma) = c\tau \circ c\sigma$;
- iv) For any 1-morphism f we have $1_{cf} \circ f = f = f \circ 1_{df}$; for any 2-morphism σ we have $1_{c\sigma} \circ \sigma = \sigma = \sigma \circ 1_{d\sigma}$, and $1_{1_{cc\sigma}} * \sigma = \sigma = \sigma * 1_{1_{dd\sigma}}$;
- v) If f, g, h are 1-morphisms such that $dg = cf, dh = cg$ then $(h \circ g) \circ f = h \circ (g \circ f)$; if ρ, σ, τ are 2-morphisms such that $d\sigma = c\rho, d\tau = c\sigma$ then $(\tau \circ \sigma) \circ \rho = \tau \circ (\sigma \circ \rho)$; if ρ, σ, τ are 2-morphisms such that $d^2\sigma = c^2\rho, d^2\tau = c^2\sigma$ then $(\tau * \sigma) * \rho = \tau * (\sigma * \rho)$;
- vi) If $\sigma, \tau, \sigma', \tau'$ are 2-morphisms such that $d\tau = c\sigma, d\tau' = c\sigma', d^2\tau = c^2\sigma'$, then $(\tau \circ \sigma) * (\tau' \circ \sigma') = (\tau * \tau') \circ (\sigma * \sigma')$;

- vii) For any objects A, B , $\text{Hom}_{\mathbb{C}}(A, B) := \{f \in \mathbb{C}_1 \mid df = A, cf = B\}$ is a class; for any 1-morphisms f, g , $\text{Hom}_{\mathbb{C}}(f, g) := \{\tau \in \mathbb{C}_2 \mid d\tau = f, c\tau = g\}$ is a class.

If $f \in \text{Hom}_{\mathbb{C}}(A, B)$ where A, B are objects, we denote $f : A \rightarrow B$, and if $\tau \in \text{Hom}_{\mathbb{C}}(f, g)$ where f, g are 1-morphisms, we denote $\tau : f \rightarrow g$.

Example 1.1.3.2. All categories, functors, natural transformations form a 2-category which we denote $\underline{\text{Cat}}$. All categories, adjunctions, natural transformations between left adjoints form a 2-category which we denote $\underline{\text{Cat}}_{ad}$. All model categories, Quillen adjunctions, natural transformations between left Quillen functors form a 2-category which we denote $\underline{\text{Model}}$.

Definition 1.1.3.3. Suppose \mathbb{C} and \mathbb{D} are 2-categories. A **(covariant) 2-functor** $F : \mathbb{C} \rightarrow \mathbb{D}$ consists of three maps $\mathbb{C}_0 \rightarrow \mathbb{D}_0, \mathbb{C}_1 \rightarrow \mathbb{D}_1, \mathbb{C}_2 \rightarrow \mathbb{D}_2$, satisfying the following conditions:

- i) Either as maps $\mathbb{C}_1 \rightarrow \mathbb{D}_0$ or as maps $\mathbb{C}_2 \rightarrow \mathbb{D}_1$ we have $Fd = dF, Fc = cF$;
- ii) For any object A we have $F(1_A) = 1_{FA}$, for any 1-morphism f we have $F(1_f) = 1_{Ff}$;
- iii) If f, g are 1-morphisms such that $dg = cf$ then $F(g \circ f) = Fg \circ Ff$; if σ, τ are 2-morphisms such that $d\tau = c\sigma$ then $F(\tau \circ \sigma) = F\tau \circ F\sigma$; if σ, τ are 2-morphisms such that $d^2\tau = c^2\sigma$ then $F(\tau * \sigma) = F\tau * F\sigma$.

A **contravariant 2-functor** $F : \mathbb{C} \rightarrow \mathbb{D}$ consists of three maps $\mathbb{C}_0 \rightarrow \mathbb{D}_0, \mathbb{C}_1 \rightarrow \mathbb{D}_1, \mathbb{C}_2 \rightarrow \mathbb{D}_2$, satisfying the following conditions:

- i) Either as maps $\mathbb{C}_1 \rightarrow \mathbb{D}_0$ or as maps $\mathbb{C}_2 \rightarrow \mathbb{D}_1$ we have $Fd = cF, Fc = dF$;
- ii) For any object A we have $F(1_A) = 1_{FA}$, For any 1-morphism f we have $F(1_f) = 1_{Ff}$;
- iii) If f, g are 1-morphisms such that $dg = cf$ then $F(g \circ f) = Ff \circ Fg$; if σ, τ are 2-morphisms such that $d\tau = c\sigma$ then $F(\tau \circ \sigma) = F\sigma \circ F\tau$; if σ, τ are 2-morphisms such that $d^2\tau = c^2\sigma$ then $F(\tau * \sigma) = F\sigma * F\tau$.

Example 1.1.3.4. $\underline{\text{Cat}}_{ad} \rightarrow \underline{\text{Cat}}, \mathbb{C} \mapsto \mathbb{C}, (F, G, \varphi) \mapsto F, \tau \mapsto \tau$ is a covariant 2-functor. $\underline{\text{Model}} \rightarrow \underline{\text{Cat}}_{ad}, \mathbb{C} \mapsto \mathbb{C}, (F, G, \varphi) \mapsto (F, G, \varphi), \tau \mapsto \tau$ is a covariant 2-functor.

Definition 1.1.3.5. Given a natural transformation $\tau : F \rightarrow F'$ between adjunctions (F, G) and (F', G') , we define its **dual natural transformation** $\tau^{\text{op}} : G' \rightarrow G$ by the composite $G'X \xrightarrow{\eta_{G'X}} GFG'X \xrightarrow{G\tau_{G'X}} GF'G'X \xrightarrow{G\varepsilon'_X} GX$.

Lemma 1.1.3.6. If $\tau : F \rightarrow F'$ is a natural transformation between adjunctions (F, G, φ) and (F', G', φ') , then the following diagram is commutative for any objects $X \in \mathbb{C}, Y \in \mathbb{D}$:

$$\begin{array}{ccc} \text{Hom}_{\mathbb{D}}(F'X, Y) & \xrightarrow{\tau_X^*} & \text{Hom}_{\mathbb{D}}(FX, Y) \\ \varphi' \downarrow & & \downarrow \varphi \\ \text{Hom}_{\mathbb{C}}(X, G'Y) & \xrightarrow{(\tau_Y^{\text{op}})^*} & \text{Hom}_{\mathbb{C}}(X, GY) \end{array}$$

In particular, τ is a natural equivalence if and only if τ^{op} is.

Lemma 1.1.3.7.

$$-^{\text{op}} : \left\{ \begin{array}{l} \underline{\text{Cat}}_{ad} \rightarrow \underline{\text{Cat}}_{ad}, \\ \mathbb{C} \mapsto \mathbb{C}^{\text{op}}, \\ (F, G, \varphi) \mapsto (G, F, \varphi^{-1}), \\ (\tau : (F, G) \rightarrow (F', G')) \mapsto \tau^{\text{op}} \end{array} \right.$$

is a contravariant 2-functor, and $(-^{\text{op}})^2$ is the identity 2-functor. Similarly

$$-^{\text{op}} : \begin{cases} \underline{\text{Model}} \rightarrow \underline{\text{Model}}, \\ \mathcal{C} \mapsto \mathcal{C}^{\text{op}}, \\ (F, G, \varphi) \mapsto (G, F, \varphi^{-1}), \\ (\tau : (F, G) \rightarrow (F', G')) \mapsto \tau^{\text{op}} \end{cases}$$

is a contravariant 2-functor, and $(-^{\text{op}})^2$ is the identity 2-functor. They are called the **duality 2-functors**.

Definition 1.1.3.8. Suppose \mathbb{C} and \mathbb{D} are 2-categories. A **pseudo-2-functor** $F : \mathbb{C} \rightarrow \mathbb{D}$ consists of three maps $\mathbb{C}_0 \rightarrow \mathbb{D}_0$, $\mathbb{C}_1 \rightarrow \mathbb{D}_1$, $\mathbb{C}_2 \rightarrow \mathbb{D}_2$, and 2-isomorphisms $\alpha_A : F(1_A) \rightarrow 1_{FA}$ for any object A , 2-isomorphisms $\mu_{gf} : Fg \circ Ff \rightarrow F(g \circ f)$ for any 1-morphisms f, g with $dg = cf$, satisfying the following conditions:

- i) Either as maps $\mathbb{C}_1 \rightarrow \mathbb{D}_0$ or as maps $\mathbb{C}_2 \rightarrow \mathbb{D}_1$ we have $Fd = dF, Fc = cF$;
- ii) For any 1-morphism f we have $F(1_f) = 1_{Ff}$; if σ, τ are 2-morphisms such that $d\tau = c\sigma$ then $F(\tau \circ \sigma) = F\tau \circ F\sigma$;
- iii) If f, g, h are 1-morphisms such that $dg = cf, dh = cg$ then the following diagrams are commutative:

$$\begin{array}{ccccc} (Fh \circ Fg) \circ Ff & \xrightarrow{\mu_{hg \circ Ff}} & F(h \circ g) \circ Ff & \xrightarrow{\mu_{(h \circ g)f}} & F((h \circ g) \circ f) \\ \parallel & & & & \parallel \\ Fh \circ (Fg \circ Ff) & \xrightarrow{Fh \circ \mu_{gf}} & Fh \circ F(g \circ f) & \xrightarrow{\mu_{h(g \circ f)}} & F(h \circ (g \circ f)) \\ \\ F(1_{cf}) \circ Ff & \xrightarrow{\mu_{1_{cf}f}} & F(1_{cf} \circ f) & Ff \circ F(1_{df}) & \xrightarrow{\mu_{f1_{df}}} & F(f \circ 1_{df}) \\ \alpha * 1_{Ff} \downarrow & & \parallel & 1_{Ff} * \alpha \downarrow & & \parallel \\ 1_{F(cf)} \circ Ff & \xlongequal{\quad} & Ff & Ff \circ 1_{F(df)} & \xlongequal{\quad} & Ff \end{array}$$

- iv) if $\sigma : f \rightarrow f', \tau : g \rightarrow g'$ are 2-morphisms such that $d^2\tau = c^2\sigma$ then the following diagram is commutative:

$$\begin{array}{ccc} Fg \circ Ff & \xrightarrow{\mu_{gf}} & F(g \circ f) \\ F\tau * F\sigma \downarrow & & \downarrow F(\tau * \sigma) \\ Fg' \circ Ff' & \xrightarrow{\mu_{g'f'}} & F(g' \circ f') \end{array}$$

Example 1.1.3.9. We have

$$\text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{Cat}}_{ad}, \mathcal{C} \mapsto \text{Ho } \mathcal{C}, (F, G, \varphi) \mapsto \text{Ho}(F, G, \varphi), \tau \mapsto L\tau$$

is a pseudo-2-functor. Moreover $\text{Ho} \circ (-^{\text{op}}) = (-^{\text{op}}) \circ \text{Ho}$.

Definition 1.1.3.10. Suppose \mathbb{C}, \mathbb{D} are 2-categories, $F, G : \mathbb{C} \rightarrow \mathbb{D}$ are pseudo-2-functors. A **2-natural transformation** $\tau : F \rightarrow G$ consists of a map $\tau_X : FX \rightarrow GX$ for any $X \in \mathbb{C}_0$, such that for any $f \in \mathbb{C}_1$, $Gf \circ \tau_{df} = \tau_{cf} \circ Ff$, and for any $\alpha \in \mathbb{C}_2$, $G\alpha * 1_{\tau_{d^2\alpha}} = 1_{\tau_{c^2\alpha}} * F\alpha$. A **2-natural isomorphism** is a 2-natural transformation τ such that τ_X is an isomorphism for any $X \in \mathbb{C}_0$.

1.2 Examples of Model Categories

1.2.1 Cofibrantly Generated Model Categories

Definition 1.2.1.1. Suppose \mathcal{C} is a category with all small colimits and λ is an ordinal. A λ -sequence X is a colimit-preserving functor $\lambda \rightarrow \mathcal{C}$. We denote $X(\alpha)$ as X_α . A **transfinite composition** of a λ -sequence is the induced map $X_0 \rightarrow \operatorname{colim}_{\beta < \lambda} X_\beta$. If I is a collection of maps of \mathcal{C} and for any $\beta + 1 < \lambda$ the map $X_\beta \rightarrow X_{\beta+1}$ is in I , we call the λ -sequence to be in I and transfinite composition a transfinite composition of maps of I .

Definition 1.2.1.2. Suppose γ is a cardinal. A γ -**filtered ordinal** is a limit ordinal α such that for any $A \subseteq \alpha, |A| \leq \gamma$ we have $\sup A < \alpha$.

Definition 1.2.1.3. Suppose \mathcal{C} is a category with all small colimits, I is a collection of maps of \mathcal{C} , A is an object in \mathcal{C} and κ is a cardinal. We say A is κ -**small relative to I** if for any κ -filtered ordinal λ and all λ -sequence X in I , the map of sets

$$\operatorname{colim}_{\beta < \lambda} \operatorname{Hom}_{\mathcal{C}}(A, X_\beta) \rightarrow \operatorname{Hom}_{\mathcal{C}}(A, \operatorname{colim}_{\beta < \lambda} X_\beta)$$

is an isomorphism. We say A is **small relative to I** if there exists some κ such that A is κ -small relative to I . We say A is **small** if A is small relative to \mathcal{C} . We say A is **finite relative to I** if there exists some finite cardinal κ such that A is κ -small relative to I . We say A is **finite** if A is finite relative to \mathcal{C} .

Example 1.2.1.4. In the category Set every object A is $|A|$ -small.

Example 1.2.1.5. If R is a ring, in the category Mod $_R$ of left R -modules every object is small, every finitely presented R -module is finite.

Remark 1.2.1.6. In the category Top not all objects are small; For example, the indiscrete space with 2 points.

Definition 1.2.1.7. Suppose \mathcal{C} is a category, I is a collection of maps of \mathcal{C} . Denote $\operatorname{RLP}(I)$ to be the class of all maps that has RLP with respect to every map in I , and $\operatorname{LLP}(I)$ to be the class of maps that has LLP with respect to every map in I . Denote $\operatorname{Cof}(I) = \operatorname{LLP}(\operatorname{RLP}(I))$ and $\operatorname{Fib}(I) = \operatorname{RLP}(\operatorname{LLP}(I))$. We call the maps in $\operatorname{Cof}(I)$ to be **I -cofibrations** and $\operatorname{Fib}(I)$ to be **I -fibrations**.

Lemma 1.2.1.8. Suppose \mathcal{C} is a category, I is a collection of maps of \mathcal{C} . Then $I \subseteq \operatorname{Cof}(I), I \subseteq \operatorname{Fib}(I)$. Moreover $\operatorname{RLP}(\operatorname{Cof}(I)) = \operatorname{RLP}(I)$ and $\operatorname{LLP}(\operatorname{Fib}(I)) = \operatorname{LLP}(I)$.

Lemma 1.2.1.9. Suppose (F, G) is an adjunction between \mathcal{C} and \mathcal{D} , I is a collection of maps of \mathcal{C} , J is a collection of maps of \mathcal{D} . Then we have $G(\operatorname{RLP}(FI)) \subseteq \operatorname{RLP}(I), F(\operatorname{Cof}(I)) \subseteq \operatorname{Cof}(FJ), F(\operatorname{LLP}(GJ)) \subseteq \operatorname{LLP}(J), G(\operatorname{Fib}(J)) \subseteq \operatorname{Fib}(GJ)$.

Definition 1.2.1.10. Suppose \mathcal{C} is a category with all small colimits and I is a collection of maps of \mathcal{C} . A **relative I -cell complex** is a transfinite composition of pushouts of elements of I . Denote $\operatorname{Cell}(I)$ to be the class of all relative I -cell complexes. An object A is called an **I -cell complex** if the map $\phi \rightarrow A$ is a relative I -cell complex.

Lemma 1.2.1.11. Suppose \mathcal{C} is a category with all small colimits and I is a collection of maps of \mathcal{C} . Then $\operatorname{Cell}(I) \subseteq \operatorname{Cof}(I)$.

Lemma 1.2.1.12. Suppose \mathcal{C} is a category with all small colimits and I is a collection of maps of \mathcal{C} . Then transfinite composition of pushouts of elements of I or isomorphisms is in $\operatorname{Cell}(I)$.

Lemma 1.2.1.13. *Suppose \mathcal{C} is a category with all small colimits and I is a collection of maps of \mathcal{C} . Then $\text{Cell}(I)$ is closed under transfinite composition.*

Lemma 1.2.1.14. *Suppose \mathcal{C} is a category with all small colimits and I is a collection of maps of \mathcal{C} . Then any pushout of coproducts of maps of I is in $\text{Cell}(I)$.*

Theorem 1.2.1.15 (The Small Object Argument). *Suppose \mathcal{C} is a category with all small colimits and I is a set of maps of \mathcal{C} . Suppose the domains of maps of I are small relative to $\text{Cell}(I)$. Then there exists a functorial factorization (γ, δ) on \mathcal{C} such that for any f a morphism in \mathcal{C} , $\gamma(f) \in \text{Cell}(I)$ and $\delta(f) \in \text{RLP}(I)$.*

Corollary 1.2.1.16. *Suppose \mathcal{C} is a category with all small colimits and I is a set of maps of \mathcal{C} . Suppose the domains of maps of I are small relative to $\text{Cell}(I)$. Then for any $f : A \rightarrow B \in \text{Cof}(I)$ there exists $g : A \rightarrow C \in \text{Cell}(I)$ such that f is a retract of g , and the retract fixes A .*

Proposition 1.2.1.17. *Suppose \mathcal{C} is a category with all small colimits and I is a set of maps of \mathcal{C} . Suppose the domains of maps of I are small relative to $\text{Cell}(I)$. Then any object that is small relative to $\text{Cell}(I)$ is small relative to $\text{Cof}(I)$.*

Definition 1.2.1.18. Suppose \mathcal{C} is a model category. \mathcal{C} is called **cofibrantly generated** if there exist sets of maps I and J of \mathcal{C} , such that:

- i) The domains of maps of I are small relative to $\text{Cell}(I)$;
- ii) The domains of maps of J are small relative to $\text{Cell}(J)$;
- iii) $\text{Fib} = \text{RLP}(J)$;
- iv) $\text{W} \cap \text{Fib} = \text{RLP}(I)$.

The sets I and J are called the **generating cofibrations** and the **generating trivial cofibrations**, respectively. A cofibrantly generated model category is called **finitely generated** if the domains and codomains of maps of I and J are finite relative to $\text{Cell}(I)$.

Proposition 1.2.1.19. *Suppose \mathcal{C} is a cofibrantly generated model category with generating cofibrations I and generating trivial cofibrations J . Then:*

- i) $\text{Cof} = \text{Cof}(I)$, $\text{W} \cap \text{Cof} = \text{Cof}(J)$;
- ii) Any cofibration is a retract of a relative I -cell complex; any trivial cofibration is a retract of a relative J -cell complex;
- iii) The domains of maps of I are small relative to Cof ; the domains of maps of J are small relative to $\text{W} \cap \text{Cof}$.

If \mathcal{C} is finitely generated then the domains and codomains of maps of I and J are finite relative to Cof .

Theorem 1.2.1.20. *Suppose \mathcal{C} is a category with all small limits and colimits, W is a class of maps of \mathcal{C} , I and J are sets of maps in \mathcal{C} . Then \mathcal{C} is a cofibrantly generated model category with weak equivalence W , generating cofibrations I and generating trivial cofibrations J if and only if the following are satisfied:*

- i) W has the 2-out-of-3 property and is closed under retracts;
- ii) The domains of maps of I are small relative to $\text{Cell}(I)$;

- iii) The domains of maps of J are small relative to $\text{Cell}(J)$;
- iv) $\text{Cell}(J) \subseteq W \cap \text{Cof}(I)$;
- v) $\text{RLP}(I) \subseteq W \cap \text{RLP}(J)$;
- vi) Either $W \cap \text{Cof}(I) \subseteq \text{Cof}(J)$ or $W \cap \text{RLP}(J) \subseteq \text{RLP}(I)$.

Proposition 1.2.1.21. Suppose $(F, G) : \mathcal{C} \rightarrow \mathcal{D}$ is an adjunction, where \mathcal{C} and \mathcal{D} are model categories. Suppose more that \mathcal{C} is a cofibrantly generated model category with generating cofibrations I and generating trivial cofibrations J . Then (F, G) is an Quillen adjunction if and only if $FI \subseteq \text{Cof}_{\mathcal{D}}$ and $FJ \subseteq W_{\mathcal{D}} \cap \text{Cof}_{\mathcal{D}}$.

Proposition 1.2.1.22. If \mathcal{C} is a cofibrantly generated model category then so is \mathcal{C}_* . If \mathcal{C} is a finitely generated model category then so is \mathcal{C}_* .

1.2.2 The Category of Modules over a Frobenius Ring

Definition 1.2.2.1. A ring R is called a **Frobenius ring** if the projective and injective R -modules coincide.

Definition 1.2.2.2. Suppose R is a ring. Two maps $f, g : M \rightarrow N$ of R -modules are called **stably equivalent**, denote $f \simeq g$, if $f - g$ factors through a projective module.

Lemma 1.2.2.3. Stable equivalence between maps is an equivalence relation that is compatible with composition.

Definition 1.2.2.4. Two R -modules M, N are called **stably equivalent** if there exists maps $f : M \rightarrow N$, $g : N \rightarrow M$ such that $fg \simeq \mathbb{1}_N$, $gf \simeq \mathbb{1}_M$. Such f, g are called **stable equivalences**.

Definition 1.2.2.5. Suppose R is a ring. Define the set $I = \{A \rightarrow R \mid A \text{ is an ideal of } R\}$, and the set $J = \{0 \rightarrow R\}$. Define $\text{Cof} = \text{Cof}(I)$, $\text{Fib} = \text{RLP}(J)$, and W to be all stable equivalences.

Lemma 1.2.2.6. A map $p \in \underline{\text{Mod}}_R$ is a fibration if and only if it is surjective.

Lemma 1.2.2.7. Suppose R is a Frobenius ring. Then a map $p \in \underline{\text{Mod}}_R$ is a trivial fibration if and only if it is surjective with injective kernel.

Proposition 1.2.2.8. A map $p \in \underline{\text{Mod}}_R$ is in $\text{RLP}(I)$ if and only if it is surjective with injective kernel. In particular, if R is a Frobenius ring, then $W \cap \text{Fib} = \text{RLP}(I)$.

Lemma 1.2.2.9. A map $i \in \underline{\text{Mod}}_R$ is a cofibration if and only if it is injective.

Proposition 1.2.2.10. A map $i \in \underline{\text{Mod}}_R$ is in $\text{Cof}(J)$ if and only if it is injective with projective cokernel. In particular, $\text{Cof}(J) \subseteq W \cap \text{Cof}$.

Theorem 1.2.2.11. If R is a Frobenius ring, then $\underline{\text{Mod}}_R$ is a cofibrantly generated model category with I being its generating cofibrations, J being its generating trivial cofibrations and weak equivalences being stable equivalences. If R is Noetherian then $\underline{\text{Mod}}_R$ is finitely generated.

Remark 1.2.2.12. If $\underline{\text{Mod}}_R$ is a cofibrantly generated model category with I being its generating cofibrations and J being its generating trivial cofibrations, then R must be a Frobenius ring and weak equivalences must be stable equivalences.

Corollary 1.2.2.13. If R is a Frobenius ring, every object in $\underline{\text{Mod}}_R$ is cofibrant and fibrant, and two maps are left or right homotopic if and only if they are stably equivalent.

Proposition 1.2.2.14. For any ring homomorphism $R \rightarrow S$ between Frobenius rings, the adjunction $(\text{Ind}, \text{Res}) : \underline{\text{Mod}}_R \rightarrow \underline{\text{Mod}}_S$ is a Quillen adjunction if and only if S is a flat R -module, it is a Quillen equivalence if and only if the homomorphism is an isomorphism.

1.2.3 The Category of Chain Complexes of Modules over a Ring

Lemma 1.2.3.1. *If R is a ring, all objects in the category $\underline{\mathbf{Ch}}_R$ of chain complexes of R -modules are small. All bounded complexes of finitely presented R -modules are finite.*

Definition 1.2.3.2. Suppose R is a ring, define the functor $S^n : \underline{\mathbf{Mod}}_R \rightarrow \underline{\mathbf{Ch}}_R$ with $S^n(M)_n = M$ and $S^n(M)_m = 0$ if $n \neq m$; define the functor $D^n : \underline{\mathbf{Mod}}_R \rightarrow \underline{\mathbf{Ch}}_R$ with $D^n(M)_n = D^n(M)_{n-1} = M$ and $D^n(M)_m = 0$ if $n \neq m, m-1$, and $d_n = 1_M$. Denote $S^n(R)$ by S^n and $D^n(R)$ by D^n .

Lemma 1.2.3.3. *S^n is left adjoint to $Z_n : X \rightarrow \ker d_n$, and D^n is left adjoint to $-_n$.*

Definition 1.2.3.4. Suppose R is a ring. Define the set $I = \{S^{n-1} \rightarrow D^n | n \in \mathbb{Z}\}$, and the set $J = \{0 \rightarrow D^n | n \in \mathbb{Z}\}$. Define $\text{Cof} = \text{Cof}(I)$, $\text{Fib} = \text{RLP}(J)$, and W to be all quasi-isomorphisms.

Lemma 1.2.3.5. *A map $p \in \underline{\mathbf{Ch}}_R$ is a fibration if and only if it is surjective.*

Proposition 1.2.3.6. *A map $p \in \underline{\mathbf{Ch}}_R$ is in $\text{RLP}(I)$ if and only if it is a surjective quasi-isomorphism. In particular, $\text{RLP}(I) = W \cap \text{Fib}$.*

Lemma 1.2.3.7. *Any cofibrant chain complex is degreewise projective. Any bounded below degreewise projective chain complex is cofibrant.*

Remark 1.2.3.8. Not every degreewise projective chain complex is cofibrant. For example take $R = \mathbb{k}[x]/(x^2)$ where \mathbb{k} is a field, and take A to be the complex with $A_n = R$, and d_n to be the multiplication by x for any n , then A is degreewise projective but not cofibrant.

Lemma 1.2.3.9. *If C is a cofibrant chain complex and K is acyclic, then any map from C to K is nullhomotopic.*

Proposition 1.2.3.10. *A map $i \in \underline{\mathbf{Ch}}_R$ is a cofibration if and only if it is injective with cofibrant cokernel.*

Proposition 1.2.3.11. *A map $i \in \underline{\mathbf{Ch}}_R$ is in $\text{Cof}(J)$ if and only if it is injective with projective cokernel. In particular, $\text{Cof}(J) \subseteq W \cap \text{Cof}$.*

Theorem 1.2.3.12. *$\underline{\mathbf{Ch}}_R$ is a cofibrantly generated model category with I being its generating cofibrations, J being its generating trivial cofibrations and weak equivalences being quasi-isomorphisms. This model structure is called the **projective model structure**.*

Corollary 1.2.3.13. *Any object in $\underline{\mathbf{Ch}}_R$ is projectively fibrant. An object in $\underline{\mathbf{Ch}}_R$ is projective if and only if it is acyclic and cofibrant. If two maps in $\underline{\mathbf{Ch}}_R$ are chain homotopic then they are right homotopic.*

Definition 1.2.3.14. Define Cof' to be all injections in $\underline{\mathbf{Ch}}_R$. Define Fib' to be $\text{RLP}(W \cap \text{Cof}')$, and call elements in Fib' **injective fibrations**. For any $X \in \underline{\mathbf{Ch}}_R$ define $|X| = |\bigcup_n X_n|$. Define $\gamma = \sup\{|R|, |\mathbb{Z}|\}$. Define I' to be the set containing a map from each isomorphism class of injections whose codomain has cardinality no larger than γ . Define $J' = I' \cap W$.

Proposition 1.2.3.15. *$\text{Cof}(I') = \text{Cof}'$. Moreover a map $p \in \underline{\mathbf{Ch}}_R$ is in $\text{RLP}(I')$ if and only if it is surjective with injective kernel.*

Corollary 1.2.3.16. *Any injective object in $\underline{\mathbf{Ch}}_R$ is injectively fibrant and acyclic. Moreover $\text{RLP}(I') \subseteq W \cap \text{Fib}'$.*

Lemma 1.2.3.17. *Any injectively fibrant chain complex is degreewise injective. Any bounded above degreewise injective chain complex is injectively fibrant.*

Remark 1.2.3.18. Not every degreewise projective chain complex is injectively fibrant. For example take $R = \mathbb{k}[x]/(x^2)$ where \mathbb{k} is a field, and take A to be the complex with $A_n = R$, and d_n to be the multiplication by x for any n , then A is degreewise injective but not injectively fibrant.

Lemma 1.2.3.19. *If C is an acyclic chain complex and K is injectively fibrant, then any map from C to K is nullhomotopic.*

Proposition 1.2.3.20. *A map $p \in \underline{\mathbf{Ch}}_R$ is in Fib' if and only if it is surjective with injectively fibrant kernel.*

Lemma 1.2.3.21. *Suppose $i : A \rightarrow B$ is a map in $\underline{\mathbf{Ch}}_R$ such that $i \in \text{Cof}' \cap \text{W}$. If C is a subcomplex of B with $|C| \leq \gamma$, then there exists a subcomplex D of B containing C such that $i : D \cap A \rightarrow D$ is in W .*

Proposition 1.2.3.22. $\text{Cof}(J') = \text{Cof}' \cap \text{W}$. $\text{RLP}(J') = \text{Fib}'$.

Theorem 1.2.3.23. $\underline{\mathbf{Ch}}_R$ is a cofibrantly generated model category with I' being its generating cofibrations, J' being its generating trivial cofibrations and weak equivalences being quasi-isomorphisms. This model structure is called the **injective model structure**.

Corollary 1.2.3.24. *Any object in $\underline{\mathbf{Ch}}_R$ is injectively cofibrant. An object in $\underline{\mathbf{Ch}}_R$ is injective if and only if it is acyclic and injectively fibrant.*

Proposition 1.2.3.25. *The identity functor from the projective model structure to the injective model structure is a Quillen equivalence.*

Proposition 1.2.3.26. *For any ring homomorphism $R \rightarrow S$, the adjunction $(\text{Ind}, \text{Res}) : \underline{\mathbf{Ch}}_R \rightarrow \underline{\mathbf{Ch}}_S$ is a Quillen adjunction between the projective model structures, it is a Quillen equivalence if and only if the homomorphism is an isomorphism.*

Proposition 1.2.3.27. *For any ring homomorphism $R \rightarrow S$, the adjunction $(\text{Ind}, \text{Res}) : \underline{\mathbf{Ch}}_R \rightarrow \underline{\mathbf{Ch}}_S$ is a Quillen adjunction between the injective model structures if and only if S is a flat R -module, it is a Quillen equivalence if and only if the homomorphism is an isomorphism.*

Proposition 1.2.3.28. $[S^n M, S^0 N] \cong \text{Ext}_R^n(M, N)$ for any R -module M, N .

1.2.4 The Category of Topological Spaces

Lemma 1.2.4.1. *Every topological space is small relative to the inclusions.*

Definition 1.2.4.2. Define an inclusion to be a **closed T_1 inclusion** if it is closed and every point not in the image is closed.

Proposition 1.2.4.3. *Every compact space is finite relative to closed T_1 inclusions.*

Definition 1.2.4.4. Define the set $I = \{S^{n-1} \rightarrow D^n \mid n \geq 0\}$ ($S^{-1} = \emptyset, D^0 = *$), and the set $J = \{D^n \rightarrow D^n \times [0, 1] \mid n \geq 0\}$. Define $\text{Cof} = \text{Cof}(I)$, $\text{Fib} = \text{RLP}(J)$, and W to be all weak homotopy equivalences. Define a map to be a **Serre fibration** if and only if it is in Fib . Define a map to be a **relative cell complex** if and only if it is in $\text{Cell}(I)$.

Lemma 1.2.4.5. *Every relative cell complex is a closed T_1 inclusion.*

Corollary 1.2.4.6. *Every map in Cof or $\text{Cof}(J)$ is a closed T_1 inclusion.*

Lemma 1.2.4.7. *Suppose X is a relative cell complex. Then every compact set of X intersects the interiors of only finitely many cells.*

Lemma 1.2.4.8. *Closed T_1 inclusions that are also weak equivalence are closed under transfinite composition.*

Proposition 1.2.4.9. $\text{Cof}(J) \subseteq W \cap \text{Cof}$.

Proposition 1.2.4.10. $\text{RLP}(I) \subseteq W \cap \text{Fib}$.

Lemma 1.2.4.11. *A map $p : X \rightarrow Y$ is in $\text{RLP}(I)$ if and only for any map $i : A \rightarrow B \in I$, $\mathcal{H}om_{\square}(i, p) : \mathcal{H}om(B, X) \rightarrow \mathcal{H}om(A, X) \times_{\mathcal{H}om(A, Y)} \mathcal{H}om(B, Y)$ is surjective. In particular, if for any map $i \in I$, $\mathcal{H}om_{\square}(i, p)$ is a trivial fibration, then $p \in \text{RLP}(I)$.*

Lemma 1.2.4.12. *If p is a fibration and $i \in I$, then $\mathcal{H}om_{\square}(i, p)$ is a fibration.*

Corollary 1.2.4.13. *Every topological space is fibrant, and for any $n \geq 0$ and Y a space the map $\mathcal{H}om(D^n, Y) \rightarrow \mathcal{H}om(S^{n-1}, Y)$ is a fibration.*

Lemma 1.2.4.14. *If p is a weak equivalence, then for any $n \geq 0$, $\mathcal{H}om(D^n, p)$ is a weak equivalence.*

Lemma 1.2.4.15. *If p is a weak equivalence, then for any $n \geq -1$, $\mathcal{H}om(S^n, p)$ is a weak equivalence.*

Proposition 1.2.4.16. *The pullback of any weak equivalence along a fibration is a weak equivalence.*

Theorem 1.2.4.17. $W \cap \text{Fib} \subseteq \text{RLP}(I)$.

Theorem 1.2.4.18. Top is a finitelyly generated model category with I being its generating cofibrations, J being its generating trivial cofibrations and weak equivalences being weak homotopy equivalences. As a corollary, Top_{*} is a finitelyly generated model category with I_+ being its generating cofibrations, J_+ being its generating trivial cofibrations and weak equivalences being weak homotopy equivalences.

Definition 1.2.4.19. Suppose X is a space. A subset U of X is called **compactly open** if for any map $f : K \rightarrow X$ with K compact Hausdorff, $f^{-1}(U)$ is open. X is called a **k -space** if all its compactly open sets are open. Define \underline{K} to be the full subcategory of Top containing all k -spaces. Define the **k -space topology** on X , denoted kX , to be defining U open in kX if and only if it is compactly open in X .

Proposition 1.2.4.20. k is a functor from Top to \underline{K} , and it has a left adjoint and right inverse i , the inclusion.

Proposition 1.2.4.21. \underline{K} has all small colimits and limits, where colimits are directly taken in Top and limits are taken by applying k to the limit in Top.

Proposition 1.2.4.22. For any $X, Y \in \underline{K}$, define $C(X, Y)$ to be the space with underlying set $\text{Hom}_{\underline{K}}(X, Y)$, with the topological subbasis

$$\{\{g : X \rightarrow Y \mid g(f(K)) \subseteq U\} \mid f : K \rightarrow X, \\ K \text{ compact, Hausdorff, } U \text{ an open set of } Y\}.$$

Define $\mathcal{H}om(X, Y) = kC(X, Y)$, then we have a natural isomorphism

$$\text{Hom}_{\underline{K}}(k(X \times Y), Z) \cong \text{Hom}_{\underline{K}}(X, \mathcal{H}om(Y, Z))$$

for any $X, Y, Z \in \underline{K}$. ¹

¹The usage of $\text{Hom}, \mathcal{H}om, \text{Map}, -^-$ are as follows. Suppose we want to say that the Hom-object of A and B is C . If C is a set, or C is a simplicial set, or we are talking about Hom-object in enriched categories, we use Hom . If in addition C is a simplicial set, we may add \bullet to distinguish it with the case that C is a set. If C is a homotopy type, we use Map . If B, C are of the same type, and a) A is of different type, or b) B, C are sets, simplicial sets or homotopy types, we use the exponential $-^-$. In other cases, we use $\mathcal{H}om$.

Theorem 1.2.4.23. \underline{K} is a finitely generated model category with I being its generating cofibrations, J being its generating trivial cofibrations and weak equivalences being weak homotopy equivalences. Moreover we have $\text{Cof}_{\underline{K}} = \text{Cof}_{\underline{\text{Top}}} \cap \underline{K}$, $\text{Fib}_{\underline{K}} = \text{Fib}_{\underline{\text{Top}}} \cap \underline{K}$. As a corollary, \underline{K}_* is a finitely generated model category with I_+ being its generating cofibrations, J_+ being its generating trivial cofibrations and weak equivalences being weak homotopy equivalences. Moreover we have $\text{Cof}_{\underline{K}_*} = \text{Cof}_{\underline{\text{Top}}_*} \cap \underline{K}_*$, $\text{Fib}_{\underline{K}_*} = \text{Fib}_{\underline{\text{Top}}_*} \cap \underline{K}_*$. Furthermore the inclusions $\underline{K} \rightarrow \underline{\text{Top}}$, $\underline{K}_* \rightarrow \underline{\text{Top}}_*$ are Quillen equivalences.

1.3 Simplicial Sets

1.3.1 The Definition of Simplicial Sets

Definition 1.3.1.1. Define the **simplicial category** Δ to be the category with objects $\{[n] := \{0, 1, \dots, n\} \mid n \in \mathbb{N}\}$ and

$$\text{Hom}_{\Delta}([k], [n]) := \{f : [k] \rightarrow [n] \mid \forall 0 \leq x \leq y \leq k, f(x) \leq f(y)\}.$$

Definition 1.3.1.2. Define Δ_+ to be the subcategory of Δ with

$$\text{Hom}_{\Delta_+}([k], [n]) := \{f \in \text{Hom}_{\Delta}([k], [n]) \mid f \text{ injective}\}.$$

Define Δ_- to be the subcategory of Δ with

$$\text{Hom}_{\Delta_-}([k], [n]) := \{f \in \text{Hom}_{\Delta}([k], [n]) \mid f \text{ surjective}\}.$$

Definition 1.3.1.3. For any $n \geq 1, 0 \leq i \leq n$, define

$$d^i : [n-1] \rightarrow [n], d^i(j) = \begin{cases} j & (j < i) \\ j+1 & (j \geq i) \end{cases};$$

For any $n \geq 1, 0 \leq i \leq n-1$, define

$$s^i : [n] \rightarrow [n-1], s^i(j) = \begin{cases} j & (j \leq i) \\ j-1 & (j > i) \end{cases}.$$

Lemma 1.3.1.4. We have the **cosimplicial identities**:

$$\begin{aligned} d^j d^i &= d^i d^{j-1} & (i < j), \\ s^j d^i &= d^i s^{j-1} & (i < j), \\ &= \mathbb{1} & (i = j, j+1), \\ &= d^{i-1} s^j & (i > j+1), \\ s^j s^i &= s^{i-1} s^j & (i > j). \end{aligned}$$

Definition 1.3.1.5. If \mathcal{C} is a category, define the category of **cosimplicial objects of \mathcal{C}** to be \mathcal{C}^{Δ} , and the category of **simplicial objects of \mathcal{C}** to be $\mathcal{C}^{\Delta^{\text{op}}}$.

Definition 1.3.1.6. Define the category of **simplicial sets** to be $\underline{\text{SSet}} := \underline{\text{Set}}^{\Delta^{\text{op}}}$.

Definition 1.3.1.7. If K is a simplicial set, denote the set of **n -simplices** to be $K_n := K[n]$. For any $n \geq 1, 0 \leq i \leq n$, define the **face maps** to be $d_i := K(d^i) : K_n \rightarrow K_{n-1}$. For any $n \geq 1, 0 \leq i \leq n-1$, define the **degeneracy maps** to be $s_i := K(s^i) : K_{n-1} \rightarrow K_n$.

Lemma 1.3.1.8. *For any simplicial set we have the **simplicial identities**:*

$$\begin{aligned} d_i d_j &= d_{j-1} d_i & (i < j), \\ d_i s_j &= s_{j-1} d_i & (i < j), \\ &= \mathbb{1} & (i = j, j+1), \\ &= s_j d_{i-1} & (i > j+1), \\ s_i s_j &= s_j s_{i-1} & (i > j); \end{aligned}$$

Conversely, any maps satisfying the simplicial identities determine a simplicial set.

Definition 1.3.1.9. Suppose x, y are simplices in a simplicial set K . y is called a **face** of x if y is the image of x under a composition of some face maps, y is called a **degeneracy** of x if y is the image of x under a composition of some degeneracy maps. x is called **nondegenerate** if x is a degeneracy only of x . K is called **finite** if the number of nondegenerate elements is finite.

Lemma 1.3.1.10. *If $K \in \underline{\mathbf{SSet}}$, then $\forall x \in K, \exists ! y \in K$, such that x is a degeneracy of y .*

Lemma 1.3.1.11. *Every simplicial set is small. Every finite simplicial set is finite.*

Definition 1.3.1.12. We may define the functor $\Delta[-] : \Delta \rightarrow \underline{\mathbf{SSet}}$, $[n] \mapsto \text{Hom}_\Delta(-, [n])$, and we call $\Delta[n]$ the **standard n -dimensional simplicial set**. We denote $i_n = \mathbb{1}_{[n]} \in \Delta[n]_n$.

Remark 1.3.1.13. The above definition generalizes to the functor $\Delta[-] : \underline{\mathbf{POSet}} \rightarrow \underline{\mathbf{SSet}}$, $P \mapsto \text{Hom}_{\underline{\mathbf{POSet}}}(-, P)$, via the inclusion $\Delta \rightarrow \underline{\mathbf{POSet}}$.

Lemma 1.3.1.14. *All nondegenerate k -simplices of $\Delta[n]$ are elements in $\text{Hom}_{\Delta_+}([k], [n])$. In particular, i_n is the only nondegenerate n -simplex of $\Delta[n]$.*

Lemma 1.3.1.15. *We have a natural isomorphism between the functors*

$$\text{Hom}_{\underline{\mathbf{SSet}}}(\Delta[n], -) \cong -_n, f \mapsto f(i_n) : \underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Set}}.$$

Definition 1.3.1.16. We define the **boundary** of $\Delta[n]$, $\partial\Delta[n]$ to be the simplicial set obtained from $\Delta[n]$ by removing all degeneracies of i_n .

Definition 1.3.1.17. We define the **r -horn** of $\Delta[n]$, $\Lambda^r[n]$ to be the simplicial set obtained from $\partial\Delta[n]$ by removing all degeneracies of $d^r \in \partial\Delta[n]_{n-1}$.

Definition 1.3.1.18. There is a functor $\Delta : \underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}$ with

$$\begin{aligned} K \mapsto & \left(\begin{array}{l} \text{Objects: } \{\Delta[n] \rightarrow K\} \\ \text{Hom}_{\Delta K}((f : \Delta[k] \rightarrow K), (g : \Delta[n] \rightarrow K)) \\ \quad = \{h \in \text{Hom}_\Delta([k], [n]) \mid g \circ \Delta[-](h) = f\} \end{array} \right), \\ (a : K \rightarrow L) \mapsto & ((f : \Delta[n] \rightarrow K) \mapsto a \circ f). \end{aligned}$$

For any simplicial set K , denote ΔK the **category of simplices** in K .

Lemma 1.3.1.19. *We have*

$$\text{colim} \left(\begin{array}{l} \Delta K \rightarrow \underline{\mathbf{SSet}} \\ (f : \Delta[n] \rightarrow K) \mapsto \Delta[n] \end{array} \right) = K.$$

Definition 1.3.1.20. There is a functor $\Delta' : \underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}$ with

$$\begin{aligned} K \mapsto & \left(\begin{array}{l} \text{Objects: } \{f : \Delta[n] \rightarrow K \mid f(i_n) \text{ nondegenerate}\} \\ \text{Hom}_{\Delta' K}((f : \Delta[k] \rightarrow K), (g : \Delta[n] \rightarrow K)) \\ \quad = \{h \in \text{Hom}_{\Delta_+}([k], [n]) \mid g \circ \Delta[-](h) = f\} \end{array} \right), \\ (a : K \rightarrow L) \mapsto & ((f : \Delta[n] \rightarrow K) \mapsto a \circ f). \end{aligned}$$

For any simplicial set K , denote $\Delta' K$ the **category of nondegenerate simplices** in K .

Lemma 1.3.1.21. *If K is **regular**, which means for any nondegenerate n -simplex the map $\Delta[n] \rightarrow K$ induced by the n -simplex is injective, then we have*

$$\operatorname{colim} \left(\begin{array}{c} \Delta' K \rightarrow \underline{\mathbf{SSet}} \\ (f : \Delta[n] \rightarrow K) \mapsto \Delta[n] \end{array} \right) = K.$$

Remark 1.3.1.22. If K is not regular then Lemma 1.3.1.21 may not be true. A counterexample is $\Delta[2]/\partial\Delta[2]$.

Proposition 1.3.1.23. *If \mathcal{C} is a category with all small colimits, then the functor $\operatorname{Adj}(\underline{\mathbf{SSet}}, \mathcal{C}) \rightarrow \mathcal{C}^\Delta$ with $(F, G, \varphi) \mapsto F \circ \Delta[-]$ is an equivalence between categories with inverse being $A^\bullet \mapsto (A^\bullet \otimes -, \operatorname{Hom}_{\mathcal{C}}(A^\bullet, -), \varphi)$, where*

$$A^\bullet \otimes K = \operatorname{colim} \left(\begin{array}{c} \Delta K \rightarrow \mathcal{C} \\ (f : \Delta[n] \rightarrow K) \mapsto A^n \end{array} \right)$$

and

$$\operatorname{Hom}_{\mathcal{C}}(A^\bullet, Y)_n = \operatorname{Hom}_{\mathcal{C}}(A^n, Y).$$

Proposition 1.3.1.24. *If \mathcal{C} is a category with all small limits, then the category $\operatorname{Adj}(\underline{\mathbf{SSet}}, \mathcal{C}^{\operatorname{op}})^{\operatorname{op}}$ is naturally equivalent to $\mathcal{C}^{\Delta^{\operatorname{op}}}$. We denote the image of $A_\bullet \in \mathcal{C}^{\Delta^{\operatorname{op}}}$ under the equivalence to be $(A_\bullet^-, \operatorname{Hom}_{\mathcal{C}}(-, A_\bullet), \varphi)$.*

Corollary 1.3.1.25. *If \mathcal{C} is a pointed category with all small colimits, then the category $\operatorname{Adj}(\underline{\mathbf{SSet}}_*, \mathcal{C})$ is naturally equivalent to \mathcal{C}^Δ . We denote the image of $A^\bullet \in \mathcal{C}^\Delta$ under the equivalence to be $(A^\bullet \wedge -, \operatorname{Hom}_{\mathcal{C}_*}(A^\bullet, -), \varphi)$. Furthermore we have a natural isomorphism $A^\bullet \wedge (-_+) \cong A^\bullet \otimes -$. Dually the category $\operatorname{Adj}(\underline{\mathbf{SSet}}_*, \mathcal{C}^{\operatorname{op}})^{\operatorname{op}}$ is naturally equivalent to $\mathcal{C}^{\Delta^{\operatorname{op}}}$. We denote the image of $A_\bullet \in \mathcal{C}^{\Delta^{\operatorname{op}}}$ under the equivalence to be $(A_{\bullet*}^-, \operatorname{Hom}_{\mathcal{C}_*}(-, A_\bullet), \varphi)$. Furthermore we have a natural isomorphism $A_{\bullet*}^{-+} \cong A_\bullet^-$.*

Example 1.3.1.26. By Proposition 1.3.1.23 the functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{SSet}}^\Delta$, $K \mapsto K \times \Delta[-]$ corresponds to a functor $\underline{\mathbf{SSet}} \rightarrow \operatorname{Adj}(\underline{\mathbf{SSet}}, \underline{\mathbf{SSet}})$ which we denote by $K \mapsto (K \times -, -^K, \varphi)$, where $(L^K)_n = \operatorname{Hom}_{\underline{\mathbf{SSet}}}(K \times \Delta[n], L)$. The left adjoint is exactly the product in $\underline{\mathbf{SSet}}$. We call $-^K$ the **function complex functor**.

Example 1.3.1.27. By Proposition 1.3.1.23 the element

$$([n] \mapsto \{(t_0, \dots, t_n) | t_i \geq 0, t_0 + \dots + t_n = 1\}) \in \underline{\mathbf{Top}}^\Delta(\underline{\mathbf{K}}^\Delta)$$

corresponds to an element in $\operatorname{Adj}(\underline{\mathbf{SSet}}, \underline{\mathbf{Top}}(\underline{\mathbf{K}}))$. We denote the adjunction by $(|-|, \operatorname{Sing}, \varphi)$ and call $|-|$ the **geometric realization** and Sing the **singular functor**.

Lemma 1.3.1.28. $|\Delta[n]| = D^n$, $|\partial\Delta[n]| = S^{n-1}$, $|\Lambda^r[n]| = D^{n-1}$.

Proposition 1.3.1.29. *As a functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{K}}$, $|-|$ preserves finite products.*

Remark 1.3.1.30. As a functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Top}}$ $|-|$ need not preserve finite products. This is because the product need not preserve colimits.

1.3.2 The Standard Model Structure on Simplicial Sets

Definition 1.3.2.1. Define the set $I = \{i : \partial\Delta[n] \rightarrow \Delta[n] \mid n \geq 0\}$, and the set $J = \{j : \Lambda^r[n] \rightarrow \Delta[n] \mid n > 0, 0 \leq r \leq n\}$. Define $\operatorname{Cof} = \operatorname{Cof}(I)$, $\operatorname{Fib} = \operatorname{RLP}(J)$, and $\operatorname{W} = |-|^{-1}(\operatorname{W}_{\underline{\mathbf{Top}}})$. Define a map to be a **Kan fibration** if and only if it is in Fib . Define a simplicial set to be a **Kan complex** if and only if it is fibrant. Define a map to be an **anodyne extension** if and only if it is in $\operatorname{Cof}(J)$.

Definition 1.3.2.2. If $p : X \rightarrow Y$ is a fibration, $v : \Delta[0] \rightarrow Y$ is a vertex in Y , define $\Delta[0] \times_Y X$ to be the **fiber of p over v** , denoted X_v .

Proposition 1.3.2.3. A map is a cofibration if and only if it is injective. In particular, any simplicial set is cofibrant, and $\text{Cof}(J) \subseteq \text{Cof}(I)$. Moreover, any cofibration is a relative I -cell complex.

Proposition 1.3.2.4. We have $|I|$ is the set of generating cofibrations of $\underline{\mathbf{K}}$ and $|J|$ is the set of generating trivial cofibrations of $\underline{\mathbf{K}}$. Thus all maps in $|\text{Cof}(I)|$ are cofibrations, and all maps in $|\text{Cof}(J)|$ are trivial cofibrations. In particular, all anodyne extensions are trivial cofibrations. Moreover the singular functor takes fibrations between k -spaces to Kan fibrations and takes trivial fibrations between k -spaces to $\text{RLP}(I)$.

Lemma 1.3.2.5. As a functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{K}}$, $|-|$ preserves all finite limits, and in particular, pullbacks.

Lemma 1.3.2.6. For any map $f \in \text{RLP}(I)$, $|f|$ is a fibration.

Lemma 1.3.2.7. For any map $f \in \text{RLP}(I)$, f is a trivial fibration.

Definition 1.3.2.8. Suppose $i : K \rightarrow L$ and $p : X \rightarrow Y$ are maps between simplicial sets. We define $P(i, p) := (K \times Y) \amalg_{K \times X} (L \times X)$, and $i \square p$ being the induced map $P(i, p) \rightarrow L \times Y$. Moreover we define p_{\square}^i to be the induced map $X^L \rightarrow X^K \times_{Y^K} Y^L$.

Lemma 1.3.2.9. We have that $f \square g \nearrow h$ if and only if $f \nearrow h_{\square}^g$ for any maps f, g, h .

Lemma 1.3.2.10. Suppose $i : \partial\Delta[n] \rightarrow \Delta[n]$, $f : \Lambda^r[1] \rightarrow \Delta[1]$ are the inclusions, where $r = 0, 1$. Then the map $i \square f$ is an anodyne extension.

Proposition 1.3.2.11. Suppose $f : \Lambda^r[1] \rightarrow \Delta[1]$ is the inclusion and $i : K \rightarrow L$ is an arbitrary injective map, where $r = 0, 1$. Then the map $i \square f$ is an anodyne extension.

Proposition 1.3.2.12. A map is an anodyne extension if and only if it is in $\text{Cof}(J')$, where $J' = \{i \square f | i \in I, f : \Lambda^r[1] \rightarrow \Delta[1], r = 0, 1\}$.

Theorem 1.3.2.13. If f is an anodyne extension and i is an arbitrary injective map, then $i \square f$ is an anodyne extension.

Theorem 1.3.2.14. If p is a fibration and i is an arbitrary injective map, then p_{\square}^i is a fibration.

Definition 1.3.2.15. Suppose X is a fibrant simplicial set and x, y are 0-simplices, we define x to be **homotopic** to y , denoted $x \sim y$, if there exists some 1-simplex z with $d_0 z = x, d_1 z = y$.

Lemma 1.3.2.16. If X is a fibrant simplicial set, then \sim is an equivalence relation on X_0 .

Definition 1.3.2.17. If X is a fibrant simplicial set, define $\pi_0 X = X / \sim$ and denote elements in $\pi_0 X$ the **path components** of X . Moreover if v is a 0-simplex of X , define $\pi_0(X, v) = (\pi_0 X, [v])$.

Lemma 1.3.2.18. $\pi_0 : \underline{\mathbf{SSet}}_f \rightarrow \underline{\mathbf{Set}}$ and $\pi_0 : (\underline{\mathbf{SSet}}_f)_* \rightarrow \underline{\mathbf{Set}}_*$ are functorial. If X is a fibrant simplicial set, then $\pi_0 X$ and $\pi_0 |X|$ are naturally isomorphic.

Definition 1.3.2.19. Suppose v is a vertex of a simplicial set X , and Y is another simplicial set. We define the **constant map at v** from Y to X to be the composite $Y \rightarrow \Delta[0] \xrightarrow{v} X$.

Definition 1.3.2.20. Suppose X is a fibrant simplicial set and v is a vertex. Let F be the fiber of the fibration $X^{\Delta[n]} \rightarrow X^{\partial\Delta[n]}$ over v . Define the **n -th homotopy group** $\pi_n(X, v)$ of X at v to be $\pi_0(F, v)$.

Lemma 1.3.2.21. *Given a map $f : (X, v) \rightarrow (Y, w)$, f induces a map $f_* : \pi_n(X, v) \rightarrow \pi_n(Y, w)$, thus $\pi_n : (\underline{\mathbf{SSet}}_f)_* \rightarrow \underline{\mathbf{Set}}_*$ is functorial.*

Definition 1.3.2.22. Suppose $f, g : K \rightarrow X$ are maps of simplicial sets, we denote a map $H : K \times \Delta[1] \rightarrow X$ to be a **homotopy** from f to g if $H|_{K \times 0} = f$ and $H|_{K \times 1} = g$. If such H exists we denote $f \simeq g$.

Lemma 1.3.2.23. *If X is fibrant, then for any simplicial set K , \simeq is an equivalence relation on $\text{Hom}_{\underline{\mathbf{SSet}}}(K, X)$, and $f \simeq g$ if and only if $f \sim g$ as vertices in X^K . Moreover*

$$\pi_n(X, v) = (\text{Hom}_{\underline{\mathbf{SSet}}}(\Delta[n], X) / \simeq, [v]).$$

Lemma 1.3.2.24. *Suppose X is fibrant, v is a vertex of f , a is an n -simplex such that $d_i a = v$ for all i . Then $[a] = [v] \in \pi_n(X, v)$ if and only if there is an $(n+1)$ -simplex x such that $d_{n+1} x = a$ and $d_i x = v$ if $i \leq n$.*

Definition 1.3.2.25. Suppose K is a subsimplicial set of L , $i : K \rightarrow L$ is the inclusion. A map $r : L \rightarrow K$ is called a **retraction** from L to K if $ri = \mathbb{1}$; A map $H : L \times \Delta[1] \rightarrow L$ is called a **deformation retraction** from L to K if $H|_{L \times 0} = \mathbb{1}$, $H|_{L \times 1}$ is a retraction, and $H|_{K \times \Delta[1]}$ is the projection onto K .

Example 1.3.2.26. The vertex n is a deformation retraction of $\Delta[n]$, and the deformation retraction restricts to a deformation retraction from $\Lambda^n[n]$ to n .

Remark 1.3.2.27. If $n > 0$ then $n \not\sim \mathbb{1}_{\Delta[n]}$, thus $\Delta[n]$ is not fibrant.

Proposition 1.3.2.28. *If X is a nonempty fibrant simplicial set with no nontrivial homotopy groups, then the map $X \rightarrow \Delta[0]$ is in $\text{RLP}(I)$.*

Definition 1.3.2.29. Suppose $p : X \rightarrow Y$ is a fibration between fibrant simplicial sets, v is a vertex of X , F is the fiber of p over $p(v)$. Define the map $\partial : \pi_n(Y, p(v)) \mapsto \pi_{n-1}(F, v)$ as follows: for any $[\alpha] \in \pi_n(Y, p(v))$, define γ to be a lift of

$$\begin{array}{ccc} \Lambda^n[n] & \xrightarrow{v} & X \\ \downarrow & \nearrow \gamma & \downarrow p \\ \Delta[n] & \xrightarrow{\alpha} & Y \end{array}$$

and define $\partial[\alpha] = [d_n \gamma]$.

Lemma 1.3.2.30. $\partial : \pi_n(Y, p(v)) \mapsto \pi_{n-1}(F, v)$ is well-defined and is natural for maps between fibrations.

Proposition 1.3.2.31. *Suppose $p : X \rightarrow Y$ is a fibration between fibrant simplicial sets, v is a vertex of X , F is the fiber of p over $p(v)$. Then we have an exact sequence of pointed sets:*

$$\begin{aligned} \cdots \xrightarrow{\partial} \pi_n(F, v) &\xrightarrow{i_*} \pi_n(X, v) \xrightarrow{p_*} \pi_n(Y, p(v)) \\ &\xrightarrow{\partial} \pi_{n-1}(F, v) \xrightarrow{i_*} \cdots \xrightarrow{i_*} \pi_0(X, v) \xrightarrow{p_*} \pi_0(Y, p(v)), \end{aligned}$$

which is called the **long exact sequence**.

Definition 1.3.2.32. We define a fibration $p : X \rightarrow Y$ to be **locally trivial** if for any simplex $y : \Delta[n] \rightarrow Y$ there exists some simplicial set F , such that the pullback fibration $y^*p : y^*X := \Delta[n] \times_Y X \rightarrow \Delta[n]$ is isomorphic over $\Delta[n]$ to the projection $\Delta[n] \times F \rightarrow \Delta[n]$.

Proposition 1.3.2.33. *If p is a locally trivial fibration such that every fiber of p is non-empty and has no nontrivial homotopy groups, then $p \in \text{RLP}(I)$.*

Proposition 1.3.2.34. *Suppose $p : X \rightarrow Y$ is a fibration and $f \simeq g : K \rightarrow Y$. Then the pullback fibrations f^*p and g^*p are **fiber homotopy equivalent**, i.e. there exists maps $r : f^*X \rightarrow g^*X$ and $s : g^*X \rightarrow f^*X$ such that $f^*p \circ s = g^*p$, $g^*p \circ r = f^*p$, and there exists homotopies $H : rs \simeq \mathbb{1}_{g^*X}$ and $H' : sr \simeq \mathbb{1}_{f^*X}$ with $g^*p \circ H = \text{pr}_1 \circ (g^*p \times \mathbb{1})$, $f^*p \circ H' = \text{pr}_1 \circ (f^*p \times \mathbb{1})$.*

Corollary 1.3.2.35. *Suppose $p : X \rightarrow Y$ is a fibration, and y is an n -simplex of Y , then the pullback fibration y^*p is fiber homotopy equivalent to the projection fibration $\Delta[n] \times F \rightarrow \Delta[n]$, where F is the fiber of p over $y(n)$.*

Definition 1.3.2.36. Suppose $p : X \rightarrow Y$ is a fibration, and x, y are two n -simplices of X . We say x and y are **p -related**, written $x \sim_p y$, if they represent vertices in the same path component in the same fiber of the map $p_{\square}^{\partial\Delta[n] \rightarrow \Delta[n]}$. We say p is a **minimal fibration** if $x \sim_p y$ implies $x = y$.

Lemma 1.3.2.37. *Suppose $p : X \rightarrow Y$ is a fibration, and x, y are two n -simplices of X . Then \sim_p is an equivalence relation, and $x \sim_p y$ if and only if $px = py$, $d_i x = d_i y$ for any $0 \leq i \leq n$, and there exists a homotopy $H : x \simeq y$ such that $H|_{\partial\Delta[n] \times \Delta[1]}$ and pH are constant homotopies.*

Lemma 1.3.2.38. *Suppose $p : X \rightarrow Y$, $q : Z \rightarrow Y$ are fibrations and q is minimal, $f, g : X \rightarrow Z$, $H : f \simeq g$ such that $qH = \text{pr}_1(p \times \mathbb{1})$. If g is an isomorphism, so is f .*

Corollary 1.3.2.39. *Any minimal fibration is locally trivial.*

Lemma 1.3.2.40. *Suppose $p : X \rightarrow Y$ is a fibration and x, y are degenerate n -simplices in X that are p -related. Then $x = y$.*

Theorem 1.3.2.41. *Any fibration p can be factored into $p'r$, where r is a retraction onto a subsimplicial set of the domain such that $r \in \text{RLP}(I)$, and p' is a minimal fibration.*

Corollary 1.3.2.42. *If p is a fibration such that every fiber of p is non-empty and has no nontrivial homotopy groups, then $p \in \text{RLP}(I)$.*

Proposition 1.3.2.43. *If p is a locally trivial fibration then $|p|$ is a fibration.*

Corollary 1.3.2.44. *If p is a fibration then $|p|$ is a fibration.*

Proposition 1.3.2.45. *There exists a natural isomorphism $\pi_n(X, v) \rightarrow \pi_n(|X|, |v|)$ for any $(X, v) \in \underline{\text{SSet}}_*$.*

Remark 1.3.2.46. Suppose X is a fibrant simplicial set, v is a vertex of X . For any $[\alpha], [\beta] \in \pi_1(X, v)$, take γ to be a lift of the following diagram:

$$\begin{array}{ccc} \Lambda^1[2] & \xrightarrow[\beta \text{ on } d_0 i_2]{\alpha \text{ on } d_2 i_2} & X \\ \downarrow & \nearrow \gamma & \downarrow \\ \Delta[2] & \longrightarrow & * \end{array}$$

and define $[\alpha] \cdot [\beta] = [d_1 \gamma]$; take δ to be the lift of a following diagram:

$$\begin{array}{ccc} \Lambda^0[2] & \xrightarrow[v \text{ on } d_1 i_2]{\alpha \text{ on } d_2 i_2} & X \\ \downarrow & \nearrow \delta & \downarrow \\ \Delta[2] & \longrightarrow & * \end{array}$$

and define $[\alpha]^{-1} = [d_0 \delta]$. Then this gives a natural group structure on $\pi_1(X, v)$ making the map $\pi_1(X, v) \rightarrow \pi_1(|X|, |v|)$ a natural group isomorphism.

Theorem 1.3.2.47. *If p is a locally trivial fibration, then $p \in \text{RLP}(I)$.*

Theorem 1.3.2.48. *$\underline{\mathbf{SSet}}$ is a finitely generated model category with I being its generating cofibrations and J being its generating trivial cofibrations. As a corollary, $\underline{\mathbf{SSet}}_*$ is a finitely generated model category with I_+ being its generating cofibrations and J_+ being its generating trivial cofibrations. This model structure is called the **standard model structure** on $\underline{\mathbf{SSet}}$.*

Theorem 1.3.2.49. *$(|-|, \text{Sing}, \varphi)$ is a Quillen equivalence between $\underline{\mathbf{SSet}}$ and $\underline{\mathbf{K}}$. $(|-|_*, \text{Sing}_*, \varphi_*)$ is a Quillen equivalence between $\underline{\mathbf{SSet}}_*$ and $\underline{\mathbf{K}}_*$.*

Proposition 1.3.2.50. *Suppose \mathcal{C} is a model category and $F : \underline{\mathbf{SSet}} \rightarrow \mathcal{C}$ is a functor that preserves colimits and cofibrations. Then F preserves trivial cofibrations if and only if $F(\Delta[n]) \rightarrow F(\Delta[0])$ is a weak equivalence for any n .*

Corollary 1.3.2.51. *Suppose \mathcal{C} is a model category and $F : \underline{\mathbf{SSet}}_* \rightarrow \mathcal{C}$ is a functor that preserves colimits and cofibrations. Then F preserves trivial cofibrations if and only if $F(\Delta[n]_+) \rightarrow F(\Delta[0]_+)$ is a weak equivalence for any n .*

1.4 Monoidal Model Categories

1.4.1 Monoidal Categories

Definition 1.4.1.1. Suppose \mathcal{C} is a category. A **monoidal structure** on \mathcal{C} consists of a quintuple (\otimes, S, a, l, r) , where $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is a functor, S is an object in \mathcal{C} , $a_{XYZ} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$, $l_X : S \otimes X \rightarrow X$, $r_X : X \otimes S \rightarrow X$ are natural isomorphisms, such that the following diagrams are commutative for any objects X, Y, Z, W :

$$\begin{array}{ccc}
((X \otimes Y) \otimes Z) \otimes W & \xrightarrow{a_{XYZ} \otimes 1_W} & (X \otimes (Y \otimes Z)) \otimes W \\
a_{(X \otimes Y)ZW} \downarrow & & \downarrow a_{X(Y \otimes Z)W} \\
(X \otimes Y) \otimes (Z \otimes W) & & \\
a_{XY(Z \otimes W)} \downarrow & & \\
X \otimes (Y \otimes (Z \otimes W)) & \xleftarrow{1_X \otimes a_{YZW}} & X \otimes ((Y \otimes Z) \otimes W)
\end{array}$$

$$\begin{array}{ccc}
(X \otimes S) \otimes Y & \xrightarrow{a_{XSY}} & X \otimes (S \otimes Y) \\
r_X \otimes 1_Y \searrow & & \swarrow 1_X \otimes l_Y \\
& X \otimes Y &
\end{array}$$

$$\begin{array}{ccc}
S \otimes S & & \\
r_S \downarrow & & \downarrow l_S \\
& S &
\end{array}$$

A category together with a monoidal structure is called a **monoidal category**.

Definition 1.4.1.2. Suppose \mathcal{C}, \mathcal{D} are monoidal categories. A **monoidal functor** between \mathcal{C} and \mathcal{D} is a triple (F, μ, α) , where $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, $\mu_{XY} : FX \otimes FY \rightarrow F(X \otimes Y)$ is a natural isomorphism, $\alpha : FS \rightarrow S$ is an isomorphism, such that the following diagrams are commutative for any objects X, Y, Z :

$$\begin{array}{ccc}
(FX \otimes FY) \otimes FZ & \xrightarrow{a_{(FX)(FY)(FZ)}} & FX \otimes (FY \otimes FZ) \\
\mu_{XY} \otimes 1_{FZ} \downarrow & & \downarrow 1_{FX} \otimes \mu_{YZ} \\
F(X \otimes Y) \otimes FZ & & FX \otimes F(Y \otimes Z) \\
\mu_{(X \otimes Y)Z} \downarrow & & \downarrow \mu_{X(Y \otimes Z)} \\
F((X \otimes Y) \otimes Z) & \xrightarrow{F(a_{XYZ})} & F(X \otimes (Y \otimes Z))
\end{array}$$

$$\begin{array}{ccc}
FS \otimes FX & \xrightarrow{\alpha \otimes \mathbb{1}_{FX}} & S \otimes FX \\
\mu_{SX} \downarrow & & \downarrow l_{FX} \\
F(S \otimes X) & \xrightarrow{F(l_X)} & FX
\end{array}
\quad
\begin{array}{ccc}
FX \otimes FS & \xrightarrow{\mathbb{1}_{FX} \otimes \alpha} & FX \otimes S \\
\mu_{XS} \downarrow & & \downarrow r_{FX} \\
F(X \otimes S) & \xrightarrow{F(r_X)} & FX
\end{array}$$

Definition 1.4.1.3. Suppose \mathcal{C}, \mathcal{D} are monoidal categories and F, F' are monoidal functors between \mathcal{C} and \mathcal{D} . A **monoidal natural transformation** between F and F' is a natural transformation $\tau : F \rightarrow F'$ such that the following diagrams are commutative for any objects X, Y :

$$\begin{array}{ccc}
FX \otimes FY & \xrightarrow{\mu_{XY}} & F(X \otimes Y) \\
\tau_X \otimes \tau_Y \downarrow & & \downarrow \tau_{X \otimes Y} \\
F'X \otimes F'Y & \xrightarrow{\mu'_{XY}} & F'(X \otimes Y)
\end{array}
\quad
\begin{array}{ccc}
FS & \xrightarrow{\alpha_S} & S \\
\tau_S \searrow & & \nearrow \alpha'_S \\
& F'S &
\end{array}$$

Lemma 1.4.1.4. *Monoidal categories, monoidal functors and monoidal natural transformations form a 2-category, which we denote Mon.*

Definition 1.4.1.5. Suppose \mathcal{C} is a category. A **symmetric monoidal structure** on \mathcal{C} is a monoidal structure \otimes together with a natural isomorphism $T_{XY} : X \otimes Y \rightarrow Y \otimes X$, such that the following diagrams are commutative for any objects X, Y, Z :

$$\begin{array}{ccc}
(X \otimes Y) \otimes Z & \xrightarrow{T_{XY} \otimes \mathbb{1}_Z} & (Y \otimes X) \otimes Z \xrightarrow{a_{YXZ}} Y \otimes (X \otimes Z) \\
a_{XYZ} \downarrow & & \downarrow \mathbb{1}_Y \otimes T_{XZ} \\
X \otimes (Y \otimes Z) & \xrightarrow{T_{X(Y \otimes Z)}} & (Y \otimes Z) \otimes X \xrightarrow{a_{YZX}} Y \otimes (Z \otimes X)
\end{array}$$

$$\begin{array}{ccc}
X \otimes Y & & S \otimes S \\
T_{YX} \uparrow & & \uparrow \mathbb{1} \\
Y \otimes X & & S \otimes S
\end{array}
\quad
\begin{array}{ccc}
X \otimes S & \xrightarrow{r_X} & X \\
T_{XS} \searrow & & \nearrow T_{SX} \\
& S \otimes X &
\end{array}$$

A category together with a symmetric monoidal structure is called a **symmetric monoidal category**.

Definition 1.4.1.6. Suppose \mathcal{C}, \mathcal{D} are symmetric monoidal categories. A **symmetric monoidal functor** between \mathcal{C} and \mathcal{D} is a monoidal functor $F : \mathcal{C} \rightarrow \mathcal{D}$, such that the following diagram is commutative for any objects X, Y :

$$\begin{array}{ccc}
FX \otimes FY & \xrightarrow{\mu_{XY}} & F(X \otimes Y) \\
T_{(FX)(FY)} \downarrow & & \downarrow FT_{XY} \\
FY \otimes FX & \xrightarrow{\mu_{YX}} & F(Y \otimes X)
\end{array}$$

Lemma 1.4.1.7. *Symmetric monoidal categories, symmetric monoidal functors and monoidal natural transformations form a 2-category, which we denote SymMon. There is a forgetful 2-functor from SymMon to Mon.*

Definition 1.4.1.8. Suppose \mathcal{C} is a monoidal category. A **(right) \mathcal{C} -module structure** on a category \mathcal{D} is a triple (\otimes, a, r) , where $\otimes : \mathcal{D} \times \mathcal{C} \rightarrow \mathcal{D}$ is a functor, $a_{XKL} : (X \otimes K) \otimes L \rightarrow X \otimes (K \otimes L)$, $r_X : X \otimes S \rightarrow X$ are natural isomorphisms, such that the following diagrams are

commutative for any $X \in \mathcal{D}, K, L, M \in \mathcal{C}$:

$$\begin{array}{ccc}
((X \otimes K) \otimes L) \otimes M & \xrightarrow{a_{XKL} \otimes \mathbb{1}_M} & (X \otimes (K \otimes L)) \otimes M \\
\downarrow a_{(X \otimes K)LM} & & \downarrow a_{X(K \otimes L)M} \\
(X \otimes K) \otimes (L \otimes M) & & \\
\downarrow a_{XK(L \otimes M)} & & \\
X \otimes (K \otimes (L \otimes M)) & \xleftarrow{\mathbb{1}_X \otimes a_{KLM}} & X \otimes ((K \otimes L) \otimes M)
\end{array}$$

$$\begin{array}{ccc}
(X \otimes S) \otimes K & \xrightarrow{a_{XSK}} & X \otimes (S \otimes K) \\
\searrow r_X \otimes \mathbb{1}_K & & \swarrow \mathbb{1}_X \otimes l_K \\
& X \otimes K &
\end{array}$$

$$\begin{array}{ccc}
(X \otimes K) \otimes S & \xrightarrow{a_{XKS}} & X \otimes (K \otimes S) \\
\searrow r_X \otimes K & & \swarrow \mathbb{1}_X \otimes r_K \\
& X \otimes K &
\end{array}$$

A category together with a \mathcal{C} -module structure is called a **(right) \mathcal{C} -module**.

Definition 1.4.1.9. Suppose \mathcal{D}, \mathcal{E} are \mathcal{C} -modules. A **(right) \mathcal{C} -module functor** between \mathcal{D} and \mathcal{E} is a pair (F, μ) , where $F : \mathcal{D} \rightarrow \mathcal{E}$ is a functor, $\mu_{XK} : FX \otimes K \rightarrow F(X \otimes K)$ is a natural isomorphism, such that the following diagrams are commutative for any objects $X \in \mathcal{D}, K, L \in \mathcal{C}$:

$$\begin{array}{ccc}
(FX \otimes K) \otimes L & \xrightarrow{\mu_{XK} \otimes \mathbb{1}_L} & F(X \otimes K) \otimes L \\
\downarrow a_{(FX)KL} & & \downarrow \mu_{(X \otimes K)L} \\
FX \otimes (K \otimes L) & & \\
\downarrow \mu_{X(K \otimes L)} & & \\
F(X \otimes (K \otimes L)) & \xleftarrow{F(a_{XKL})} & F((X \otimes K) \otimes L)
\end{array}$$

$$\begin{array}{ccc}
FX \otimes S & \xrightarrow{\mu_{XS}} & F(X \otimes S) \\
\searrow r_{FX} & & \swarrow Fr_X \\
& X \otimes S &
\end{array}$$

Definition 1.4.1.10. Suppose \mathcal{D}, \mathcal{E} are \mathcal{C} -modules and F, F' are \mathcal{C} -module functors between \mathcal{D} and \mathcal{E} . A **(right) \mathcal{C} -module natural transformation** between F and F' is a natural transformation $\tau : F \rightarrow F'$ such that the following diagram is commutative for any objects $X \in \mathcal{D}, K \in \mathcal{C}$:

$$\begin{array}{ccc}
FX \otimes K & \xrightarrow{\mu_{XK}} & F(X \otimes K) \\
\downarrow \tau_X \otimes \mathbb{1}_K & & \downarrow \tau_{X \otimes K} \\
F'X \otimes K & \xrightarrow{\mu'_{XK}} & F'(X \otimes K)
\end{array}$$

Lemma 1.4.1.11. \mathcal{C} -modules, \mathcal{C} -module functors and \mathcal{C} -module natural transformations form a 2-category, which we denote $\underline{\mathbf{Mod}}_{\mathcal{C}}$. A monoidal functor between two monoidal categories \mathcal{C} and \mathcal{D} induces a forgetful 2-functor from $\underline{\mathbf{Mod}}_{\mathcal{D}}$ to $\underline{\mathbf{Mod}}_{\mathcal{C}}$.

Definition 1.4.1.12. Suppose \mathcal{C} is a monoidal category. A **\mathcal{C} -algebra structure** on a category \mathcal{D} is a monoidal structure on \mathcal{D} together with a monoidal functor $i : \mathcal{C} \rightarrow \mathcal{D}$. A category together with a \mathcal{C} -algebra structure is called a **\mathcal{C} -algebra**.

Definition 1.4.1.13. Suppose \mathcal{D}, \mathcal{E} are \mathcal{C} -algebras. A **\mathcal{C} -algebra functor** between \mathcal{D} and \mathcal{E} is a monoidal functor $F : \mathcal{D} \rightarrow \mathcal{E}$ together with a monoidal natural transformation $\rho : F \circ i_{\mathcal{D}} \rightarrow i_{\mathcal{E}}$.

Definition 1.4.1.14. Suppose \mathcal{D}, \mathcal{E} are \mathcal{C} -algebras and F, F' are \mathcal{C} -algebra functors between \mathcal{D} and \mathcal{E} . A **\mathcal{C} -algebra natural transformation** between F and F' is a monoidal natural transformation $\tau : F \rightarrow F'$ such that the following diagram is commutative for any object $K \in \mathcal{C}$:

$$\begin{array}{ccc} F(i(K)) & \xrightarrow{\tau_{i(K)}} & F'(i(K)) \\ \rho_K \searrow & & \swarrow \rho'_K \\ & i(K) & \end{array}$$

Lemma 1.4.1.15. \mathcal{C} -algebras, \mathcal{C} -algebra functors and \mathcal{C} -algebra natural transformations form a 2-category, which we denote $\underline{\mathbf{Alg}}_{\mathcal{C}}$. A monoidal functor between two monoidal categories \mathcal{C} and \mathcal{D} induces a forgetful 2-functor from $\underline{\mathbf{Alg}}_{\mathcal{D}}$ to $\underline{\mathbf{Alg}}_{\mathcal{C}}$.

Proposition 1.4.1.16. For any monoidal category \mathcal{C} , we have a forgetful 2-functor from $\underline{\mathbf{Alg}}_{\mathcal{C}}$ to $\underline{\mathbf{Module}}_{\mathcal{C}}$, where the \otimes -structure is given by $X \otimes K = X \otimes iK$ for any object X in some \mathcal{C} -algebra.

Definition 1.4.1.17. Suppose \mathcal{C} is a symmetric monoidal category. A **symmetric \mathcal{C} -algebra structure** on a category \mathcal{D} is a symmetric monoidal structure on \mathcal{D} together with a symmetric monoidal functor $i : \mathcal{C} \rightarrow \mathcal{D}$. A category together with a symmetric \mathcal{C} -algebra structure is called a **symmetric \mathcal{C} -algebra**.

Definition 1.4.1.18. Suppose \mathcal{D}, \mathcal{E} are symmetric \mathcal{C} -algebras. A **symmetric \mathcal{C} -algebra functor** between \mathcal{D} and \mathcal{E} is a \mathcal{C} -algebra functor $F : \mathcal{D} \rightarrow \mathcal{E}$ that is also a symmetric monoidal functor.

Lemma 1.4.1.19. Symmetric \mathcal{C} -algebras, symmetric \mathcal{C} -algebra functors and \mathcal{C} -algebra natural transformations form a 2-category, which we denote $\underline{\mathbf{SymAlg}}_{\mathcal{C}}$. There is a forgetful 2-functor from $\underline{\mathbf{SymAlg}}_{\mathcal{C}}$ to $\underline{\mathbf{Alg}}_{\mathcal{C}}$ for any monoidal category \mathcal{C} . A monoidal functor between two monoidal categories \mathcal{C} and \mathcal{D} induces a forgetful 2-functor from $\underline{\mathbf{SymAlg}}_{\mathcal{D}}$ to $\underline{\mathbf{SymAlg}}_{\mathcal{C}}$.

Definition 1.4.1.20. Suppose $\mathcal{C}, \mathcal{D}, \mathcal{E}$ are categories. An **adjunction of 2 variables** from $\mathcal{C} \times \mathcal{D}$ to \mathcal{E} is a quintuple $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r, \varphi_l, \varphi_r)$, where

$$\otimes : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}, \mathcal{H}om_l : \mathcal{C}^{\text{op}} \times \mathcal{E} \rightarrow \mathcal{D}, \mathcal{H}om_r : \mathcal{D}^{\text{op}} \times \mathcal{E} \rightarrow \mathcal{C}$$

are functors,

$$(\varphi_l)_{CDE} : \text{Hom}_{\mathcal{E}}(C \otimes D, E) \rightarrow \text{Hom}_{\mathcal{D}}(D, \mathcal{H}om_l(C, E))$$

$$(\varphi_r)_{CDE} : \text{Hom}_{\mathcal{E}}(C \otimes D, E) \rightarrow \text{Hom}_{\mathcal{C}}(C, \mathcal{H}om_r(D, E))$$

are natural isomorphisms.

Definition 1.4.1.21. Suppose \star is one of the structures given above. \star is called **closed** if all the structure bifunctors are adjunctions of 2 variables, and all the structure functors are left adjoints.

Definition 1.4.1.22. If \mathcal{C} is a closed monoidal category and \mathcal{D} is a closed \mathcal{C} -module, then we denote X^K to be $\mathcal{H}om_r(K, X)$ for $X \in \mathcal{D}, K \in \mathcal{C}$, and $\mathcal{H}om(X, Y)$ to be $\mathcal{H}om_l(X, Y)$ for $X, Y \in \mathcal{D}$.

Lemma 1.4.1.23. For any 2-category given above, all such categories with closed such structures, all such functors with closed such structures and all such natural transformations with closed such structures form a 2-category. We denote such 2-category by adding a Clo before the original one.

Proposition 1.4.1.24. *If \mathcal{C} is a closed symmetric monoidal category, then there is a duality 2-functor $-^{\text{op}}$ on $\underline{\text{CloMod}}_{\mathcal{C}}$, which maps a closed \mathcal{C} -module*

$$(\mathcal{D}, \otimes, \mathcal{H}om_l, \mathcal{H}om_r, a, r)$$

to

$$(\mathcal{D}^{\text{op}}, \otimes^{\text{op}}, \mathcal{H}om_l^{\text{op}}, \mathcal{H}om_r^{\text{op}}, a^{\text{op}}, r^{\text{op}}),$$

where $X \otimes^{\text{op}} K = \mathcal{H}om_r(K, X)$, $\mathcal{H}om_l^{\text{op}}(X, Y) = \mathcal{H}om_l(Y, X)$, $\mathcal{H}om_r^{\text{op}}(K, X) = X \otimes K$, a^{op} is defined by the natural isomorphism

$$\begin{aligned} & \text{Hom}_{\mathcal{D}}(Y, \mathcal{H}om_r(L, \mathcal{H}om_r(K, X))) \\ & \cong \text{Hom}_{\mathcal{D}}(Y \otimes L, \mathcal{H}om_r(K, X)) \\ & \cong \text{Hom}_{\mathcal{D}}((Y \otimes L) \otimes K, X) \\ & \cong \text{Hom}_{\mathcal{D}}(Y \otimes (L \otimes K), X) \\ & \cong \text{Hom}_{\mathcal{D}}(Y \otimes (K \otimes L), X) \\ & \cong \text{Hom}_{\mathcal{D}}(Y, \mathcal{H}om_r(K \otimes L, X)), \end{aligned}$$

r^{op} is defined by the natural isomorphism

$$\text{Hom}_{\mathcal{D}}(Y, \mathcal{H}om_r(S, X)) \cong \text{Hom}_{\mathcal{D}}(Y \otimes S, X) \cong \text{Hom}_{\mathcal{D}}(Y, X);$$

and maps a closed \mathcal{C} -module functor (F, G, μ) to (G, F, μ^{op}) , where μ^{op} is defined by the natural isomorphism

$$\begin{aligned} & \text{Hom}_{\mathcal{D}}(X, \mathcal{H}om_r(K, GY)) \\ & \cong \text{Hom}_{\mathcal{D}}(X \otimes K, GY) \\ & \cong \text{Hom}_{\mathcal{E}}(F(X \otimes K), Y) \\ & \cong \text{Hom}_{\mathcal{E}}(FX \otimes K, Y) \\ & \cong \text{Hom}_{\mathcal{E}}(FX, \mathcal{H}om_r(K, Y)) \\ & \cong \text{Hom}_{\mathcal{D}}(X, G \mathcal{H}om_r(K, Y)); \end{aligned}$$

and maps a closed \mathcal{C} -module natural transformation τ to τ^{op} . Moreover $(-^{\text{op}})^2 = 1$.

1.4.2 Monoidal Model Categories

Definition 1.4.2.1. Suppose that $\mathcal{C}, \mathcal{D}, \mathcal{E}$ are model categories. An adjunction of 2 variables $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r, \varphi_l, \varphi_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is called a **Quillen adjunction of 2 variables** if for any $f : A \rightarrow B \in \text{Cof}_{\mathcal{C}}, g : C \rightarrow D \in \text{Cof}_{\mathcal{D}}$, the **pushout product**

$$f \square g : A \otimes D \amalg_{A \otimes C} B \otimes C \rightarrow B \otimes D$$

is a cofibration, and is trivial if f or g is. If so \otimes is called a **Quillen bifunctor**.

Lemma 1.4.2.2. Suppose $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is an adjunction of 2 variables between model categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$. Then the following statements are equivalent:

- i) \otimes is a Quillen bifunctor;
- ii) For any $f : A \rightarrow B \in \text{Cof}_{\mathcal{C}}, p : X \rightarrow Y \in \text{Fib}_{\mathcal{E}}$, the map

$$\mathcal{H}om_{\square, l}(f, p) : \mathcal{H}om_l(A, X) \times_{\mathcal{H}om_l(B, X)} \mathcal{H}om_l(B, Y) \rightarrow \mathcal{H}om_l(A, Y)$$

is a fibration, and is trivial if f or p is;

iii) For any $g : C \rightarrow D \in \text{Cof}_{\mathcal{D}}, p : X \rightarrow Y \in \text{Fib}_{\mathcal{E}}$, the map

$$\mathcal{H}om_{\square, r}(g, p) : \mathcal{H}om_r(C, X) \times_{\mathcal{H}om_r(D, X)} \mathcal{H}om_r(D, Y) \rightarrow \mathcal{H}om_r(C, Y)$$

is a fibration, and is trivial if g or p is.

Lemma 1.4.2.3. Suppose $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is a Quillen adjunction of 2 variables between model categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$. If C is cofibrant in \mathcal{C} , then $(C \otimes -, \mathcal{H}om_l(C, -)) : \mathcal{D} \rightarrow \mathcal{E}$ is a Quillen adjunction. If D is cofibrant in \mathcal{D} , then $(- \otimes D, \mathcal{H}om_r(D, -)) : \mathcal{C} \rightarrow \mathcal{E}$ is a Quillen adjunction. If E is fibrant in \mathcal{E} , then $(\mathcal{H}om_l(-, E), \mathcal{H}om_r(-, E)) : \mathcal{C} \rightarrow \mathcal{D}^{\text{op}}$ is a Quillen adjunction.

Lemma 1.4.2.4. Suppose $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is an adjunction of 2 variables between categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$. Suppose I, I' are collections of maps in \mathcal{C}, \mathcal{D} , respectively. Then $\text{Cof}(I) \square \text{Cof}(I') \subseteq \text{Cof}(I \square I')$.

Corollary 1.4.2.5. Suppose $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is an adjunction of 2 variables between model categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$. Suppose more that \mathcal{C}, \mathcal{D} are cofibrantly generated model categories, with generating cofibrations I, I' , generating trivial cofibrations J, J' , respectively. Then \otimes is a Quillen bifunctor, if and only if $I \square I' \subseteq \text{Cof}_{\mathcal{E}}, I \square J' \subseteq W_{\mathcal{E}} \cap \text{Cof}_{\mathcal{E}}, J \square I' \subseteq W_{\mathcal{E}} \cap \text{Cof}_{\mathcal{E}}$.

Remark 1.4.2.6. If \mathcal{C} is a closed monoidal category, then $(\square, \mathcal{H}om_{\square, l}, \mathcal{H}om_{\square, r})$ forms a closed monoidal structure on $\mathcal{C}^{[1]}$.

Definition 1.4.2.7. A **monoidal model category** is a model category together with a closed monoidal structure, such that:

- i) The monoidal structure bifunctor \otimes is a Quillen bifunctor;
- ii) For any X cofibrant, the natural transformation $QS \otimes X \rightarrow S \otimes X$ is a weak equivalence;
- iii) For any X cofibrant, the natural transformation $X \otimes QS \rightarrow X \otimes S$ is a weak equivalence.

A **symmetric monoidal model category** is a monoidal model category that is also a symmetric monoidal category.

Lemma 1.4.2.8. Suppose \mathcal{C} is a model category that is also a closed monoidal category. Then for any X cofibrant, the natural transformation $QS \otimes X \rightarrow X$ is a weak equivalence, if and only if for any fibrant X , the natural transformation $X \rightarrow \mathcal{H}om_l(QS, X)$ is a weak equivalence; For any X cofibrant, the natural transformation $X \otimes QS \rightarrow X$ is a weak equivalence, if and only if for any fibrant X , the natural transformation $X \rightarrow \mathcal{H}om_r(QS, X)$ is a weak equivalence.

Proposition 1.4.2.9. Suppose \mathcal{C} is a monoidal model category with the unit being the terminal object $*$. Suppose more that $*$ is cofibrant. Then \mathcal{C}_* is a monoidal model category, with $X \wedge Y$ defined to be the pushout

$$\begin{array}{ccc} X \amalg Y & \xrightarrow{(\mathbb{1}_X \otimes w) \amalg (v \otimes \mathbb{1}_Y)} & X \otimes Y \\ \downarrow & & \downarrow \\ * & \xrightarrow{\quad} & X \wedge Y \end{array}$$

for any $(X, v), (Y, w) \in \mathcal{C}_*$, with the unit of \wedge being $*_+$; $\mathcal{H}om_{l,*}(X, Y)$ defined to be the pullback

$$\begin{array}{ccc} \mathcal{H}om_{l,*}(X, Y) & \xrightarrow{\quad} & \mathcal{H}om_l(X, Y) \\ \downarrow & & \downarrow \mathcal{H}om_l(v, Y) \\ * & \xrightarrow{\mathcal{H}om_l(X, w)} & \mathcal{H}om_l(*, Y) \end{array}$$

for any $(X, v), (Y, w) \in \mathcal{C}_*$, with the basepoint of $\mathcal{H}om_{l,*}(X, Y)$ being induced by the maps $* \rightarrow *$ and

$$\varphi_l(X \otimes * \cong X \rightarrow * \xrightarrow{w} Y) : * \rightarrow \mathcal{H}om_l(X, Y);$$

$\mathcal{H}om_{r,*}(X, Y)$ defined similarly. If \mathcal{C} is a symmetric monoidal model category, so is \mathcal{C}_* .

Proposition 1.4.2.10. SSet is a symmetric monoidal model category, with $\otimes = \times$, $\mathcal{H}om_l = \mathcal{H}om_r = -^-$. SSet_{*} is a symmetric monoidal model category.

Proposition 1.4.2.11. K is a symmetric monoidal model category, with $\otimes = \times$, $\mathcal{H}om_l = \mathcal{H}om_r = \mathcal{H}om$. K_{*} is a symmetric monoidal model category.

Remark 1.4.2.12. Top is not a monoidal model category.

Proposition 1.4.2.13. If R is a commutative ring, then Ch_R with the projective model structure is a symmetric monoidal model category, where $X \otimes Y$ is defined to be

$$(X \otimes Y)_n = \bigoplus_{k \in \mathbb{Z}} X_k \otimes_R Y_{n-k},$$

$$d_n(x \otimes y) = (dx) \otimes y + (-1)^{|x|} x \otimes (dy);$$

$\mathcal{H}om(X, Y) := \mathcal{H}om_l(X, Y) = \mathcal{H}om_r(X, Y)$ is defined to be

$$\mathcal{H}om(X, Y)_n = \prod_{k \in \mathbb{Z}} \text{Hom}(X_k, Y_{n+k}),$$

$$(d_n(f))(x) = d(f(x)) + (-1)^{n+1} f(dx);$$

The associativity isomorphism T is defined to be $T(x \otimes y) = (-1)^{|x||y|} y \otimes x$.

Proposition 1.4.2.14. If R is a ring, then the functor $\otimes : \underline{\text{Ch}}_{R^{\text{op}}} \times \underline{\text{Ch}}_R \rightarrow \underline{\text{Ch}}_{\mathbb{Z}}$ defined in Proposition 1.4.2.13 is a Quillen bifunctor.

Remark 1.4.2.15. Ch_R with the injective model structure is not a monoidal model category. For example, in Ch_ℤ,

$$(S^0(\mathbb{Z}) \rightarrow S^0(\mathbb{Q})) \square (0 \rightarrow S^0(\mathbb{Z}/2\mathbb{Z})) = (S^0(\mathbb{Z}/2\mathbb{Z}) \rightarrow 0),$$

which is certainly not injective.

Definition 1.4.2.16. Suppose \mathcal{C}, \mathcal{D} are monoidal model categories. A **monoidal Quillen adjunction** from \mathcal{C} to \mathcal{D} is a Quillen adjunction (F, G) such that F is monoidal, and the map $FQS \rightarrow QS$ is a weak equivalence. If so F is called a **monoidal Quillen functor**. A **symmetric monoidal Quillen functor** is a monoidal Quillen functor that is a symmetric monoidal functor.

Lemma 1.4.2.17. The composition of two monoidal Quillen functors is a monoidal Quillen functor.

Lemma 1.4.2.18. Monoidal model categories, monoidal Quillen functors and monoidal natural transformations form a 2-category, which we denote MonModel. Symmetric monoidal model categories, symmetric monoidal Quillen functors and monoidal natural transformations form a 2-category, which we denote SymMonModel.

Example 1.4.2.19. $-_+ : \underline{\text{SSet}} \rightarrow \underline{\text{SSet}}_*$, $-_+ : \underline{\text{K}} \rightarrow \underline{\text{K}}_*$ are monoidal Quillen adjunctions.

Example 1.4.2.20. If $R \rightarrow S$ is a ring homomorphism, then the induced Quillen adjunction $\underline{\text{Ch}}_R \rightarrow \underline{\text{Ch}}_S$ is a monoidal Quillen adjunction.

Proposition 1.4.2.21. $|-| : \underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{K}}, |-| : \underline{\mathbf{SSet}}_* \rightarrow \underline{\mathbf{K}}_*$ are monoidal Quillen adjunctions.

Definition 1.4.2.22. Suppose \mathcal{C} is a monoidal model category. A **\mathcal{C} -model category** is a model category \mathcal{D} together with a closed right \mathcal{C} -module structure, such that $\otimes : \mathcal{D} \times \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen bifunctor, and for any X cofibrant, the natural transformation $X \otimes QS \rightarrow X \otimes S$ is a weak equivalence. A $\underline{\mathbf{SSet}}$ -model category is called a **simplicial model category**.

Definition 1.4.2.23. Suppose \mathcal{C} is a monoidal model category. A **\mathcal{C} -Quillen functor** between \mathcal{C} -model categories is a Quillen functor that is also a closed \mathcal{C} -module functor.

Example 1.4.2.24. $\underline{\mathbf{SSet}}, \underline{\mathbf{SSet}}_*, \underline{\mathbf{K}}, \underline{\mathbf{K}}_*$ are simplicial model categories.

Example 1.4.2.25. For any ring R , $\underline{\mathbf{Ch}}_R$ is a $\underline{\mathbf{Ch}}_{\mathbb{Z}}$ -model category.

Lemma 1.4.2.26. If \mathcal{C} is a monoidal model category, all \mathcal{C} -model categories, \mathcal{C} -Quillen functors and \mathcal{C} -module natural transformations form a 2-category, which we denote $\underline{\mathbf{Model}}_{\mathcal{C}}$. Any monoidal Quillen functor between monoidal model categories \mathcal{C} and \mathcal{D} induces a forgetful 2-functor from $\underline{\mathbf{Model}}_{\mathcal{D}}$ to $\underline{\mathbf{Model}}_{\mathcal{C}}$.

Lemma 1.4.2.27. If \mathcal{C} is a pointed monoidal model category, every \mathcal{C} -model category is pointed.

Proposition 1.4.2.28. Suppose \mathcal{C} is a monoidal model category with the unit being the terminal object $*$. Suppose more that $*$ is cofibrant. If \mathcal{D} is a \mathcal{C} -model category, then \mathcal{D}_* is a \mathcal{C}_* -model category, where the \mathcal{C}_* -module structure is given similar to Proposition 1.4.2.9.

Lemma 1.4.2.29. If \mathcal{C} is a symmetric monoidal model category, the duality 2-functor on $\underline{\mathbf{Model}}$ and the duality 2-functor on $\underline{\mathbf{Mod}}_{\mathcal{C}}$ induces a duality 2-functor on $\underline{\mathbf{Model}}_{\mathcal{C}}$.

Definition 1.4.2.30. Suppose \mathcal{C} is a monoidal model category. A **monoidal \mathcal{C} -model category** is a monoidal model category \mathcal{D} together with a monoidal Quillen functor $\mathcal{C} \rightarrow \mathcal{D}$. A **monoidal \mathcal{C} -Quillen functor** between monoidal \mathcal{C} -model categories is a Quillen functor that is also a closed \mathcal{C} -algebra functor. We have similar definitions for a **symmetric monoidal \mathcal{C} -model category** and a **symmetric monoidal \mathcal{C} -Quillen functor**.

Lemma 1.4.2.31. If \mathcal{C} is a monoidal model category, all monoidal \mathcal{C} -model categories, monoidal \mathcal{C} -Quillen functors and \mathcal{C} -algebra natural transformations form a 2-category, which we denote $\underline{\mathbf{MonModel}}_{\mathcal{C}}$. Any monoidal Quillen functor between monoidal model categories \mathcal{C} and \mathcal{D} induces a forgetful 2-functor from $\underline{\mathbf{MonModel}}_{\mathcal{D}}$ to $\underline{\mathbf{MonModel}}_{\mathcal{C}}$. We have a similar atatement for the symmetric case.

Example 1.4.2.32. $\underline{\mathbf{K}}$ is a symmetric monoidal $\underline{\mathbf{SSet}}$ -model category.

Proposition 1.4.2.33. Suppose $(\otimes, \mathcal{H}om_l, \mathcal{H}om_r, \varphi_l, \varphi_r) : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ is a Quillen adjunction of 2 variables between model categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$. Then the derived functors defines an adjunction of 2 variables

$$(\otimes^L, R\mathcal{H}om_l, R\mathcal{H}om_r, \text{Ho } \varphi_l, \text{Ho } \varphi_r) : \text{Ho } \mathcal{C} \times \text{Ho } \mathcal{D} \rightarrow \text{Ho } \mathcal{E},$$

where $\text{Ho } \varphi_l$ is given by

$$\begin{aligned} [C \otimes^L D, E] &= [QC \otimes QD, E] \\ &\cong [QC \otimes QD, RE] \\ &\cong [QD, \mathcal{H}om_l(QC, RE)] \\ &\cong [D, \mathcal{H}om_l(QC, RE)] = [D, (R\mathcal{H}om_l)(C, E)]; \end{aligned}$$

and $\text{Ho } \varphi_r$ is given similarly.

Theorem 1.4.2.34. *The pseudo-2-functor $\text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{Cat}}_{ad}$ lifts to a pseudo-2-functor*

$$\text{Ho} : \underline{\text{MonModel}} \rightarrow \underline{\text{CloMon}},$$

where a monoidal model category

$$(\mathcal{C}, \otimes, \mathcal{H}om_l, \mathcal{H}om_r, \varphi_l, \varphi_r, a, l, r, S)$$

is mapped to

$$(\text{Ho } \mathcal{C}, \otimes^L, R \mathcal{H}om_l, R \mathcal{H}om_r, \text{Ho } \varphi_l, \text{Ho } \varphi_r, a, l, r, QS),$$

where a is given by

$$\begin{aligned} (X \otimes^L Y) \otimes^L Z &= Q(QX \otimes QY) \otimes QZ \\ &\cong (QX \otimes QY) \otimes QZ \\ &\cong QX \otimes (QY \otimes QZ) \\ &\cong QX \otimes Q(QY \otimes QZ) = X \otimes^L (Y \otimes^L Z), \end{aligned}$$

l is given by

$$S \otimes^L X = QS \otimes QX \cong S \otimes QX \cong QX \cong X,$$

r is given similarly; a monoidal Quillen adjunction

$$(F, G, \varphi, \mu, \alpha)$$

is mapped to

$$(LF, RG, \text{Ho } \varphi, \mu, \alpha),$$

where μ is given by

$$\begin{aligned} (LF)X \otimes^L (LF)Y &= QFQX \otimes QFQY \\ &\cong FQX \otimes FQY \\ &\cong F(QX \otimes QY) \\ &\cong FQ(QX \otimes QY) = (LF)(X \otimes^L Y), \end{aligned}$$

α is given by

$$(LF)S = FQS \cong FS \cong S;$$

a monoidal natural transformation τ is mapped to $L\tau$.

Theorem 1.4.2.35. *The pseudo-2-functor $\text{Ho} : \underline{\text{MonModel}} \rightarrow \underline{\text{CloMon}}$ lifts to a pseudo-2-functor*

$$\text{Ho} : \underline{\text{SymMonModel}} \rightarrow \underline{\text{CloSymMon}},$$

where the natural isomorphism T is given by

$$X \otimes^L Y = QX \otimes QY \cong QY \otimes QX = Y \otimes^L X.$$

Theorem 1.4.2.36. *Suppose \mathcal{C} is a monoidal model category, then the pseudo-2-functor $\text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{Cat}}_{ad}$ lifts to a pseudo-2-functor*

$$\text{Ho} : \underline{\text{Model}}_{\mathcal{C}} \rightarrow \underline{\text{CloMod}}_{\text{Ho } \mathcal{C}},$$

where the natural isomorphisms a, r, μ are defined similarly to that of Theorem 1.4.2.34; lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{MonModel}}_{\mathcal{C}} \rightarrow \underline{\text{CloAlg}}_{\text{Ho } \mathcal{C}},$$

where (\mathcal{D}, i) is mapped to $(\text{Ho } \mathcal{D}, Li)$, (F, ρ) is mapped to $(LF, L\rho)$; and lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{SymMonModel}}_{\mathcal{C}} \rightarrow \underline{\text{CloSymAlg}}_{\text{Ho } \mathcal{C}}$$

if \mathcal{C} is symmetric. In the latter case

$$\text{Ho} \circ (-^{\text{op}}) = (-^{\text{op}}) \circ \text{Ho} : \underline{\text{SymMonModel}}_{\mathcal{C}} \rightarrow \underline{\text{CloSymAlg}}_{\text{Ho } \mathcal{C}}.$$

Proposition 1.4.2.37. $L|-| : \text{Ho } \underline{\text{SSet}} \rightarrow \text{Ho } \underline{\text{K}}$ is a equivalence between closed symmetric monoidal categories. $L|-|_* : \text{Ho } \underline{\text{SSet}}_* \rightarrow \text{Ho } \underline{\text{K}}_*$ is a equivalence between closed symmetric monoidal categories.

1.5 Framings

1.5.1 Diagrams over Reedy Categories

Definition 1.5.1.1. Suppose \mathcal{B} is a small category, and λ is an ordinal. A functor $f : \mathcal{B} \rightarrow \lambda$ is called a **linear extension** if the image of a nonidentity map is nonidentity. If so define $f(i)$ to be the **degree** of i for any object $i \in \mathcal{B}$. If such linear extension exists we call \mathcal{B} a **direct category**. If \mathcal{B}^{op} is a direct category we call \mathcal{B} an **inverse category**.

Definition 1.5.1.2. Suppose \mathcal{C} is a category with all small colimits, \mathcal{B} is a direct category, $i \in \mathcal{B}$. Define the **latching space functor** $L_i : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ as the composition

$$\mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}^{\mathcal{B}_i} \xrightarrow{\text{colim}} \mathcal{C},$$

where \mathcal{B}_i is defined to be the category of all nonidentity maps in \mathcal{B} with codomain i . Dually, suppose \mathcal{C} is a category with all small limits, \mathcal{B} is an inverse category, $i \in \mathcal{B}$. Define the **matching space functor** $M_i : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ as the composition

$$\mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}^{\mathcal{B}^i} \xrightarrow{\text{lim}} \mathcal{C},$$

where \mathcal{B}^i is defined to be the category of all nonidentity maps in \mathcal{B} with domain i .

Lemma 1.5.1.3. If \mathcal{C} is a category with all small colimits, \mathcal{B} is a direct category, $i \in \mathcal{B}$, then there exists a natural transformation $L_i \rightarrow -_i := -(i)$; Dually if \mathcal{C} is a category with all small limits, \mathcal{B} is an inverse category, $i \in \mathcal{B}$, then there exists a natural transformation $-_i \rightarrow M_i$.

Proposition 1.5.1.4. Suppose \mathcal{C} is a model category, \mathcal{B} is a direct category, and we have a commutative diagram in $\mathcal{C}^{\mathcal{B}}$:

$$\begin{array}{ccc} A & \longrightarrow & X \\ f \downarrow & & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

such that $p_i : X_i \rightarrow Y_i$ is a fibration for any $i \in \mathcal{B}$, $g_i : A_i \amalg_{L_i A} L_i B \rightarrow B_i$ is a cofibration for any $i \in \mathcal{B}$. If f_i is a trivial fibration for any i or g_i is a trivial cofibration for any i , then the above diagram has a lift.

Corollary 1.5.1.5. Suppose \mathcal{C} is a model category, \mathcal{B} is a direct category, $f : A \rightarrow B \in \mathcal{C}^{\mathcal{B}}$. If the map $g_i : A_i \amalg_{L_i A} L_i B \rightarrow B_i$ is a (trivial) cofibration for any $i \in \mathcal{B}$, then $\text{colim } f : \text{colim } A \rightarrow \text{colim } B$ is a (trivial) cofibration.

Theorem 1.5.1.6. Suppose \mathcal{C} is a model category, \mathcal{B} is a direct category. Then there exists a model structure on $\mathcal{C}^{\mathcal{B}}$ where a map is a weak equivalence if and only if it is a degreewise weak equivalence; a map is a fibration if and only if it is a degreewise fibration; a map $f : A \rightarrow B$ is a (trivial) cofibration if and only if $g_i : A_i \amalg_{L_i A} L_i B \rightarrow B_i$ is a (trivial) cofibration for any $i \in \mathcal{B}$. Dually suppose \mathcal{C} is a model category, \mathcal{B} is an inverse category. Then there exists a model structure on $\mathcal{C}^{\mathcal{B}}$ where a map is a weak equivalence if and only if it is a degreewise weak equivalence; a map is a cofibration if and only if it is a degreewise cofibration; a map $p : X \rightarrow Y$ is a (trivial) fibration if and only if $q_i : X_i \rightarrow Y_i \times_{M_i Y} M_i X$ is a (trivial) fibration for any $i \in \mathcal{B}$.

Corollary 1.5.1.7. Suppose \mathcal{C} is a model category, \mathcal{B} is a direct category. Then the colimit functor $\text{colim} : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ is a left Quillen functor, left adjoint to the constant diagram functor. Dually suppose \mathcal{C} is a model category, \mathcal{B} is an inverse category. Then the limit functor $\text{lim} : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ is a right Quillen functor, right adjoint to the constant diagram functor.

Corollary 1.5.1.8. Suppose \mathcal{C} is a model category, \mathcal{B} is a direct category. If $f : A \rightarrow B$ is a (trivial) cofibration in $\mathcal{C}^{\mathcal{B}}$, then for any $i \in \mathcal{B}$, f_i is a (trivial) cofibration. Dually suppose \mathcal{C} is a model category, \mathcal{B} is an inverse category. If $p : X \rightarrow Y$ is a (trivial) fibration in $\mathcal{C}^{\mathcal{B}}$, then for any $i \in \mathcal{B}$, p_i is a (trivial) fibration.

Remark 1.5.1.9. Suppose \mathcal{C} is a cofibrantly generated model category with generating cofibrations I and generating trivial cofibrations J , and \mathcal{B} is a direct category. Then $\mathcal{C}^{\mathcal{B}}$ is a cofibrantly generated model category with generating cofibrations $\bigcup_{i \in \mathcal{B}} F_i I$ and generating trivial cofibrations $\bigcup_{i \in \mathcal{B}} F_i J$, where F_i is constructed to be a left adjoint of $-_i$.

Definition 1.5.1.10. Suppose \mathcal{B} is a small category. \mathcal{B} is called a **Reedy category** if there exists subcategories \mathcal{B}_+ , \mathcal{B}_- and a **degree function** $d : \text{Obj}(\mathcal{B}) \rightarrow \lambda$ where λ is some ordinal, such that all nonidentity morphisms in \mathcal{B}_+ raise degrees, all nonidentity morphisms in \mathcal{B}_- lower degrees, and for every map $f \in \mathcal{B}$, there exists unique maps $g \in \mathcal{B}_-$, $h \in \mathcal{B}_+$, such that $f = hg$. (In particular, \mathcal{B}_+ is a direct category and \mathcal{B}_- is an inverse category.)

Example 1.5.1.11. $\Delta, \Delta^{\text{op}}$ are Reedy categories. For any simplicial set K , ΔK is a Reedy category.

Lemma 1.5.1.12. If \mathcal{B} is a Reedy category, then \mathcal{B}^{op} is a Reedy category, where $(\mathcal{B}^{\text{op}})_+ = (\mathcal{B}_-)^{\text{op}}$ and $(\mathcal{B}^{\text{op}})_- = (\mathcal{B}_+)^{\text{op}}$.

Definition 1.5.1.13. Suppose \mathcal{C} is a category with all small colimits and limits, \mathcal{B} is a Reedy category, $i \in \mathcal{B}$. Define the **latching space functor** $L_i : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ as the composition

$$\mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}^{\mathcal{B}_+} \xrightarrow{L_i} \mathcal{C},$$

and the **matching space functor** $M_i : \mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}$ as the composition

$$\mathcal{C}^{\mathcal{B}} \rightarrow \mathcal{C}^{\mathcal{B}_-} \xrightarrow{M_i} \mathcal{C}.$$

Lemma 1.5.1.14. If \mathcal{C} is a category with all small colimits and limits, \mathcal{B} is a Reedy category, $i \in \mathcal{B}$, then there exists natural transformations $L_i \rightarrow -_i, -_i \rightarrow M_i$.

Example 1.5.1.15. Suppose \mathcal{C} is a category with all small colimits and limits. Then for any $A \in \mathcal{C}^{\Delta}$, $L_1 A = A[0] \amalg A[0]$, $M_1 A = A[0]$; for any $A \in \mathcal{C}^{\Delta^{\text{op}}}$, $L_1 A = A[0]$, $M_1 A = A[0] \times A[0]$.

Lemma 1.5.1.16. Suppose \mathcal{C} is a category with all small colimits and limits, \mathcal{B} is a Reedy category, define \mathcal{B}_{β} to be the full subcategory of \mathcal{B} of all objects with degree less than β for an ordinal β . Then given a functor $X : \mathcal{B}_{\beta} \rightarrow \mathcal{C}$, an extension of X to $X' : \mathcal{B}_{\beta+1} \rightarrow \mathcal{C}$ is equivalent to a choice of factorizations of maps $L_i X \rightarrow M_i X$ for all i 's with degree β ; given a natural transformation τ between two functors $X, Y : \mathcal{B}_{\beta} \rightarrow \mathcal{C}$ and extensions of X, Y to $X', Y' : \mathcal{B}_{\beta+1} \rightarrow \mathcal{C}$, an extension of τ to $\tau' : X' \rightarrow Y'$ is equivalent to a choice of maps $X'_i \rightarrow Y'_i$ making the following diagram commutative:

$$\begin{array}{ccccc} L_i X & \longrightarrow & X'_i & \longrightarrow & M_i X \\ L_i \tau \downarrow & & \downarrow & & \downarrow M_i \tau \\ L_i Y & \longrightarrow & Y'_i & \longrightarrow & M_i Y \end{array}$$

for all i 's with degree β .

Proposition 1.5.1.17. Suppose \mathcal{C} is a category with all small colimits and limits. Then there exists functors $l^\bullet, r^\bullet : \mathcal{C} \rightarrow \mathcal{C}^\Delta$ such that $l^\bullet(-)[0] = r^\bullet(-)[0] = \mathbb{1}$, $L_{[n]}l^\bullet = l^\bullet(-)[n]$, $M_{[n]}r^\bullet = r^\bullet(-)[n]$. Moreover we have $(l^\bullet, -[0])$, $(-[0], r^\bullet)$ are adjoint pairs, and for any $A \in \mathcal{C}$, $l^\bullet A[n] = \Pi_{n+1}A$, $r^\bullet A[n] = A$. Furthermore, $(l^\bullet, -[0])$, $(-[0], r^\bullet)$ are Quillen adjunctions. Dually there exists functors $l_\bullet, r_\bullet : \mathcal{C} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$ such that $l_\bullet(-)[0] = r_\bullet(-)[0] = \mathbb{1}$, $L_{[n]}l_\bullet = l_\bullet(-)[n]$, $M_{[n]}r_\bullet = r_\bullet(-)[n]$. Moreover we have $(l_\bullet, -[0])$, $(-[0], r_\bullet)$ are adjoint pairs, and for any $A \in \mathcal{C}$, $l_\bullet A[n] = A$, $r_\bullet A[n] = \Pi_{n+1}A$. Furthermore, $(l_\bullet, -[0])$, $(-[0], r_\bullet)$ are Quillen adjunctions.

Theorem 1.5.1.18. Suppose \mathcal{C} is a model category, \mathcal{B} is a Reedy category. Then there exists a model structure on $\mathcal{C}^\mathcal{B}$ where a map is a weak equivalence if and only if it is a degreewise weak equivalence; a map $f : A \rightarrow B$ is a (trivial) cofibration if and only if $g_i : A_i \amalg_{L_i A} L_i B \rightarrow B_i$ is a (trivial) cofibration for any $i \in \mathcal{B}$; a map $p : X \rightarrow Y$ is a (trivial) fibration if and only if $q_i : X_i \rightarrow Y_i \times_{M_i Y} M_i X$ is a (trivial) fibration for any $i \in \mathcal{B}$. Such model structure is called the **Reedy model structure**.

Proposition 1.5.1.19. If \mathcal{C} is a model category, \mathcal{B} is a Reedy category, then the Reedy model structure on $\mathcal{C}^{\mathcal{B}^{\text{op}}}$ is the same as the Reedy model structure on $(\mathcal{C}^{\text{op}})^\mathcal{B}$. Moreover if $\mathcal{B}_1, \mathcal{B}_2$ are Reedy categories, then the Reedy model structure on $(\mathcal{C}^{\mathcal{B}_2})^{\mathcal{B}_1}$ is the same as the Reedy model structure on $(\mathcal{C}^{\mathcal{B}_1})^{\mathcal{B}_2}$.

Remark 1.5.1.20. If \mathcal{C} is a model category, \mathcal{B} is a Reedy category, the colimit functor $\mathcal{C}^\mathcal{B} \rightarrow \mathcal{C}$ need not be left Quillen, and the limit functor $\mathcal{C}^\mathcal{B} \rightarrow \mathcal{C}$ need not be right Quillen.

Lemma 1.5.1.21 (The Cube Lemma). Suppose \mathcal{C} is a model category, P_i, Q_i, R_i ($i = 0, 1$) are cofibrant objects in \mathcal{C} , $f_i : P_i \rightarrow Q_i, g_i : P_i \rightarrow R_i$ ($i = 0, 1$) are morphisms in \mathcal{C} , such that f_0, f_1 are cofibrations. Define $S_i = Q_i \amalg_{P_i} R_i$ ($i = 0, 1$). Suppose more there are morphisms $p : P_0 \rightarrow P_1, q : Q_0 \rightarrow Q_1, r : R_0 \rightarrow R_1$, which induce a morphism $s : S_0 \rightarrow S_1$. If p, q, r are weak equivalences, then s is a weak equivalence; if $p, (f_1, q) : P_1 \amalg_{P_0} Q_0 \rightarrow Q_1, r$ are cofibrations, then s is a cofibration.

Definition 1.5.1.22. Suppose \mathcal{C} is a model category, A, B are objects in \mathcal{C} .

- i) A **cosimplicial frame** on A is a factorization $l^\bullet A \rightarrow A^* \rightarrow r^\bullet A$ in the category \mathcal{C}^Δ , such that the map $l^\bullet A \rightarrow A^*$ is a cofibration, the map $A^* \rightarrow r^\bullet A$ is a weak equivalence, and the factorization restricts to identity on degree 0. Dually, a **simplicial frame** on A is a factorization $l_\bullet A \rightarrow A_* \rightarrow r_\bullet A$ in the category $\mathcal{C}^{\Delta^{\text{op}}}$, such that the map $l_\bullet A \rightarrow A_*$ is a weak equivalence, the map $A_* \rightarrow r_\bullet A$ is a fibration, and the factorization restricts to identity on degree 0.
- ii) Suppose A^*, B^* are cosimplicial frames on A, B , respectively. A **map of cosimplicial frames** between A^*, B^* is simply a morphism $A^* \rightarrow B^*$ in the category \mathcal{C}^Δ . Given a map $f : A \rightarrow B$, a **map of cosimplicial frames over f** (or A , if f is the identity map on A) is a morphism $A^* \rightarrow B^*$ in the category \mathcal{C}^Δ making the following diagram commutative:

$$\begin{array}{ccccc} l^\bullet A & \longrightarrow & A^* & \longrightarrow & r^\bullet A \\ \downarrow l^\bullet f & & \downarrow & & \downarrow r^\bullet f \\ l^\bullet B & \longrightarrow & B^* & \longrightarrow & r^\bullet B \end{array}$$

Dually, suppose A_*, B_* are simplicial frames on A, B , respectively. A **map of simplicial frames** between A_*, B_* is simply a morphism $A_* \rightarrow B_*$ in the category $\mathcal{C}^{\Delta^{\text{op}}}$. Given a map $f : A \rightarrow B$, a **map of simplicial frames over f** (or A , if f is the identity map on A) is a morphism $A_* \rightarrow B_*$ in the category $\mathcal{C}^{\Delta^{\text{op}}}$ making the following diagram commutative:

$$\begin{array}{ccccc} l_\bullet A & \longrightarrow & A_* & \longrightarrow & r_\bullet A \\ \downarrow l_\bullet f & & \downarrow & & \downarrow r_\bullet f \\ l_\bullet B & \longrightarrow & B_* & \longrightarrow & r_\bullet B \end{array}$$

iii) A **left framing** on \mathcal{C} is a functor $-^* : \mathcal{C} \rightarrow \mathcal{C}^\Delta$, such that $-^*[0]$ is naturally isomorphic to the identity functor on \mathcal{C} , and A^* is a cosimplicial frame on A whenever A is cofibrant. Dually, a **right framing** on \mathcal{C} is a functor $-_* : \mathcal{C} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$, such that $-_*[0]$ is naturally isomorphic to the identity functor on \mathcal{C} , and A_* is a simplicial frame on A whenever A is fibrant.

iv) A **framing** on \mathcal{C} is a left framing together with a right framing.

Lemma 1.5.1.23. *Suppose \mathcal{C} is a model category, A is an object in \mathcal{C} . A cosimplicial frame on A is precisely a cosimplicial object A^* such that $A^*[0]$ is isomorphic to A , the induced map $L_n A^* \rightarrow A^*[n]$ is a cofibration for any positive n , and the induced map $A^*[n] \rightarrow A^*[0]$ is a weak equivalence for any n . Dually a simplicial frame on A is precisely a simplicial object A_* such that $A_*[0]$ is isomorphic to A , the induced map $A_*[0] \rightarrow A_*[n]$ is a weak equivalence for any n , and the induced map $A_*[n] \rightarrow M_n A_*$ is a fibration for any positive n .*

Lemma 1.5.1.24. *Any cosimplicial frame of a cofibrant object is cofibrant, and any simplicial frame of a fibrant object is fibrant.*

Lemma 1.5.1.25. *Any left framing on \mathcal{C} is a right framing on \mathcal{C}^{op} , and any right framing on \mathcal{C} is a left framing on \mathcal{C}^{op} .*

Theorem 1.5.1.26. *Suppose \mathcal{C} is a model category. Define $-^\circ : \mathcal{C} \rightarrow \mathcal{C}^\Delta$ by induction, with $-^\circ[0]$ being the identity functor, and $-^\circ[n] = c \circ \alpha(L_n -^\circ \rightarrow M_n -^\circ)$. Then for any $A \in \mathcal{C}$, A° is a cosimplicial frame on A , thus $-^\circ$ is a left framing on \mathcal{C} . Dually, define $-_\circ : \mathcal{C} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$ by induction, with $-_\circ[0]$ being the identity functor, and $-_\circ[n] = d \circ \delta(L_n -_\circ \rightarrow M_n -_\circ)$. Then for any $A \in \mathcal{C}$, A_\circ is a simplicial frame on A , thus $-_\circ$ is a right framing on \mathcal{C} .*

Proposition 1.5.1.27. *Suppose \mathcal{C} is a simplicial model category. Then $- \otimes \Delta[-] : \mathcal{C} \rightarrow \mathcal{C}^\Delta$ is a left framing on \mathcal{C} , and $-\Delta[-] : \mathcal{C} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$ is a right framing on \mathcal{C} .*

Remark 1.5.1.28. Suppose \mathcal{C} is a simplicial model category. If A is not cofibrant, then $A \otimes \Delta[-]$ may not be a cosimplicial frame on A . Dually if A is not fibrant, then $A \Delta[-]$ may not be a simplicial frame on A . Also, the functors $- \otimes \Delta[-], -^\circ$ are usually not the same, and the functors $-\Delta[-], -_\circ$ are usually not the same.

Definition 1.5.1.29. Suppose \mathcal{C} is a model category. By Proposition 1.3.1.23, $-^\circ$ induces functors $\mathcal{C} \times \underline{\mathbf{SSet}} \rightarrow \mathcal{C}, (A, K) \mapsto A^\circ \otimes K$, and $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \underline{\mathbf{SSet}}, (A, Y) \mapsto \text{Hom}_{\mathcal{C}}(A^\circ, Y)$. We denote $A^\circ \otimes K$ by $A \otimes K$ (if there is no confusion), $\text{Hom}_{\mathcal{C}}(A^\circ, Y)$ by $\text{Map}(A, Y)_l$, and call $(A, Y) \mapsto \text{Map}(A, Y)_l$ the **left function complex functor**. Dually by Proposition 1.3.1.24, $-_\circ$ induces functors $\mathcal{C} \times \underline{\mathbf{SSet}}^{\text{op}} \rightarrow \mathcal{C}, (A, K) \mapsto A_\circ^K$, and $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \underline{\mathbf{SSet}}, (A, Y) \mapsto \text{Hom}_{\mathcal{C}}(A, Y_\circ)$. We denote A_\circ^K by A^K , $\text{Hom}_{\mathcal{C}}(A, Y_\circ)$ by $\text{Map}(A, Y)_r$, and call $(A, Y) \mapsto \text{Map}(A, Y)_r$ the **right function complex functor**.

Remark 1.5.1.30. $A \otimes -$ is a left adjoint, but $- \otimes K$ need not be.

1.5.2 Homotopy Categories as $\text{Ho } \underline{\mathbf{SSet}}$ -Modules

Proposition 1.5.2.1. *Suppose \mathcal{C} is a model category, and $g : K \rightarrow L$ is a cofibration in $\underline{\mathbf{SSet}}$. If $f : A^\bullet \rightarrow B^\bullet$ is a cofibration in \mathcal{C}^Δ , then $f \square g : A^\bullet \otimes L \amalg_{A^\bullet \otimes K} B^\bullet \otimes K \rightarrow B^\bullet \otimes L$ is a cofibration that is trivial if f is. Dually if $p : X_\bullet \rightarrow Y_\bullet$ is a fibration in $\mathcal{C}^{\Delta^{\text{op}}}$, then $p_\square^q : X_\bullet^L \rightarrow Y_\bullet^L \times_{Y_\bullet^K} X_\bullet^K$ is a fibration that is trivial if p is.*

Corollary 1.5.2.2. *Suppose \mathcal{C} is a model category, and K is a simplicial set. Then the functor $- \otimes K : \mathcal{C}^\Delta \rightarrow \mathcal{C}$ preserves cofibrations and trivial cofibrations, and the functor $-^K : \mathcal{C}^{\Delta^{\text{op}}} \rightarrow \mathcal{C}$ preserves fibrations and trivial fibrations.*

Proposition 1.5.2.3. Suppose \mathcal{C} is a model category, and $g : K \rightarrow L$ is a trivial cofibration in $\underline{\mathbf{SSet}}$. If $f : A^* \rightarrow B^*$ is a cofibration of cosimplicial frames on cofibrant objects, then $f \square g : A^* \otimes L \amalg_{A^* \otimes K} B^* \otimes K \rightarrow B^* \otimes L$ is a trivial cofibration. Dually if $p : X_* \rightarrow Y_*$ is a fibration of simplicial frames on fibrant objects, then $p \square^g : X_*^L \rightarrow Y_*^L \times_{Y_*^K} X_*^K$ is a trivial fibration.

Corollary 1.5.2.4. Suppose \mathcal{C} is a model category, A is cofibrant, Y is fibrant. Then for any cosimplicial frame A^* on A , the adjunction $(A^* \otimes -, \text{Hom}_{\mathcal{C}}(A^*, -)) : \underline{\mathbf{SSet}} \rightarrow \mathcal{C}$ is a Quillen adjunction. In particular, the adjunction $(A \otimes -, \text{Map}(A, -)_l) : \underline{\mathbf{SSet}} \rightarrow \mathcal{C}$ is a Quillen adjunction. Dually for any simplicial frame Y_* on Y , the adjunction $(Y_*^-, \text{Hom}_{\mathcal{C}}(-, Y_*)) : \underline{\mathbf{SSet}} \rightarrow \mathcal{C}^{\text{op}}$ is a Quillen adjunction. In particular, the adjunction $(Y^-, \text{Map}(-, Y)_r) : \underline{\mathbf{SSet}} \rightarrow \mathcal{C}^{\text{op}}$ is a Quillen adjunction.

Corollary 1.5.2.5. Suppose \mathcal{C} is a model category, and K is a simplicial set. If $A^* \rightarrow B^*$ is a map of cosimplicial frames on cofibrant objects that is a weak equivalence in degree 0, then the map $A^* \otimes K \rightarrow B^* \otimes K$ is a weak equivalence. Dually if $X_* \rightarrow Y_*$ is a map of simplicial frames on fibrant objects that is a weak equivalence in degree 0, then the map $X_*^K \rightarrow Y_*^K$ is a weak equivalence.

Corollary 1.5.2.6. Suppose \mathcal{C} is a model category, and K is a simplicial set. Then the functor $-\otimes K : \mathcal{C} \rightarrow \mathcal{C}$ preserves weak equivalences between cofibrant objects, and the functor $-^K : \mathcal{C} \rightarrow \mathcal{C}$ preserves weak equivalences between fibrant objects.

Definition 1.5.2.7. We call elements in the category $\underline{\mathbf{SSet}}^{\Delta^{\text{op}}}$ **bisimplicial sets**. There is a functor $\underline{\mathbf{SSet}}^{\Delta^{\text{op}}} \rightarrow \underline{\mathbf{SSet}}$ induced by the diagonal functor $\Delta \rightarrow \Delta \times \Delta$. We call this functor diag .

Lemma 1.5.2.8. diag preserves colimits. Furthermore every bisimplicial set is cofibrant with respect to the Reedy model structure.

Proposition 1.5.2.9. Suppose \mathcal{C} is a category with all small colimits and limits. Define the functor $F_n : \mathcal{C} \rightarrow \mathcal{C}^{\Delta^{\text{op}}}$ with $(F_n K)_m = \amalg_{\Delta[n]_m} K$ and the simplicial structure given by the simplicial structure of $\Delta[n]$ for any $K \in \mathcal{C}$. Then F_n is left adjoint to $-[n]$. Furthermore for any $X \in \mathcal{C}^{\Delta^{\text{op}}}$, X is the coequalizer of the following diagram:

$$\coprod_{[k] \rightarrow [n]} F_k X_n \begin{array}{c} \xrightarrow{F_k X_n \rightarrow F_n X_n, \text{ induced by the map } \Delta[k] \rightarrow \Delta[n]} \\ \xrightarrow{F_k X_n \rightarrow F_k X_k, \text{ induced by the map } X_n \rightarrow X_k} \end{array} \coprod_k F_k X_k.$$

Proposition 1.5.2.10. For any simplicial set K , we have $\text{diag } F_n K = K \otimes \Delta[n]$. Therefore diag is left adjoint to the functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{SSet}}^{\Delta^{\text{op}}}, K \mapsto K^{\Delta[-]}$.

Lemma 1.5.2.11. The functor $K \mapsto K^{\Delta[-]}$ preserves fibrations and trivial fibrations, hence the adjunction $(\text{diag}, K \mapsto K^{\Delta[-]})$ is a Quillen adjunction.

Proposition 1.5.2.12. Suppose X is a bisimplicial set such that for any map $[k] \rightarrow [n]$ in Δ the induced map $X[n] \rightarrow X[k]$ is a weak equivalence. Then the map $X[0] \rightarrow \text{diag } X$ is a weak equivalence.

Proposition 1.5.2.13. Suppose \mathcal{C} is a model category, A^* is a cosimplicial frame on a cofibrant object, Y_* is a simplicial frame on a fibrant object. Then there exists weak equivalences

$$\text{Hom}_{\mathcal{C}}(A^*, Y) \rightarrow \text{diag } \text{Hom}_{\mathcal{C}}(A^*, Y_*) \leftarrow \text{Hom}_{\mathcal{C}}(A, Y_*).$$

Corollary 1.5.2.14. Suppose \mathcal{C} is a model category, A is cofibrant, Y is fibrant. Then $\text{Map}(-, Y)_l$ preserves weak equivalences between cofibrant objects, and $\text{Map}(A, -)_r$ preserves weak equivalences between fibrant objects.

Theorem 1.5.2.15. Suppose \mathcal{C} is a model category. Then the left derived functors of $- \otimes - : \mathcal{C} \times \underline{\mathbf{SSet}} \rightarrow \mathcal{C}$ and $-^- : \mathcal{C}^{\text{op}} \times \underline{\mathbf{SSet}} \rightarrow \mathcal{C}^{\text{op}}$ exist, which will be denoted $- \otimes^L -$ and $-^-_R$. The right derived functors of $\text{Map}(-, -)_l, \text{Map}(-, -)_r : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \underline{\mathbf{SSet}}$ exist, which will be denoted $R\text{Map}(-, -)_l, R\text{Map}(-, -)_r$, and are naturally isomorphic. There exist natural isomorphisms

$$[X \otimes^L K, Y] \cong [K, R\text{Map}(X, Y)_l] \cong [K, R\text{Map}(X, Y)_r] \cong [Y_R^K, X],$$

constructing an adjunction of 2 variables $\text{Ho } \mathcal{C} \times \text{Ho } \underline{\mathbf{SSet}} \rightarrow \text{Ho } \mathcal{C}$. Finally $- \otimes^L \Delta[0]$ is naturally isomorphic to the identity.

Lemma 1.5.2.16. Suppose \mathcal{C} is a model category, $f : A \rightarrow B$ is a morphism in \mathcal{C} . Then for any cosimplicial frame A^* on A , there exists a map $A^* \rightarrow B^\circ$ of cosimplicial frames over f . Dually for any simplicial frame B_* on B , there exists a map $A_\circ \rightarrow B_*$ of simplicial frames over f .

Lemma 1.5.2.17. Suppose \mathcal{C} is a model category, A^*, B^* are cosimplicial frames on cofibrant objects A, B , respectively. Suppose $f, g : A^* \rightarrow B^*$ are two maps of cosimplicial frames that agree on degree 0. Then the natural transformation $A^* \otimes^L - \rightarrow B^* \otimes^L -$ induced by f and g are equal.

Theorem 1.5.2.18. Suppose \mathcal{C} is a model category. Then for any $A \in \mathcal{C}$ and $K \in \underline{\mathbf{SSet}}$, $A \otimes (K \times \Delta[-])$ is a cosimplicial frame on $A \otimes K$ whenever A is cofibrant, which induces a weak equivalence

$$A \otimes (K \times L) \cong (A \otimes (K \times \Delta[-])) \otimes L \rightarrow (A \otimes K) \otimes L$$

that is natural on L . Define a_{AKL} to be the isomorphism in $\text{Ho } \mathcal{C}$:

$$\begin{aligned} (A \otimes^L K) \otimes^L L &= Q(QA \otimes QK) \otimes QL \xrightarrow{q_{QA \otimes QK} \otimes QL} (QA \otimes QK) \otimes QL \\ &\rightarrow QA \otimes (QK \times QL) \xrightarrow{QA \otimes q_{QK \times QL}^{-1}} QA \otimes Q(QK \times QL) = A \otimes^L (K \times^L L), \end{aligned}$$

then a is natural in A, K, L , and

$$(\text{Ho } \mathcal{C}, - \otimes^L -, R\text{Map}(-, -), -^-_R, a, r : - \otimes^L \Delta[0] \cong \mathbb{1})$$

gives $\text{Ho } \mathcal{C}$ a closed $\text{Ho } \underline{\mathbf{SSet}}$ -module structure.

Lemma 1.5.2.19. Suppose \mathcal{C}, \mathcal{D} are model categories, $(F, G) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction, A^* is a cosimplicial frame on a cofibrant object A in \mathcal{C} , Y_* is a simplicial frame on a fibrant object Y in \mathcal{D} . Then FA^* is a cosimplicial frame on FA and GY_* is a simplicial frame on GY .

Theorem 1.5.2.20. Suppose \mathcal{C}, \mathcal{D} are model categories, $(F, G, \varphi) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction. Then for any $A \in \mathcal{C}$, there is a weak equivalence

$$F(A \otimes K) \cong F(A^\circ) \otimes K \rightarrow FA \otimes K$$

whenever A is cofibrant and is natural on K . Define μ_{AK} to be the isomorphism in $\text{Ho } \mathcal{C}$:

$$\begin{aligned} (LFA) \otimes^L K &= QFQA \otimes QK \xrightarrow{q_{FQA}^{-1} \otimes QK} FQA \otimes QK \\ &\rightarrow F(QA \otimes QK) \xrightarrow{Fq_{QA \otimes QK}} FQ(QA \otimes QK) = (LF)(A \otimes^L K), \end{aligned}$$

then μ is natural in A, K , and

$$(LF, RG, \text{Ho } \varphi, \mu)$$

gives LF a closed $\text{Ho } \underline{\mathbf{SSet}}$ -module functor structure.

Theorem 1.5.2.21. *The pseudo-2-functor $\text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{Cat}}_{ad}$ lifts to a pseudo-2-functor*

$$\text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{CloMod}}_{\text{Ho } \underline{\text{SSet}}}$$

which is given by

$$\begin{aligned} \mathcal{C} &\mapsto (\text{Ho } \mathcal{C}, - \otimes^L -, R \text{Map}(-, -), -\bar{R}, a, r), \\ (F, G, \varphi) &\mapsto (LF, RG, \text{Ho } \varphi, \mu), \\ \tau &\mapsto L\tau. \end{aligned}$$

Moreover $\text{Ho} \circ (-^{\text{op}}) = (-^{\text{op}}) \circ \text{Ho} : \underline{\text{Model}} \rightarrow \underline{\text{CloMod}}_{\text{Ho } \underline{\text{SSet}}}$, and the identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\text{Model}}_{\underline{\text{SSet}}} \xrightarrow{\text{Ho}} \underline{\text{CloMod}}_{\text{Ho } \underline{\text{SSet}}}$$

and

$$\underline{\text{Model}}_{\underline{\text{SSet}}} \rightarrow \underline{\text{Model}} \xrightarrow{\text{Ho}} \underline{\text{CloMod}}_{\text{Ho } \underline{\text{SSet}}}.$$

Proposition 1.5.2.22. *Suppose \mathcal{C} is a model category. Then for any $X \in \mathcal{C}$, $X \otimes^L - : \text{Ho } \underline{\text{SSet}} \rightarrow \text{Ho } \mathcal{C}$ is a closed $\text{Ho } \underline{\text{SSet}}$ -module functor, where the structure map μ_{KL} is given by a_{XKL} . On the other hand, for any $K \in \underline{\text{SSet}}$, define μ_{XL} to be the natural isomorphism*

$$(X \otimes^L K) \otimes^L L \rightarrow X \otimes^L (K \times^L L) \rightarrow X \otimes^L (L \times^L K) \rightarrow (X \otimes^L L) \otimes^L K,$$

then $- \otimes^L K$ together with the structure map given above is a closed $\text{Ho } \underline{\text{SSet}}$ -module functor.

Proposition 1.5.2.23. *Suppose \mathcal{C} is a simplicial model category. Then for any $K \in \underline{\text{SSet}}$, the closed $\text{Ho } \underline{\text{SSet}}$ -module functors $- \otimes^L K$ and $- \circ \otimes^L K$ share the same structure map μ .*

Proposition 1.5.2.24. *Suppose \mathcal{C} is a model category. Then for any $X \in \mathcal{C}$, $K, L \in \underline{\text{SSet}}$, the associativity isomorphism is a $\text{Ho } \underline{\text{SSet}}$ -module natural transformation in the following cases:*

$$\begin{aligned} (X \otimes^L K) \otimes^L - &\rightarrow X \otimes^L (K \times^L -), \\ (X \otimes^L -) \otimes^L L &\rightarrow X \otimes^L (- \times^L L), \\ (- \otimes^L K) \otimes^L L &\rightarrow - \otimes^L (K \times^L L). \end{aligned}$$

Proposition 1.5.2.25. *Suppose \mathcal{C} is a monoidal model category. Then for any $X, Y, Z \in \mathcal{C}$, $X \otimes^L -, - \otimes^L Y : \text{Ho } \mathcal{C} \rightarrow \text{Ho } \mathcal{C}$ are closed $\text{Ho } \underline{\text{SSet}}$ -module functors, constructing following natural isomorphisms for $K \in \underline{\text{SSet}}$:*

$$m_{XYK}^l : (X \otimes^L Y) \otimes^L K \rightarrow X \otimes^L (Y \otimes^L K)$$

and

$$m_{XYK}^r : (X \otimes^L Y) \otimes^L K \rightarrow (X \otimes^L K) \otimes^L Y.$$

Furthermore

$$\begin{aligned} l : S \otimes^L - &\rightarrow \mathbb{1}, \\ r : - \otimes^L S &\rightarrow \mathbb{1}, \\ a_{XY-} : (X \otimes^L Y) \otimes^L - &\rightarrow X \otimes^L (Y \otimes^L -), \\ a_{X-Z} : (X \otimes^L -) \otimes^L Z &\rightarrow X \otimes^L (- \otimes^L Z), \\ a_{-YZ} : (- \otimes^L Y) \otimes^L Z &\rightarrow - \otimes^L (Y \otimes^L Z) \end{aligned}$$

are $\text{Ho } \underline{\text{SSet}}$ -module natural transformations.

Theorem 1.5.2.26. *The pseudo-2-functor $\text{Ho} : \underline{\text{MonModel}} \rightarrow \underline{\text{CloMon}}$ lifts to a pseudo-2-functor*

$$\text{Ho} : \underline{\text{MonModel}} \rightarrow \underline{\text{CloAlg}}_{\text{Ho } \underline{\text{SSet}}},$$

where a monoidal model category (\mathcal{C}, \otimes) is mapped to $(\text{Ho } \mathcal{C}, \otimes^L, (i, \mu, \alpha))$, where $i : \text{Ho } \underline{\text{SSet}} \rightarrow \text{Ho } \mathcal{C}$ is given by $K \mapsto S \otimes^L K$, μ is given by

$$\begin{aligned} iK \otimes^L iL &= (S \otimes^L K) \otimes^L (S \otimes^L L) \xrightarrow{(m^i_{(S \otimes^L K)SL})^{-1}} ((S \otimes^L K) \otimes^L S) \otimes^L L \\ &\xrightarrow{r_{S \otimes^L K \otimes^L L}} (S \otimes^L K) \otimes^L L \xrightarrow{a_{SKL}} S \otimes^L (K \times^L L) = i(K \times^L L), \end{aligned}$$

and α is given by

$$i(\Delta[0]) = S \otimes^L \Delta[0] \xrightarrow{r_S} S;$$

a monoidal Quillen adjunction $(F, G) : \mathcal{C} \rightarrow \mathcal{D}$ is mapped to (LF, RG, ρ) , where ρ is given by

$$(LF)i_{\mathcal{C}}K = (LF)(S \otimes^L K) \xrightarrow{\mu_{SK}^{-1}} (LFS) \otimes^L K \xrightarrow{\alpha_{S \otimes^L K}} S \otimes^L K = i_{\mathcal{D}}K;$$

a monoidal natural transformation τ is mapped to $L\tau$. Furthermore, the identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\text{MonModel}}_{\underline{\text{SSet}}} \xrightarrow{\text{Ho}} \underline{\text{CloAlg}}_{\text{Ho } \underline{\text{SSet}}}$$

and

$$\underline{\text{MonModel}}_{\underline{\text{SSet}}} \rightarrow \underline{\text{MonModel}} \xrightarrow{\text{Ho}} \underline{\text{CloAlg}}_{\text{Ho } \underline{\text{SSet}}}.$$

Theorem 1.5.2.27. *Suppose \mathcal{C} is a monoidal model category. Then for any $K \in \underline{\text{SSet}}$, the natural isomorphism*

$$X \otimes^L (S \otimes^L K) \xrightarrow{(m^i_{XSK})^{-1}} (X \otimes^L S) \otimes^L K \xrightarrow{r_{X \otimes^L K}} X \otimes^L K$$

is a $\text{Ho } \underline{\text{SSet}}$ -module natural transformation between $\text{Ho } \underline{\text{SSet}}$ -module functors $- \otimes^L (S \otimes^L K)$ and $- \otimes^L K$.²

Theorem 1.5.2.28. *The pseudo-2-functor*

$$\text{Ho} : \underline{\text{SymMonModel}} \rightarrow \underline{\text{CloSymMon}}$$

lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{SymMonModel}} \rightarrow \underline{\text{CloSymAlg}}_{\text{Ho } \underline{\text{SSet}}},$$

where the structures are given in Theorem 1.5.2.26. The identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\text{SymMonModel}}_{\underline{\text{SSet}}} \xrightarrow{\text{Ho}} \underline{\text{CloSymAlg}}_{\text{Ho } \underline{\text{SSet}}}$$

and

$$\underline{\text{SymMonModel}}_{\underline{\text{SSet}}} \rightarrow \underline{\text{SymMonModel}} \xrightarrow{\text{Ho}} \underline{\text{CloSymAlg}}_{\text{Ho } \underline{\text{SSet}}}.$$

²This theorem remains unproved in class. Its proof is in Corollaire 6.7 of *Propriétés universelles et extensions de Kan dérivées*.

Definition 1.5.2.29. Suppose \mathcal{C} is a pointed model category. By Proposition 1.3.1.25, $-^\circ$ induces functors $\mathcal{C} \times \underline{\mathbf{SSet}}_* \rightarrow \mathcal{C}$, $(A, K) \mapsto A^\circ \wedge K$, and $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \underline{\mathbf{SSet}}_*$, $(A, Y) \mapsto \text{Hom}_{\mathcal{C}*}(A^\circ, Y)$. We denote $A^\circ \wedge K$ by $A \wedge K$ (if there is no confusion), $\text{Hom}_{\mathcal{C}*} \mathcal{C}(A^\circ, Y)$ by $\text{Map}_*(A, Y)_l$. Dually $-_\circ$ induces functors $\mathcal{C} \times \underline{\mathbf{SSet}}_*^{\text{op}} \rightarrow \mathcal{C}$, $(A, K) \mapsto A_{\circ*}^K$, and $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \underline{\mathbf{SSet}}_*$, $(A, Y) \mapsto \text{Hom}_{\mathcal{C}*}(A, Y_\circ)$. We denote $A_{\circ*}^K$ by A_*^K , $\text{Hom}_{\mathcal{C}*}(A, Y_\circ)$ by $\text{Map}_*(A, Y)_r$.

Proposition 1.5.2.30. Suppose \mathcal{C} is a pointed model category, and $g : K \rightarrow L$ is a cofibration in $\underline{\mathbf{SSet}}_*$. If $f : A^\bullet \rightarrow B^\bullet$ is a cofibration in \mathcal{C}^Δ , then $f \square_* g : A^\bullet \wedge L \amalg_{A^\bullet \wedge K} B^\bullet \wedge K \rightarrow B^\bullet \wedge L$ is a cofibration that is trivial if f is. Dually if $p : X_\bullet \rightarrow Y_\bullet$ is a fibration in $\mathcal{C}^{\Delta^{\text{op}}}$, then $p_{\square*}^g : X_{**}^L \rightarrow Y_{**}^L \times_{Y_{**}^K} X_{**}^K$ is a fibration that is trivial if p is.

Proposition 1.5.2.31. Suppose \mathcal{C} is a pointed model category, and $g : K \rightarrow L$ is a trivial cofibration in $\underline{\mathbf{SSet}}_*$. If $f : A^* \rightarrow B^*$ is a cofibration of cosimplicial frames on cofibrant objects, then $f \square_* g : A^* \wedge L \amalg_{A^* \wedge K} B^* \wedge K \rightarrow B^* \wedge L$ is a trivial cofibration. Dually if $p : X_* \rightarrow Y_*$ is a fibration of simplicial frames on fibrant objects, then $p_{\square*}^g : X_{**}^L \rightarrow Y_{**}^L \times_{Y_{**}^K} X_{**}^K$ is a trivial fibration.

Proposition 1.5.2.32. Suppose \mathcal{C} is a pointed model category, A^* is a cosimplicial frame on a cofibrant object, Y_* is a simplicial frame on a fibrant object. Then there exists weak equivalences

$$\text{Hom}_{\mathcal{C}*}(A^*, Y) \rightarrow \text{diag } \text{Hom}_{\mathcal{C}*}(A^*, Y_*) \leftarrow \text{Hom}_{\mathcal{C}*}(A, Y_*).$$

Theorem 1.5.2.33. The pseudo-2-functor $\text{Ho} : \underline{\mathbf{Model}}_* \rightarrow \underline{\mathbf{Cat}}_{ad*}$ lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\mathbf{Model}}_* \rightarrow \underline{\mathbf{CloMod}}_{\text{Ho } \underline{\mathbf{SSet}}_*},$$

where the structures are similar to that given above. Moreover $\text{Ho} \circ (-^{\text{op}}) = (-^{\text{op}}) \circ \text{Ho} : \underline{\mathbf{Model}}_* \rightarrow \underline{\mathbf{CloMod}}_{\text{Ho } \underline{\mathbf{SSet}}_*}$, and the identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\mathbf{Model}}_{\underline{\mathbf{SSet}}_*} \xrightarrow{\text{Ho}} \underline{\mathbf{CloMod}}_{\text{Ho } \underline{\mathbf{SSet}}_*}$$

and

$$\underline{\mathbf{Model}}_{\underline{\mathbf{SSet}}_*} \rightarrow \underline{\mathbf{Model}}_* \xrightarrow{\text{Ho}} \underline{\mathbf{CloMod}}_{\text{Ho } \underline{\mathbf{SSet}}_*}.$$

Theorem 1.5.2.34. The pseudo-2-functor $\text{Ho} : \underline{\mathbf{MonModel}}_* \rightarrow \underline{\mathbf{CloMon}}_*$ lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\mathbf{MonModel}}_* \rightarrow \underline{\mathbf{CloAlg}}_{\text{Ho } \underline{\mathbf{SSet}}_*},$$

where the structures are similar to that given above. Furthermore, the identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\mathbf{MonModel}}_{\underline{\mathbf{SSet}}_*} \xrightarrow{\text{Ho}} \underline{\mathbf{CloAlg}}_{\text{Ho } \underline{\mathbf{SSet}}_*}$$

and

$$\underline{\mathbf{MonModel}}_{\underline{\mathbf{SSet}}_*} \rightarrow \underline{\mathbf{MonModel}}_* \xrightarrow{\text{Ho}} \underline{\mathbf{CloAlg}}_{\text{Ho } \underline{\mathbf{SSet}}_*}.$$

Theorem 1.5.2.35. Suppose \mathcal{C} is a pointed monoidal model category. Then for any $K \in \underline{\mathbf{SSet}}_*$, the natural isomorphism

$$X \wedge^L (S \wedge^L K) \xrightarrow{(m_{XSK}^L)^{-1}} (X \wedge^L S) \wedge^L K \xrightarrow{r_{X \wedge^L K}} X \wedge^L K$$

is a $\text{Ho } \underline{\mathbf{SSet}}_*$ -module natural transformation between $\text{Ho } \underline{\mathbf{SSet}}_*$ -module functors $- \wedge^L (S \wedge^L K)$ and $- \wedge^L K$.³

³This theorem remains unproved in class as well.

Theorem 1.5.2.36. *The pseudo-2-functor*

$$\mathrm{Ho} : \underline{\mathrm{SymMonModel}}_* \rightarrow \underline{\mathrm{CloSymMon}}_*$$

lifts to a pseudo-2-functor

$$\mathrm{Ho} : \underline{\mathrm{SymMonModel}}_* \rightarrow \underline{\mathrm{CloSymAlg}}_{\mathrm{Ho} \, \underline{\mathrm{SSet}}_*},$$

where the structures are similar to that given above. The identity functor gives a 2-natural isomorphism between the pseudo-2-functors

$$\underline{\mathrm{SymMonModel}}_{\underline{\mathrm{SSet}}_*} \xrightarrow{\mathrm{Ho}} \underline{\mathrm{CloSymAlg}}_{\mathrm{Ho} \, \underline{\mathrm{SSet}}_*}$$

and

$$\underline{\mathrm{SymMonModel}}_{\underline{\mathrm{SSet}}_*} \rightarrow \underline{\mathrm{SymMonModel}}_* \xrightarrow{\mathrm{Ho}} \underline{\mathrm{CloSymAlg}}_{\mathrm{Ho} \, \underline{\mathrm{SSet}}_*}.$$

1.6 Pointed Model Categories

1.6.1 The Suspension and Loop Functors

Definition 1.6.1.1. Suppose \mathcal{C} is a pointed model category. Define the **suspension functor** Σ to be the functor $-\wedge^L S^1 : \mathrm{Ho} \, \mathcal{C} \rightarrow \mathrm{Ho} \, \mathcal{C}$, and the **loop functor** Ω to be the functor $-_{*R}^{S^1} : \mathrm{Ho} \, \mathcal{C} \rightarrow \mathrm{Ho} \, \mathcal{C}$.

Lemma 1.6.1.2. (Σ, Ω) is an adjoint pair.

Lemma 1.6.1.3. Suppose \mathcal{C} is a pointed model category. For any $X \in \mathcal{C}$, ΣX is the cofiber of the map $QX \amalg QX \rightarrow \mathrm{Cyl}(QX)$, and is naturally isomorphic in $\mathrm{Ho} \, \mathcal{C}$ to the cofiber of the map $X \amalg X \rightarrow \mathrm{Cyl} X$ if X is cofibrant. Dually for any $X \in \mathcal{C}$, ΩX is the fiber of the map $\mathrm{Path}(RX) \rightarrow RX \times RX$, and is naturally isomorphic in $\mathrm{Ho} \, \mathcal{C}$ to the fiber of the map $\mathrm{Path} X \rightarrow X \times X$ if X is fibrant.

Proposition 1.6.1.4. Suppose \mathcal{C} is a pointed model category, A is cofibrant in \mathcal{C} , X is fibrant in \mathcal{C} . Then for any nonnegative integer t there exists natural isomorphisms

$$[A, \Omega^t X] \cong [\Sigma^t A, X] \cong \pi_t \mathcal{H}om_*(A, X)_l \cong \pi_t \mathcal{H}om_*(A, X)_r.$$

Lemma 1.6.1.5. Suppose \mathcal{C} is a pointed model category, A is cofibrant in \mathcal{C} , X is fibrant in \mathcal{C} . Then any element in $[\Sigma A, X]$ can be represented by a map $k : A \rightarrow X^{\Delta[1]} = \mathrm{Path} X$ such that $p_0 k = p_1 k = 0$, and two such maps k, k' represent the same element in $[\Sigma A, X]$ if and only if there exists a map $K : A \rightarrow X^{\Delta[1] \times \Delta[1]}$ such that $p_0^0 K = k, p_0^1 K = k', p_1^0 K = p_1^1 K = 0$, where

$$p_0^0, p_0^1, p_1^0, p_1^1 : X^{\Delta[1] \times \Delta[1]} \rightarrow X^{\Delta[1]}$$

are induced by the inclusions

$$\begin{aligned} i_0^0 &: \Delta[1] \times \{0\} \rightarrow \Delta[1] \times \Delta[1], \\ i_0^1 &: \Delta[1] \times \{1\} \rightarrow \Delta[1] \times \Delta[1], \\ i_1^0 &: \{0\} \times \Delta[1] \rightarrow \Delta[1] \times \Delta[1], \\ i_1^1 &: \{1\} \times \Delta[1] \rightarrow \Delta[1] \times \Delta[1], \end{aligned}$$

respectively. Dually any element in $[\Sigma A, X]$ can be represented by a map $h : \mathrm{Cyl} A = A \otimes \Delta[1] \rightarrow X$ such that $hi_0 = hi_1 = 0$, and two such maps h, h' represent the same element in $[\Sigma A, X]$ if and only if there exists a map $H : A \otimes (\Delta[1] \times \Delta[1]) \rightarrow X$ such that $Hi_0^0 = h, Hi_0^1 = h', Hi_1^0 = Hi_1^1 = 0$, where

$$i_0^0, i_0^1, i_1^0, i_1^1 : A \otimes \Delta[1] \rightarrow A \otimes (\Delta[1] \times \Delta[1])$$

are induced by the inclusions given above.

Lemma 1.6.1.6. Suppose \mathcal{C} is a pointed model category, A is cofibrant in \mathcal{C} , X is fibrant in \mathcal{C} . Then two maps $k, k' : A \rightarrow \text{Path } X$ with $p_0 k = p_1 k = p_0 k' = p_1 k' = 0$ represent the same element in $[\Sigma A, X]$ if and only if there exists a map $K : \text{Cyl } A \rightarrow \text{Path } X$ such that $p_0 K = p_1 K = 0, Ki_0 = k, Ki_1 = k'$. Dually two maps $h, h' : \text{Cyl } A \rightarrow X$ with $hi_0 = hi_1 = h'i_0 = h'i_1 = 0$ represent the same element in $[\Sigma A, X]$ if and only if there exists a map $H : \text{Cyl } A \rightarrow \text{Path } X$ such that $p_0 H = h, p_1 H = h', Hi_0 = Hi_1 = 0$.

Lemma 1.6.1.7. Suppose \mathcal{C} is a pointed model category, A is cofibrant in \mathcal{C} , X is fibrant in \mathcal{C} . Then two maps $k : A \rightarrow \text{Path } X$ with $p_0 k = p_1 k = 0$ and $h : \text{Cyl } A \rightarrow X$ with $hi_0 = hi_1 = 0$ represent the same element in $[\Sigma A, X]$ if and only if there exists a map $H : \text{Cyl } A \rightarrow \text{Path } X$ such that $p_0 H = h, p_1 H = 0, Hi_0 = k, Hi_1 = 0$. Such H is called a **correspondence** between h and k .

Corollary 1.6.1.8. Suppose \mathcal{C} is a pointed model category. Then for any X and positive integer t , $\Sigma^t X$ is naturally a cogroup object in $\text{Ho } \mathcal{C}$ that is naturally abelian if $t \geq 2$. Dually $\Omega^t X$ is naturally a group object in $\text{Ho } \mathcal{C}$ that is naturally abelian if $t \geq 2$.

Remark 1.6.1.9. Suppose \mathcal{C} is a pointed model category, A is cofibrant in \mathcal{C} , X is fibrant in \mathcal{C} . Then the group structure on $[\Sigma A, X]$ may be expressed as follows: for any $k, k' : A \rightarrow \text{Path } X$ with $p_0 k = p_1 k = p_0 k' = p_1 k' = 0$ representing elements in $[\Sigma A, X]$, define K to be a lift of the following diagram:

$$\begin{array}{ccc} * & \xrightarrow{\quad} & X^{\Delta[2]} \\ \downarrow & \nearrow K & \downarrow \\ A & \xrightarrow[k' \text{ on } X^{\Delta\{1,2\}}]{k \text{ on } X^{\Delta\{0,1\}}} & X^{\Lambda^1[2]} \end{array}$$

then $[k] \cdot [k'] = [X^{d_1} K]$; define H to be a lift of the following diagram:

$$\begin{array}{ccc} * & \xrightarrow{\quad} & X^{\Delta[2]} \\ \downarrow & \nearrow H & \downarrow \\ A & \xrightarrow[0 \text{ on } X^{\Delta\{0,2\}}]{k \text{ on } X^{\Delta\{0,1\}}} & X^{\Lambda^0[2]} \end{array}$$

then $[k]^{-1} = [X^{d_0} H]$.

Definition 1.6.1.10. Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, X is fibrant in \mathcal{C} . Then for any map $h : A \rightarrow \text{Path } X$ with $p_0 h = p_1 h = 0$ representing an element in $[\Sigma A, X]$ and map $u : C \rightarrow X$ representing an element in $[C, X]$, take α to be a lift of the following diagram:

$$\begin{array}{ccc} A & \xrightarrow{h} & \text{Path } X \\ f \downarrow & \nearrow \alpha & \downarrow p_0 \\ B & \xrightarrow{u} C & \xrightarrow{g} X \end{array}$$

and take w to be the map induced as follows:

$$\begin{array}{ccccc} A & \xrightarrow{h} & \text{Path } X & & \\ \downarrow & \searrow f & \nearrow \alpha & & \downarrow p_1 \\ & B & & & \\ & \downarrow g & & & \\ * & \longrightarrow & C & \xrightarrow{\quad w \quad} & X \end{array}$$

and define $[u] \odot [h] = [w]$. Dually suppose $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, and A is cofibrant in \mathcal{C} . Then for any map $h : \text{Cyl } A \rightarrow B$ with $hi_0 = hi_1 = 0$ representing an element in $[A, \Omega B]$ and map $u : A \rightarrow F$ representing an element in $[A, F]$, take α to be a lift of the following diagram:

$$\begin{array}{ccccc} A & \xrightarrow{u} & F & \xrightarrow{i} & E \\ i_0 \downarrow & & \alpha \nearrow & & \downarrow p \\ \text{Cyl } A & \xrightarrow{h} & & & B \end{array}$$

and take w to be the map induced as follows:

$$\begin{array}{ccccc} A & \xrightarrow{w} & F & \longrightarrow & * \\ i_1 \downarrow & & i \downarrow & & \downarrow \\ \text{Cyl } A & \xrightarrow{h} & E & \xrightarrow{p} & B \end{array}$$

and define $[u] \odot [h] = [w]$.

Lemma 1.6.1.11. *Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$. Then there exists cylinder objects B', C' for B, C , and maps $\text{Cyl } A \xrightarrow{f'} B' \xrightarrow{g'} C'$, such that f' is a cofibration with cofiber g' , the following diagram is commutative:*

$$\begin{array}{ccccc} A \amalg A & \xrightarrow{(i_0, i_1)} & \text{Cyl } A & \xrightarrow{\sigma} & A \\ f \amalg f \downarrow & & \downarrow f' & & \downarrow f \\ B \amalg B & \xrightarrow{(i_0, i_1)} & B' & \xrightarrow{\sigma} & B \\ g \amalg g \downarrow & & \downarrow g' & & \downarrow g \\ C \amalg C & \xrightarrow{(i_0, i_1)} & C' & \xrightarrow{\sigma} & C \end{array}$$

and the map $(B \amalg B) \amalg_{A \amalg A} \text{Cyl } A \rightarrow B'$ is a cofibration. Dually there is a statement for fibration of fibrant objects.

Lemma 1.6.1.12. *Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, X is fibrant in \mathcal{C} . Then the map $\odot : [C, X] \times [\Sigma A, X] \rightarrow [C, X]$ is well-defined. Dually suppose $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, and A is cofibrant in \mathcal{C} . Then the map $\odot : [A, F] \times [A, \Omega B] \rightarrow [A, F]$ is well-defined.*

Lemma 1.6.1.13. *Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, X is fibrant in \mathcal{C} . Then the map $\odot : [C, X] \times [\Sigma A, X] \rightarrow [C, X]$ is natural in X . Dually suppose $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, and A is cofibrant in \mathcal{C} . Then the map $\odot : [A, F] \times [A, \Omega B] \rightarrow [A, F]$ is natural in A .*

Theorem 1.6.1.14. *Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, X is fibrant in \mathcal{C} . Then the map $\odot : [C, X] \times [\Sigma A, X] \rightarrow [C, X]$ defines a right $[\Sigma A, X]$ -action on $[C, X]$, constructing a right ΣA -coaction on C in $\text{Ho } \mathcal{C}$. Dually suppose $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, and A is cofibrant in \mathcal{C} . Then the map $\odot : [A, F] \times [A, \Omega B] \rightarrow [A, F]$ defines a right $[A, \Omega B]$ -action on $[A, F]$, constructing a right ΩB -action on F in $\text{Ho } \mathcal{C}$.*

Proposition 1.6.1.15. Suppose \mathcal{C} is a pointed model category, $f_i : A_i \rightarrow B_i$ are cofibrations of cofibrant objects with cofibers $g_i : B_i \rightarrow C_i$ for $i = 1, 2$, $p : A_1 \rightarrow A_2$, $q : B_1 \rightarrow B_2$ are maps in \mathcal{C} such that $qf_1 = f_2p$. Then the induced map $r : C_1 \rightarrow C_2$ is Σp -equivariant in $\text{Ho } \mathcal{C}$, namely the following diagram is commutative in $\text{Ho } \mathcal{C}$:

$$\begin{array}{ccc} C_1 & \longrightarrow & C_1 \amalg \Sigma A_1 \\ r \downarrow & & \downarrow r \amalg \Sigma p \\ C_2 & \longrightarrow & C_2 \amalg \Sigma A_2 \end{array}$$

Dually there is a statement for fibrations of fibrant objects.

Definition 1.6.1.16. Suppose \mathcal{C} is a pointed model category. A **cofiber sequence** in $\text{Ho } \mathcal{C}$ is a diagram $X \rightarrow Y \rightarrow Z$ in $\text{Ho } \mathcal{C}$ together with a right ΣX -coaction on Z in $\text{Ho } \mathcal{C}$, such that there exists a commutative diagram in $\text{Ho } \mathcal{C}$:

$$\begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & Z \\ p \downarrow & & \downarrow q & & \downarrow r \\ A & \xrightarrow{f} & B & \xrightarrow{g} & C \end{array}$$

such that $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, p, q, r are isomorphisms and r is Σp -equivariant. Dually a **fiber sequence** in $\text{Ho } \mathcal{C}$ is a diagram $X \rightarrow Y \rightarrow Z$ in $\text{Ho } \mathcal{C}$ together with a right ΩZ -coaction on X in $\text{Ho } \mathcal{C}$, such that there exists a commutative diagram in $\text{Ho } \mathcal{C}$:

$$\begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & Z \\ s \downarrow & & \downarrow t & & \downarrow u \\ F & \xrightarrow{i} & E & \xrightarrow{p} & B \end{array}$$

such that $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, s, t, u are isomorphisms and s is Ωu -equivariant.

Lemma 1.6.1.17. Suppose \mathcal{C} is a pointed model category. For any cofiber sequence or fiber sequence $X \rightarrow Y \rightarrow Z$, the composite of the two maps is 0.

Definition 1.6.1.18. Suppose \mathcal{C} is a pointed model category. For any cofiber sequence $X \rightarrow Y \rightarrow Z$, define the **boundary map** $\partial : Z \rightarrow \Sigma X$ to be the composite

$$Z \xrightarrow{\text{coaction}} Z \amalg \Sigma X \xrightarrow{0 \amalg 1} \Sigma X.$$

Dually for any fiber sequence $X \rightarrow Y \rightarrow Z$, define the **boundary map** $\partial : \Omega Z \rightarrow X$ to be the composite

$$\Omega Z \xrightarrow{(0, 1)} X \times \Omega Z \xrightarrow{\text{action}} X.$$

Lemma 1.6.1.19. Suppose \mathcal{C} is a pointed model category. For any cofiber sequence $X \rightarrow Y \rightarrow Z$ and $\theta \in [\Sigma X, A]$, $\theta \partial = [0] \odot \theta$. Dually for any fiber sequence $X \rightarrow Y \rightarrow Z$ and $\theta \in [A, \Omega Z]$, $\partial \theta = [0] \odot \theta$.

1.6.2 Cofiber and Fiber Sequences

Proposition 1.6.2.1. Suppose \mathcal{C} is a pointed model category. Then any diagram isomorphic to a cofiber sequence is a cofiber sequence, and any diagram isomorphic to a fiber sequence is a fiber sequence.

Proposition 1.6.2.2. Suppose \mathcal{C} is a pointed model category. Then for any $X \in \mathcal{C}$, the diagram $* \rightarrow X \xrightarrow{1} X$ together with the trivial coaction of $\Sigma* = *$ on X is a cofiber sequence, and the diagram $X \xrightarrow{1} X \rightarrow *$ together with the trivial action of $\Omega* = *$ on X is a fiber sequence.

Proposition 1.6.2.3. Suppose \mathcal{C} is a pointed model category. Then for any $f : X \rightarrow Y \in \text{Ho } \mathcal{C}$, there exists a cofiber sequence $X \xrightarrow{f} Y \rightarrow Z$, and there exists a fiber sequence $W \rightarrow X \xrightarrow{f} Y$.

Proposition 1.6.2.4. Suppose \mathcal{C} is a pointed model category. For any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$, the sequence $Y \xrightarrow{g} Z \xrightarrow{\partial} \Sigma X$ is a cofiber sequence, where the coaction is given by

$$\Sigma X \xrightarrow{m} \Sigma X \amalg \Sigma X \xrightarrow{1 \amalg \Sigma f} \Sigma X \amalg \Sigma Y \xrightarrow{1 \amalg i} \Sigma X \amalg \Sigma Y,$$

where m is the comultiplication and i is the coinverse. Dually for any fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$, the sequence $\Omega Z \xrightarrow{\partial} X \xrightarrow{f} Y$ is a fiber sequence, where the action is given by

$$\Omega Z \times \Omega Y \xrightarrow{1 \times i} \Omega Z \times \Omega Y \xrightarrow{1 \times \Omega g} \Omega Z \times \Omega Z \xrightarrow{m} \Omega Z,$$

where m is the multiplication and i is the inverse.

Proposition 1.6.2.5. Suppose \mathcal{C} is a pointed model category. If we have a commutative diagram in $\text{Ho } \mathcal{C}$ as follows:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ \alpha \downarrow & & \beta \downarrow & & \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' \end{array}$$

where the two rows are cofiber sequences, then there exists a map $\gamma : Z \rightarrow Z'$ making the following diagram commutative:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \longrightarrow & Z \amalg \Sigma X \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow \gamma \amalg \Sigma \alpha \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & \longrightarrow & Z' \amalg \Sigma X' \end{array}$$

Dually if we have a commutative diagram in $\text{Ho } \mathcal{C}$ as follows:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ & & \beta \downarrow & & \downarrow \alpha \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' \end{array}$$

where the two rows are fiber sequences, then there exists a map $\gamma : X \rightarrow X'$ making the following diagram commutative:

$$\begin{array}{ccccccc} X \times \Omega Z & \longrightarrow & X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ \gamma \times \Omega \alpha \downarrow & & \downarrow \alpha & & \downarrow \beta & & \downarrow \alpha \\ X' \times \Omega Z' & \longrightarrow & X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' \end{array}$$

Proposition 1.6.2.6. Suppose \mathcal{C} is a pointed model category. If $v : X \rightarrow Y, u : Y \rightarrow Z$ are maps in $\text{Ho } \mathcal{C}$, then there exists cofiber sequences

$$X \xrightarrow{v} Y \xrightarrow{d} U, \quad X \xrightarrow{uv} Z \xrightarrow{a} V, \quad Y \xrightarrow{u} Z \xrightarrow{f} W, \quad U \xrightarrow{r} V \xrightarrow{s} W,$$

such that the following diagram is commutative:

$$\begin{array}{ccccccc}
X & \xrightarrow{v} & Y & \xrightarrow{d} & U & \longrightarrow & U \amalg \Sigma X \\
\downarrow \mathbb{1} & & \downarrow u & & \downarrow r & & \downarrow r \amalg \mathbb{1} \\
X & \xrightarrow{uv} & Z & \xrightarrow{a} & V & \longrightarrow & V \amalg \Sigma X \\
\downarrow v & & \downarrow \mathbb{1} & & \downarrow s & & \downarrow s \amalg \Sigma v \\
Y & \xrightarrow{u} & Z & \xrightarrow{f} & W & \longrightarrow & W \amalg \Sigma Y \\
& & & & \downarrow & \nearrow \mathbb{1} \amalg \Sigma d & \\
& & & & W \amalg \Sigma U & &
\end{array}$$

Dually there is a statement for fiber sequences.

Proposition 1.6.2.7. Suppose \mathcal{C} is a pointed model category, $X \xrightarrow{f} Y \xrightarrow{g} Z$ is a cofiber sequence, $X' \xrightarrow{i} Y' \xrightarrow{p} Z'$ is a fiber sequence. If we have a commutative diagram as follows:

$$\begin{array}{ccccccc}
X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{\partial} & \Sigma X \\
\alpha \downarrow & & \beta \downarrow & & & & \downarrow -\varphi^{-1}\alpha \\
\Omega Z' & \xrightarrow{\partial} & X' & \xrightarrow{i} & Y' & \xrightarrow{p} & Z'
\end{array}$$

where φ is the natural isomorphism $[\Sigma X, Z] \cong [X, \Omega Z]$ and the negative sign is taken with respect to the group structure of $[\Sigma X, Z]$, then there exists a fill-in map $\gamma : Z \rightarrow Y'$ making the diagram commutative. Dually if we have a commutative diagram as follows:

$$\begin{array}{ccccccc}
X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{\partial} & \Sigma X \\
-\varphi\delta \downarrow & & & & \downarrow \gamma & & \downarrow \delta \\
\Omega Z' & \xrightarrow{\partial} & X' & \xrightarrow{i} & Y' & \xrightarrow{p} & Z'
\end{array}$$

then there exists a fill-in map $\beta : Y \rightarrow X'$ making the diagram commutative.

Proposition 1.6.2.8. Suppose $(F, G) : \mathcal{C} \rightarrow \mathcal{D}$ is a Quillen adjunction between pointed model categories. Then $LF : \text{Ho } \mathcal{C} \rightarrow \text{Ho } \mathcal{D}$ maps any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ to a cofiber sequence $LF X \xrightarrow{LF f} LF Y \xrightarrow{LF g} LF Z$, where the coaction is given by

$$LF Z \rightarrow LF(Z \amalg \Sigma X) \cong LF Z \amalg LF \Sigma X \xrightarrow{\mathbb{1} \amalg \mu^{-1}} LF Z \amalg \Sigma LF X,$$

where μ is the structure isomorphism $(LF X) \wedge^L S^1 \cong LF(X \wedge^L S^1)$ of LF as a $\text{Ho } \underline{\mathbf{S}}\mathbf{Set}_*$ -module functor. Dually $RG : \text{Ho } \mathcal{D} \rightarrow \text{Ho } \mathcal{C}$ maps any fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ to a fiber sequence $RG X \xrightarrow{RG f} RG Y \xrightarrow{RG g} RG Z$, where the action is given by

$$RG X \times \Omega RG Z \xrightarrow{\mathbb{1} \times (\mu^{\text{op}})^{-1}} RG X \times RG \Omega Z \cong RG(X \times \Omega Z) \rightarrow RG X,$$

where μ^{op} is the dual natural transformation of μ .

Proposition 1.6.2.9. Suppose \mathcal{C} is a pointed model category. Then for any $A \in \text{Ho } \mathcal{C}$, $A \wedge^L -$ maps cofiber sequences in $\text{Ho } \underline{\mathbf{S}}\mathbf{Set}_*$ into cofiber sequences in $\text{Ho } \mathcal{C}$, A_{*R}^- maps cofiber sequences in $\text{Ho } \underline{\mathbf{S}}\mathbf{Set}_*$ into fiber sequences in $\text{Ho } \mathcal{C}$, $R\text{Map}_*(-, A)_l \cong R\text{Map}_*(-, A)_r$ maps fiber sequences in $\text{Ho } \mathcal{C}$ into fiber sequences in $\text{Ho } \underline{\mathbf{S}}\mathbf{Set}_*$, $R\text{Map}_*(A, -)_l \cong R\text{Map}_*(A, -)_r$ maps cofiber sequences in $\text{Ho } \mathcal{C}$ into fiber sequences in $\text{Ho } \underline{\mathbf{S}}\mathbf{Set}_*$.

Lemma 1.6.2.10. Suppose \mathcal{C} is a pointed model category, $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$. Take a cylinder object A' of A , and define $r : A' \rightarrow A''$ to be the cofiber of $A \amalg A \xrightarrow{(i_0, i_1)} A'$. Define \tilde{B} to be the pushout of the diagram:

$$\begin{array}{ccc} A \amalg A & \xrightarrow{(i_0, i_1)} & A' \\ f \amalg f \downarrow & & \downarrow \\ B \amalg B & \longrightarrow & \tilde{B} \end{array}$$

and factor the map $(f\sigma, 1 \amalg 1) : \tilde{B} \rightarrow B$ into a cofibration $\tilde{B} \rightarrow B'$ and a trivial fibration $\sigma : B' \rightarrow B$. Then B' is a cylinder object of B , and it induces a map $f' : A' \rightarrow B'$ commuting with i_0, i_1, σ . Let B_f to be the pushout of the diagram:

$$\begin{array}{ccc} A & \xrightarrow{i_0} & A' \\ f \downarrow & & \downarrow \\ B & \xrightarrow{f} & B_f \end{array}$$

Then the following diagram is a pushout square:

$$\begin{array}{ccc} A & \xrightarrow{i_1} & A' \\ jf \downarrow & & \downarrow \\ B_f & \longrightarrow & \tilde{B} \end{array}$$

Let \tilde{C} to be the pushout of the diagram:

$$\begin{array}{ccc} B_f & \xrightarrow{(i_0, f')} & B' \\ (g, r) \downarrow & & \downarrow k \\ Cof \amalg A'' & \xrightarrow{\pi} & \tilde{C} \end{array}$$

then π is a weak equivalence, and $ki_1f = 0 : A \rightarrow \tilde{C}$, which induces a map $i_1 := (ki_1, 0) : C \rightarrow \tilde{C}$. Take a map $t : A' \rightarrow \text{Cyl } A$ commuting with i_0, i_1, σ , and call the induced map $A'' \rightarrow A \wedge S^1$ also by t . Now if we define $\psi : C \rightarrow C \amalg \Sigma A$ to be the composite in $\text{Ho } \mathcal{C}$:

$$C \xrightarrow{i_1} \tilde{C} \xrightarrow{\pi^{-1}} C \amalg A'' \xrightarrow{1 \amalg t} C \amalg (A \wedge S^1) \cong C \amalg (QA \wedge QS^1) = C \amalg \Sigma A,$$

then ψ coincides with the right coaction given in Theorem 1.6.1.14. Dually there is a statement for fibration of fibrant objects.

Lemma 1.6.2.11. Suppose \mathcal{C} is a pointed model category. If $f : A \rightarrow B$ is a cofibration of cofibrant objects with cofiber $g : B \rightarrow C$, then there exists cosimplicial frames B^*, C^* for B, C and a cofibration $f^* : A^\circ \rightarrow B^*$ in \mathcal{C}^Δ covering f with cofiber $g^* : B^* \rightarrow C^*$ covering g . Dually if $p : E \rightarrow B$ is a fibration of fibrant objects with fiber $i : F \rightarrow E$, then there exists simplicial frames F_*, E_* for F, E and a fibration $p_* : E_* \rightarrow B_\circ$ in $\mathcal{C}^{\Delta^{\text{op}}}$ covering p with fiber $i_* : F_* \rightarrow E_*$ covering i .

Proposition 1.6.2.12. Suppose \mathcal{C}, \mathcal{D} are pointed model categories. Then any functor between \mathcal{C} and \mathcal{D} preserving cofibrations, weak equivalence between cofibrant objects and colimits preserves cofiber sequences, where the coaction is simply given by acting the functor on the map ψ given in Lemma 1.6.2.10, and any functor between \mathcal{C} and \mathcal{D} preserving fibrations, weak equivalence between fibrant objects and limits preserves fiber sequences, where the action is simply given by acting the functor on the map given in the dual statement in Lemma 1.6.2.10.

Proposition 1.6.2.13. *Suppose \mathcal{C} is a pointed model category. Then for any $K \in \underline{\mathbf{SSet}}_*$, $-\wedge^L K$ maps a cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in $\mathbf{Ho} \mathcal{C}$ into a cofiber sequence $X \wedge^L K \xrightarrow{f \wedge^L 1} Y \wedge^L K \xrightarrow{g \wedge^L 1} Z \wedge^L K$ in $\mathbf{Ho} \mathcal{C}$, where the coaction is given by*

$$\begin{aligned} Z \wedge^L K &\rightarrow (Z \amalg \Sigma X) \wedge^L K \\ &\cong Z \wedge^L K \amalg \Sigma X \wedge^L K \xrightarrow{1 \amalg \mu^{-1}} Z \wedge^L K \amalg \Sigma(X \wedge^L K), \end{aligned}$$

where μ is the structure isomorphism $(X \wedge^L K) \wedge^L S^1 \cong (X \wedge^L S^1) \wedge^L K$ of $-\wedge^L K$ as a $\mathbf{Ho} \underline{\mathbf{SSet}}_*$ -module functor. Dually $-^K_{*R}$ maps a fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in $\mathbf{Ho} \mathcal{C}$ into a fiber sequence $X^K_{*R} \xrightarrow{f^1_{*R}} Y^K_{*R} \xrightarrow{g^1_{*R}} Z^K_{*R}$ in $\mathbf{Ho} \mathcal{C}$, where the action is given by

$$Z^K_{*R} \times \Omega X^K_{*R} \xrightarrow{1 \times (\mu^{\text{op}})^{-1}} Z^K_{*R} \times (\Omega X)^K_{*R} \cong (Z \times \Omega X)^K_{*R} \rightarrow Z^K_{*R},$$

where μ^{op} is the dual natural transformation of μ .

1.6.3 Pre-Triangulated Categories

Definition 1.6.3.1. Suppose $(\mathcal{S}, - \wedge -, \text{Map}_*(-, -), -^{\text{op}}_*)$ is a closed $\mathbf{Ho} \underline{\mathbf{SSet}}_*$ -module. Define the **suspension functor** Σ to be the functor $- \wedge S^1 : \mathcal{S} \rightarrow \mathcal{S}$, and the **loop functor** Ω to be the functor $-^{S^1}_* : \mathcal{S} \rightarrow \mathcal{S}$.

Lemma 1.6.3.2. *Suppose \mathcal{S} is a closed $\mathbf{Ho} \underline{\mathbf{SSet}}_*$ -module. Then for any X and positive integer t , $\Sigma^t X$ is naturally a cogroup object in \mathcal{S} that is naturally abelian if $t \geq 2$. Dually $\Omega^t X$ is naturally a group object in \mathcal{S} that is naturally abelian if $t \geq 2$.*

Definition 1.6.3.3. Suppose \mathcal{S} is a closed $\mathbf{Ho} \underline{\mathbf{SSet}}_*$ -module. A **pre-triangulation** on \mathcal{S} is a collection of **cofiber sequences** or **left triangles**, and a collection of **fiber sequences** or **right triangles**, satisfying the following conditions:

- i) A cofiber sequence is a diagram of the form $X \xrightarrow{f} Y \xrightarrow{g} Z$ with a right ΣX -coaction on Z .
- ii) The dual statement of i) holds.
- iii) Any diagram isomorphic to a cofiber sequence is a cofiber sequence.
- iv) The dual statement of iii) holds.
- v) For any X , the diagram $* \rightarrow X \xrightarrow{1} X$ together with the trivial coaction of $\Sigma * = *$ on X is a cofiber sequence.
- vi) The dual statement of v) holds.
- vii) For any $f : X \rightarrow Y$, there exists a cofiber sequence $X \xrightarrow{f} Y \rightarrow Z$.
- viii) The dual statement of vii) holds.
- ix) For any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$, the sequence $Y \xrightarrow{g} Z \xrightarrow{\partial} \Sigma X$ is a cofiber sequence, where ∂ is the map $Z \rightarrow Z \amalg \Sigma X \xrightarrow{0 \amalg 1} \Sigma X$ and the coaction is given by

$$\Sigma X \xrightarrow{m} \Sigma X \amalg \Sigma X \xrightarrow{1 \amalg \Sigma f} \Sigma X \amalg \Sigma Y \xrightarrow{1 \amalg i} \Sigma X \amalg \Sigma Y,$$

where m is the comultiplication and i is the coinverse.

x) The dual statement of ix) holds.

xi) For any commutative diagram as follows:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ \alpha \downarrow & & \beta \downarrow & & \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' \end{array}$$

where the two rows are cofiber sequences, there exists a map $\gamma : Z \rightarrow Z'$ making the following diagram commutative:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \longrightarrow & Z \amalg \Sigma X \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow \gamma \amalg \Sigma \alpha \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & \longrightarrow & Z' \amalg \Sigma X' \end{array}$$

xii) The dual statement of xi) holds.

xiii) For any maps $v : X \rightarrow Y, u : Y \rightarrow Z$, there exists cofiber sequences

$$X \xrightarrow{v} Y \xrightarrow{d} U, \quad X \xrightarrow{uv} Z \xrightarrow{a} V, \quad Y \xrightarrow{u} Z \xrightarrow{f} W, \quad U \xrightarrow{r} V \xrightarrow{s} W,$$

such that the following diagram is commutative:

$$\begin{array}{ccccccc} X & \xrightarrow{v} & Y & \xrightarrow{d} & U & \longrightarrow & U \amalg \Sigma X \\ \mathbb{1} \downarrow & & u \downarrow & & r \downarrow & & \downarrow r \amalg \mathbb{1} \\ X & \xrightarrow{uv} & Z & \xrightarrow{a} & V & \longrightarrow & V \amalg \Sigma X \\ v \downarrow & & \mathbb{1} \downarrow & & s \downarrow & & \downarrow s \amalg \Sigma v \\ Y & \xrightarrow{u} & Z & \xrightarrow{f} & W & \longrightarrow & W \amalg \Sigma Y \\ & & & & \downarrow & \nearrow \mathbb{1} \amalg \Sigma d & \\ & & & & W \amalg \Sigma U & & \end{array}$$

xiv) The dual statement of xiii) holds.

xv) For any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ and fiber sequence $X' \xrightarrow{i} Y' \xrightarrow{p} Z'$ and a commutative diagram as follows:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{\partial} & \Sigma X \\ \alpha \downarrow & & \beta \downarrow & & & & \downarrow -\varphi^{-1}\alpha \\ \Omega Z' & \xrightarrow{\partial} & X' & \xrightarrow{i} & Y' & \xrightarrow{p} & Z' \end{array}$$

where φ is the natural isomorphism $[\Sigma X, Z] \cong [X, \Omega Z]$ and the negative sign is taken with respect to the group structure of $[\Sigma X, Z]$, there exists a fill-in map $\gamma : Z \rightarrow Y'$ making the diagram commutative.

xvi) The dual statement of xv) holds.

xvii) For any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in \mathcal{S} and $K \in \text{Ho } \underline{\mathbf{S}}\text{Set}_*$, the sequence $X \wedge K \xrightarrow{f \wedge \mathbb{1}} Y \wedge K \xrightarrow{g \wedge \mathbb{1}} Z \wedge K$ is a cofiber sequence, where the coaction is given by

$$\begin{aligned} Z \wedge K &\rightarrow (Z \amalg \Sigma X) \wedge K \\ &\cong Z \wedge K \amalg \Sigma X \wedge K \xrightarrow{\mathbb{1} \amalg \mu^{-1}} Z \wedge K \amalg \Sigma(X \wedge K), \end{aligned}$$

where μ is the structure isomorphism $(X \wedge K) \wedge S^1 \cong (X \wedge S^1) \wedge K$ of $-\wedge K$ as a $\text{Ho } \underline{\mathbf{SSet}}_*$ -module functor.

xviii) The dual statement of xvii) holds.

xix) For any cofiber sequence $K \xrightarrow{f} L \xrightarrow{g} M$ in $\text{Ho } \underline{\mathbf{SSet}}_*$ and $X \in \mathcal{S}$, the sequence $X \wedge K \xrightarrow{1 \wedge f} X \wedge L \xrightarrow{1 \wedge g} X \wedge M$ is a cofiber sequence, where the coaction is given by

$$\begin{aligned} X \wedge M &\rightarrow X \wedge (M \amalg \Sigma K) \\ &\cong X \wedge M \amalg X \wedge \Sigma K \xrightarrow{1 \amalg \mu^{-1}} X \wedge M \amalg \Sigma(X \wedge K), \end{aligned}$$

where μ is the structure isomorphism $(X \wedge S^1) \wedge K \cong (X \wedge K) \wedge S^1$ of $-\wedge S^1$ as a $\text{Ho } \underline{\mathbf{SSet}}_*$ -module functor.

xx) The dual statement of xix) holds.

xxi) For any fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in \mathcal{S} and $A \in \mathcal{S}$, the sequence $\text{Map}_*(A, X) \xrightarrow{\text{Map}_*(A, f)} \text{Map}_*(A, Y) \xrightarrow{\text{Map}_*(A, g)} \text{Map}_*(A, Z)$ is a fiber sequence, where the action is given by

$$\begin{aligned} \text{Map}_*(A, X) \times \Omega \text{Map}_*(A, Z) &\xrightarrow{1 \times \mu^{\text{op}}} \text{Map}_*(A, X) \times \text{Map}_*(A, \Omega Z) \\ &\cong \text{Map}_*(A, X \times \Omega Z) \rightarrow \text{Map}_*(A, X), \end{aligned}$$

where μ^{op} is the dual of the structure isomorphism $(A \wedge K) \wedge S^1 \cong A \wedge (K \wedge^L S^1)$ of $A \wedge -$ as a $\text{Ho } \underline{\mathbf{SSet}}_*$ -module functor.

xxii) The dual statement of xxi) holds.

A **pre-triangulated category** is a closed $\text{Ho } \underline{\mathbf{SSet}}_*$ -module with all small products and coproducts together with a pre-triangulation.

Proposition 1.6.3.4. *Suppose \mathcal{S} is a pre-triangulated category.*

i) *Suppose $X \xrightarrow{f} Y \xrightarrow{g} Z$ is a cofiber sequence, W is an object. Then there exists a long exact sequences of pointed sets:*

$$\begin{aligned} \dots &\xrightarrow{(\Sigma \partial)^*} [\Sigma Z, W] \xrightarrow{(\Sigma g)^*} [\Sigma Y, W] \xrightarrow{(\Sigma f)^*} [\Sigma X, W] \\ &\xrightarrow{\partial^*} [Z, W] \xrightarrow{g^*} [Y, W] \xrightarrow{f^*} [X, W], \end{aligned}$$

with the following addition properties:

- *For any $a, b \in [Z, W]$, $g^*a = g^*b$ if and only if there is an $x \in [\Sigma X, W]$ such that $a \odot x = b$;*
- *For any $a, d \in [\Sigma X, W]$, $\partial^*c = \partial^*d$ if and only if there is an $y \in [\Sigma Y, W]$ such that $c \cdot (\Sigma f)^*y = d$.*

ii) *Suppose $X \xrightarrow{f} Y \xrightarrow{g} Z$ and $X' \xrightarrow{f'} Y' \xrightarrow{g'} Z'$ are cofiber sequences and we have a commutative diagram as follows:*

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \longrightarrow & Z \amalg \Sigma X \\ a \downarrow & & b \downarrow & & c \downarrow & & \downarrow c \amalg \Sigma a \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & \longrightarrow & Z' \amalg \Sigma X' \end{array}$$

Then if a, b are isomorphisms, so is c .

Dually there is a statement for fiber sequences.

Definition 1.6.3.5. Suppose \mathcal{S} and \mathcal{T} are pre-triangulated categories. A functor $F : \mathcal{S} \rightarrow \mathcal{T}$ is called a **right exact functor** if it is a left adjoint, it is a $\text{Ho } \underline{\text{SSet}}_*$ -module functor, and it maps any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ to a cofiber sequence $FX \xrightarrow{Ff} FY \xrightarrow{Fg} FZ$, where the coaction is given by

$$FZ \rightarrow F(Z \amalg \Sigma X) \cong FZ \amalg F\Sigma X \xrightarrow{1 \amalg \mu^{-1}} FZ \amalg \Sigma FX,$$

where μ is the structure isomorphism $(FX) \wedge S^1 \cong F(X \wedge^L S^1)$ of F as a $\text{Ho } \underline{\text{SSet}}_*$ -module functor. Dually a functor $G : \mathcal{T} \rightarrow \mathcal{S}$ is called a **left exact functor** if it is a right adjoint, its left adjoint is a $\text{Ho } \underline{\text{SSet}}_*$ -module functor, and it maps any fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ to a cofiber sequence $GX \xrightarrow{Gf} GY \xrightarrow{Gg} GZ$, where the action is given by

$$GX \times \Omega GZ \xrightarrow{1 \times (\mu^{\text{op}})^{-1}} GX \times G\Omega Z \cong G(X \times \Omega Z) \rightarrow GX,$$

where μ^{op} is the dual natural transformation of the structure isomorphism of its left adjoint as a $\text{Ho } \underline{\text{SSet}}_*$ -module functor. An adjunction $(F, G) : \mathcal{S} \rightarrow \mathcal{T}$ is called an **exact adjunction** if F is right exact and G is left exact.⁴

Lemma 1.6.3.6. *Pre-triangulated categories, exact adjunctions and $\text{Ho } \underline{\text{SSet}}_*$ -module natural transformations form a 2-category, which we denote $\underline{\text{PreTr}}$. There exists a duality 2-functor $-^{\text{op}}$ on $\underline{\text{PreTr}}$.*

Theorem 1.6.3.7. *The pseudo-2-functor*

$$\text{Ho} : \underline{\text{Model}}_* \rightarrow \underline{\text{CloModule}}_{\text{Ho } \underline{\text{SSet}}_*}$$

lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{Model}}_* \rightarrow \underline{\text{PreTr}}.$$

Moreover $\text{Ho} \circ (-^{\text{op}}) = (-^{\text{op}}) \circ \text{Ho} : \underline{\text{Model}}_* \rightarrow \underline{\text{PreTr}}$.

Definition 1.6.3.8. A **closed monoidal pre-triangulated category** is a closed $\text{Ho } \underline{\text{SSet}}_*$ -algebra (\mathcal{S}, \wedge, i) with all small products and coproducts, together with a pre-triangulation, such that for any $X \in \mathcal{S}$, the functors $X \wedge -, - \wedge X : \mathcal{S} \rightarrow \mathcal{S}$ are right exact, and the functors $\mathcal{H}om_l(X, -), \mathcal{H}om_r(X, -) : \mathcal{S} \rightarrow \mathcal{S}, \mathcal{H}om_l(-, X), \mathcal{H}om_r(-, X) : \mathcal{S}^{\text{op}} \rightarrow \mathcal{S}$ are left exact. We define a **closed symmetric monoidal pre-triangulated category** to be a closed monoidal pre-triangulated category that is also a closed symmetric $\text{Ho } \underline{\text{SSet}}_*$ -algebra.

Definition 1.6.3.9. For any closed monoidal pre-triangulated category \mathcal{S} , define $S^n = \Sigma^n S = i(S^n)$ for any nonnegative integer n .

Lemma 1.6.3.10. *Suppose \mathcal{S} is a closed symmetric monoidal pre-triangulated category, then the following diagram is commutative for any nonnegative integers m, n :*

$$\begin{array}{ccc} S^m \wedge S^n & \xrightarrow{\mu} & S^{m+n} \\ T \downarrow & & \downarrow (-1)^{mn} \mathbf{1} \\ S^n \wedge S^m & \xrightarrow{\mu} & S^{m+n} \end{array}$$

where μ is the structure isomorphism of the monoidal functor i and the negative sign is taken with respect to the cogroup structure of S^{m+n} .

⁴The notations here are OPPOSITE from the notations in the original book, since the notations given here coincide with our usual thoughts.

Definition 1.6.3.11. We define a **closed monoidal exact adjunction** between closed monoidal pre-triangulated categories to be an exact adjunction that is also a closed $\text{Ho } \underline{\text{SSet}}_*$ -algebra functor. A **closed symmetric monoidal exact adjunction** between closed monoidal pre-triangulated categories is an exact adjunction that is also a closed symmetric $\text{Ho } \underline{\text{SSet}}_*$ -algebra functor.

Lemma 1.6.3.12. *Closed monoidal pre-triangulated categories, closed monoidal exact adjunctions and $\text{Ho } \underline{\text{SSet}}_*$ -algebra natural transformations form a 2-category, which we will denote $\underline{\text{CloMonPreTr}}$. Closed symmetric monoidal pre-triangulated categories, closed symmetric monoidal exact adjunctions and $\text{Ho } \underline{\text{SSet}}_*$ -algebra natural transformations form a 2-category, which we denote $\underline{\text{CloSymMonPreTr}}$.*

Theorem 1.6.3.13. *The pseudo-2-functor*

$$\text{Ho} : \underline{\text{MonModel}}_* \rightarrow \underline{\text{CloAlg}}_{\text{Ho } \underline{\text{SSet}}_*}$$

lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{MonModel}}_* \rightarrow \underline{\text{CloMonPreTr}}.$$

The pseudo-2-functor

$$\text{Ho} : \underline{\text{SymMonModel}}_* \rightarrow \underline{\text{CloSymAlg}}_{\text{Ho } \underline{\text{SSet}}_*}$$

lifts to a pseudo-2-functor

$$\text{Ho} : \underline{\text{SymMonModel}}_* \rightarrow \underline{\text{CloSymMonPreTr}}.$$

1.7 Triangulated Categories and Stable Homotopy Categories

1.7.1 Triangulated Categories

Definition 1.7.1.1. A **triangulated category** is a pre-triangulated category such that Σ is an equivalence of categories. A pointed model category is **stable** if its homotopy category is triangulated. A **closed (symmetric) monoidal triangulated category** is a closed (symmetric) monoidal pre-triangulated category that is also a triangulated category. The full sub-2-category of $\underline{\text{PreTr}}(\underline{\text{CloMonPreTr}}, \underline{\text{CloSymMonPreTr}})$ consisting of all triangulated categories will be denoted $\underline{\text{Tr}}(\underline{\text{CloMonTr}}, \underline{\text{CloSymMonTr}})$.

Lemma 1.7.1.2. *Triangulated categories are additive.*

Remark 1.7.1.3. Any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ is completely determined by the maps f, g, ∂ , and for any cofiber sequences $X \xrightarrow{f} Y \xrightarrow{g} Z$ and $X' \xrightarrow{f'} Y' \xrightarrow{g'} Z'$ and maps $a : X \rightarrow X', b : Y \rightarrow Y', c : Z \rightarrow Z'$, c is Σa -equivariant if and only if $\Sigma a \circ \partial = \partial \circ c$. Dually any fiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ is completely determined by the maps f, g, ∂ , and for any fiber sequences $X \xrightarrow{f} Y \xrightarrow{g} Z$ and $X' \xrightarrow{f'} Y' \xrightarrow{g'} Z'$ and maps $a : X \rightarrow X', b : Y \rightarrow Y', c : Z \rightarrow Z'$, a is Ωc -equivariant if and only if $a \circ \partial = \partial \circ \Omega c$.

Proposition 1.7.1.4. *Any triangulated category is a classical triangulated category.*

Lemma 1.7.1.5. *Suppose \mathcal{S} is a triangulated category, and $\Sigma X \xrightarrow{\Sigma f} \Sigma Y \xrightarrow{\Sigma g} \Sigma Z \xrightarrow{\Sigma h} \Sigma^2 X$ is a cofiber sequence. Then so is $X \xrightarrow{-f} Y \xrightarrow{-g} Z \xrightarrow{-h} \Sigma X$.*

Proposition 1.7.1.6. Suppose \mathcal{S} is a triangulated category. Then $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X$ is a cofiber sequence, if and only if $\Omega Z \xrightarrow{-\eta_X^{-1} \circ (\Omega h)} X \xrightarrow{f} Y \xrightarrow{\varepsilon_Z^{-1} \circ g} \Omega \Sigma Z$ is a cofiber sequence.

Lemma 1.7.1.7. Suppose \mathcal{S} is a triangulated category. Then for any cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X$ and object W , the sequence

$$[W, X] \xrightarrow{f_*} [W, Y] \xrightarrow{g_*} [W, Z] \xrightarrow{h_*} [W, \Sigma X]$$

is exact.

Theorem 1.7.1.8. Suppose \mathcal{S} is a triangulated category. Then $\Omega Z \xrightarrow{f} X \xrightarrow{g} Y \xrightarrow{h} Z$ is a fiber sequence, if and only if $\Omega Z \xrightarrow{f} X \xrightarrow{g} Y \xrightarrow{-\varepsilon_Z^{-1} \circ h} \Omega \Sigma Z$ is a cofiber sequence.

Definition 1.7.1.9. Suppose \mathcal{C} is a model category, and we have a commutative square in \mathcal{C} :

$$\begin{array}{ccc} W & \xrightarrow{u} & X \\ v \downarrow & & \downarrow g \\ Z & \xrightarrow{f} & Y \end{array}$$

Take the following pullback square:

$$\begin{array}{ccc} W' & \longrightarrow & X' \\ \downarrow & & \downarrow \delta(r_Y \circ g) \\ Z' & \xrightarrow{\delta(r_Y \circ f)} & RY \end{array}$$

The original square is called a **homotopy pullback square** if the induced map $W \rightarrow W'$ is a weak equivalence. Dually take the following pushout square:

$$\begin{array}{ccc} QW & \xrightarrow{\alpha(u \circ q_W)} & X'' \\ \alpha(v \circ q_W) \downarrow & & \downarrow \\ Z'' & \longrightarrow & Y'' \end{array}$$

The original square is called a **homotopy pushout square** if the induced map $Y'' \rightarrow Y$ is a weak equivalence.

Proposition 1.7.1.10. Suppose \mathcal{C} is a pointed model category. Then homotopy pullback squares and homotopy pushout square coincide if and only if \mathcal{C} is stable.

Proposition 1.7.1.11. Suppose \mathcal{S}, \mathcal{T} are triangulated categories. An adjunction $(F, G) : \mathcal{S} \rightarrow \mathcal{T}$ is an exact adjunction if and only if F is right exact, if and only if G is left exact.

Lemma 1.7.1.12. Suppose \mathcal{S} is a closed symmetric monoidal triangulated category. Then the following diagram is commutative for any integers m, n :

$$\begin{array}{ccc} S^m \wedge S^n & \xrightarrow{\mu} & S^{m+n} \\ T \downarrow & & \downarrow (-1)^{mn} \mathbf{1} \\ S^n \wedge S^m & \xrightarrow{\mu} & S^{m+n} \end{array}$$

where $S^{-k} = \Omega^k S$ for any $k > 0$, μ is the structure isomorphism of the monoidal functor i , combined with the unit or counit isomorphisms, if necessary.

1.7.2 Stable Homotopy Categories

Definition 1.7.2.1. Suppose \mathcal{S} is a (pre-)triangulated category, \mathcal{G} is a set of objects. \mathcal{G} is called a **set of weak generators** if $[\Sigma^n G, X] = 0$ for any $G \in \mathcal{G}$ and (nonnegative) integers n implies $X \cong *$. (When $n < 0$, take $\Sigma^n G = \Omega^{-n} G$)

Definition 1.7.2.2. Suppose \mathcal{S} is a pre-triangulated category. An object X is called **algebraically finite** if for any set $\{Y_\alpha\}_{\alpha \in K}$ of objects, the induced map

$$\operatorname{colim}_{S \subseteq K, S \text{ finite}} \left[X, \coprod_{\alpha \in S} Y_\alpha \right] \rightarrow \left[X, \coprod_{\alpha \in K} Y_\alpha \right]$$

is an isomorphism.

Definition 1.7.2.3. An **algebraic stable homotopy category** is a closed symmetric monoidal triangulated category with a set of algebraically finite weak generators.

Proposition 1.7.2.4. *Suppose \mathcal{C} is a pointed model category, λ is an ordinal, X is a λ -sequence of cofibrations of cofibrant objects with colimit X , and Y is fibrant. If $[X_\beta, Y] = 0$ for any $\beta < \lambda$, then $[X, Y] = 0$.⁵*

Theorem 1.7.2.5. *Suppose \mathcal{C} is a cofibrantly generated pointed model category with generating cofibrations I . Then the set of cofibers of maps of I is a set of weak generators for $\operatorname{Ho} \mathcal{C}$.*

Lemma 1.7.2.6. *Suppose \mathcal{C} is a finitely generated model category, λ is an ordinal, X, Y are λ -sequences of cofibrations, $p : X \rightarrow Y$ is a natural transformation such that p_α is a (trivial) fibration for any $\alpha < \lambda$. Then $\operatorname{colim} p_\alpha$ is a (trivial) fibration.*

Corollary 1.7.2.7. *Suppose \mathcal{C} is a finitely generated model category with generating cofibrations I , \mathcal{D} is a subcategory such that domains and codomains of I are small relative to \mathcal{D} . Then transfinite compositions of weak equivalences in \mathcal{D} are weak equivalences.*

Theorem 1.7.2.8. *Suppose \mathcal{C} is a finitely generated pointed model category. Then any cofibrant object that is finite relative to cofibrations is algebraically finite in $\operatorname{Ho} \mathcal{C}$.*

Theorem 1.7.2.9. *The pseudo-2-functor Ho lifts to a pseudo-2-functor Ho from the 2-category of finitely generated stable symmetric monoidal model categories to the 2-category of algebraic stable homotopy categories.*

⁵This proposition remains unproved in class.

Chapter 2

Infinity Categories (Semesters 2 & 3)

The following contents are given in the lectures in the spring and autumn semester of 2019. 24 lectures with a total time of 121 hours have been given so far in order to finish the book *Higher Topos Theory*. Infinity categories are really abstract things, so motivations, together with sketches of hard proofs are given in order to understand them better. However, details of proofs to the statements are still omitted. A few statements remain unproved in class due to lack of corresponding knowledge; they are marked with \dagger . These statements do not include those that are too boring to present the proofs, which do not contain any technical difficulties, however.

2.0 Preliminaries

2.0.1 Preliminaries for Ordinary Categories

2.0.2 Preliminaries for Model Categories

2.1 Introduction to Infinity Categories

2.1.1 Foundations of Infinity Categories

We use some simple examples to give the motivation of ∞ -categories.

Example 2.1.1.1. Suppose X is a topological space. The **fundamental groupoid** of X , denoted $\Pi_{\leq 1}(X)$, is the category with:

- objects being all points in X ; and
- $\text{Hom}_{\Pi_{\leq 1}(X)}(a, b)$ being the homotopy classes of all paths from a to b .

This is some definition that we shall already have seen in algebraic topology. This gives rise to a functor $\Pi_{\leq 1} : \mathbf{Top} \rightarrow \mathbf{Gpd}$. Moreover, for any map $f : X \rightarrow Y$ of topological spaces, $\Pi_{\leq 1}(f)$ is an equivalence of categories, if and only if it is fully faithful and essentially surjective, where the first assertion indicates that $\pi_1(f)$ is a bijection, and the second assertion indicates that $\pi_0(f)$ is a bijection.

However, a topological space, contains far more information than its 0th and 1st homotopy groups. There also not only exist homotopies between paths, there also exist homotopies between homotopies, and homotopies between the homotopies between homotopies, etc. How shall we encode such higher structures? We may think of a “category”, with not only morphisms between objects, but also morphisms between morphisms, and morphisms between the morphisms between morphisms, etc. Specifically, we would like to define $\Pi(X)$ as follows:

- Objects being all points a, b, \dots in X ;
- $\text{Hom}_{\Pi(X)}(a, b)$ being all paths γ, δ, \dots from a to b ;
- $\text{Hom}_{\Pi(X)}(\gamma, \delta)$ being all homotopies H, K, \dots from γ to δ ;
- \dots

And we would like to obtain the property that for any map $f : X \rightarrow Y$ of topological spaces, $\Pi(f)$ is an “equivalence” of “categories”, if and only if f induces isomorphisms on all homotopy groups; that is, f is a weak homotopy equivalence. The theory of ∞ -categories are introduced to make this thought precise and to study their properties.

Example 2.1.1.2. We recall homotopies between maps in model categories: Given a model category \mathcal{C} and cofibrant-fibrant objects A, B , maps $f, g : A \rightarrow B$, a homotopy from f to g is equivalent to a map $h : A \otimes \Delta[1] \rightarrow B$, such that $h|_{A \otimes 0} = f, h|_{A \otimes 1} = g$. However, cosimplicial frames not only allow us to construct the cylinder object of A , but also higher structures. For example, a map $A \otimes \Delta[2] \rightarrow B$ may be viewed as a homotopy between the three homotopies given by its restriction on $A \otimes \partial\Delta[2]$, a map $A \otimes \Delta[3] \rightarrow B$ may be viewed as a homotopy between the four homotopies between homotopies given by its restriction on $A \otimes \partial\Delta[3]$. The theory of ∞ -categories gives us a tool on how to formalize this kind of higher structures, and to study their properties.

Notice, that in these two examples, higher morphisms are “invertible” (homotopies are invertible, at least up to higher homotopies). This also coincides with most usages of ∞ -categories, as in most applications we may see. Therefore, we only talk about ∞ -categories, whose “ k -morphisms” are invertible whenever $k \geq 2$.¹

So how shall we define them? The most naïve approach is to define them as what we did in Section 1.1.3. However, notice one thing: in both examples, associativity and unit axioms fails to be strict, but only hold up to homotopy, i.e. higher morphisms. To fix this issue, one must add certain extra diagrams to give the correct notion of higher categories, rather than the ones given in Section 1.1.3. (The reason we introduce them in the first chapter is to make the language we use concise. However, pseudo-2-functors, DO give an approach to functors between ∞ -categories, as they keep the homotopy-coherent ideas. From now on, we will call that kind of 2-category introduced in Section 1.1.3 **strict 2-categories**.) On the other hand, even the explicit definition of a 3-category is EXTREMELY complicated, and things are just getting worse when we pass to 4-categories and beyond.

Fortunately, we have an alternative approach to the theory of ∞ -categories.

From our naïve thought, if we take the “Hom-thing” between two object in an ∞ -category, it will result in an $(\infty, 0)$ -category. In other words, ∞ -categories are categories enriched over $(\infty, 0)$ -categories. Consequencely, we only have to make a model for $(\infty, 0)$ -categories.

Notice that in Example 2.1.1.1, all morphisms, including 1-morphisms, are invertible. This means that $\Pi(X)$ “should be an $(\infty, 0)$ -category”. On the other hand, there is a general axiom, known as the **homotopy hypothesis**, accepted by higher categorists, which is roughly stated as follows:

- Every $(\infty, 0)$ -category has the form $\Pi(X)$ for some topological space X . Moreover two $(\infty, 0)$ -categories are equivalent, if and only if their corresponding topological spaces are weakly equivalent.

¹In higher category theory, we use the term (∞, n) -category to indicate an ∞ -category whose k -morphisms are invertible whenever $k \geq n + 1$. In other words, what we are focusing about is $(\infty, 1)$ -categories.

This axiom ensures that we have a one-to-one correspondence between $(\infty, 0)$ -categories and equivalences classes of topological spaces, where two spaces are equivalent if and only if they are weakly equivalent. That is, we use objects in $\text{Ho } \underline{\mathbf{Top}} \simeq \text{Ho } \underline{\mathbf{K}} \simeq \text{Ho } \underline{\mathbf{SSet}}$ to describe $(\infty, 0)$ -categories. Because of such important role they play in higher category theory, we give them a special name: **homotopy types**. We write $\underline{\mathbf{H}}$ for the category of homotopy types.

Now we have a schematic view of $(\infty, 1)$ -categories: loosely speaking, they are just $\underline{\mathbf{H}}$ -enriched categories. However, doing computations in $\underline{\mathbf{H}}$ -enriched categories are really difficult. Moreover, not all $\underline{\mathbf{H}}$ -enriched categories are $(\infty, 1)$ -categories. This is because of the difference between homotopy-commutative diagrams and homotopy-coherent diagrams, as we will discuss in the next section. Therefore, we need to give correct models for $(\infty, 1)$ -categories. Moreover, we require these models to be easy to compute on. We shall now give three different models for us to work with.

Remark 2.1.1.3. Although we are giving different models, they are actually representing the same thing: $(\infty, 1)$ -categories. Therefore it is important to show that these models are equivalent, at least up to equivalence between categories. We will develop how equivalences between categories are defined in these different models, and how one model can be translated into another, and show that under equivalences between categories, all these models are equivalent.

The first model is topological categories.

Definition 2.1.1.4. A **topological category** is a category enriched over $\underline{\mathbf{K}}$, the category of all k -spaces.²

By the definition of a topological category, if we apply the localization functor $\underline{\mathbf{K}} \rightarrow \underline{\mathbf{H}}$, a topological category becomes an $\underline{\mathbf{H}}$ -enriched category. We will now specify this localization functor with the symbol h . Therefore, topological categories are indeed models for ∞ -categories.

Definition 2.1.1.5. The **homotopy category** of a topological category \mathcal{C} is $h\mathcal{C}$. Two topological categories are said to be **(weakly) equivalent** if their homotopy categories are equivalent (viewed as $\underline{\mathbf{H}}$ -enriched categories).

Note that an equivalence between topological categories, at least in higher category theory, does not mean it is an equivalence between $\underline{\mathbf{K}}$ -enriched categories. This is because essentially the morphisms spaces we are dealing with are actually homotopy types, not real spaces.

The second model is simplicial categories.

Definition 2.1.1.6. A **simplicial category** is a category enriched over $\underline{\mathbf{SSet}}$, the category of all simplicial sets.³

By the definition of a simplicial category, if we apply the localization functor $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{H}}$, a simplicial category becomes an $\underline{\mathbf{H}}$ -enriched category. We will, again, specify this localization functor with the symbol h . Therefore, simplicial categories are also indeed models for ∞ -categories.

Definition 2.1.1.7. The **homotopy category** of a simplicial category \mathcal{C} is $h\mathcal{C}$. Two simplicial categories are said to be **(weakly) equivalent** if their homotopy categories are equivalent (viewed as $\underline{\mathbf{H}}$ -enriched categories).

²The reason we use k -spaces here, is that $\underline{\mathbf{K}}$ is a closed monoidal category, and the geometric realization plays well with k -spaces. From now on, unless otherwise stated, when we say “topological spaces”, what we actually mean is “ k -spaces.”

³Note that a simplicial category is not equivalent to a simplicial object in the category of the categories.

We now consider the relations between these two definitions. We already know that there exists an adjunction $(|-|, \text{Sing}) : \underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{K}}$, that is a closed symmetric monoidal Quillen equivalence. Consequently, we get an adjunction $(|-|, \text{Sing}) : \underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}} \rightarrow \underline{\mathbf{Cat}}_{\underline{\mathbf{K}}}$. By Section 2.0.2, this adjunction is a Quillen equivalence between the canonical model structures on both sides. Thus both the left and right adjoints preserve equivalences, and both the unit morphisms and counit morphisms are equivalences. Thus the two models can be regarded as equivalent models for $(\infty, 1)$ -categories.

However, these two models have disadvantages. As we have said before, in most $(\infty, 1)$ -categories (that we actually work with), compositions of morphisms is associative only up to homotopy. But in a topological category, composition of morphisms are associative up to nose. Therefore we must figure out how to transform a homotopy-coherent diagram into a strictly commutative diagram. Although this is always possible (See Section [ERROR! SECTION NOT FOUND]), it is convenient for us to get a more flexible model of $(\infty, 1)$ -categories.

We note that in some cases, simplicial sets can be regarded as categories. We may regard 0-simplices of the simplicial set as objects of the category, and 1-simplices of the simplicial set as morphisms of the category. For example, the simplicial set $\Delta[n]$ can be regarded as the category, that comes from the partially ordered set with $n + 1$ objects $[n]$. This gives rise to the following definition:

Definition 2.1.1.8. We consider the functor $\Delta \rightarrow \underline{\mathbf{Cat}}, [n] \mapsto [n]$. This gives a cosimplicial object in the category of all categories. By Proposition 1.3.1.23, this gives rise to an adjunction $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}$. We call the right adjoint of this adjunction the **nerve functor**, denoted N .

Now, for any category \mathcal{C} , all n -simplices in $N\mathcal{C}$ are simply all composable sequences of maps

$$X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} \dots \xrightarrow{f_{n-1}} X_n.$$

The nerve functor enjoys the following property:

Proposition 2.1.1.9. *Suppose K is a simplicial set. The following conditions are equivalent:*

- i) *There exists a category \mathcal{C} and an isomorphism of simplicial sets $K \cong N(\mathcal{C})$;*
- ii) *Any map $\Lambda^i[n] \rightarrow K$, where $0 < i < n$, extends uniquely to a map $\Delta[i] \rightarrow K$.*

Let us take a pause here to see what does this proposition means. What is a map $\Delta[2] \rightarrow N(\mathcal{C})$? it means taking three morphisms $f : X \rightarrow Y, g : Y \rightarrow Z$ and $h : X \rightarrow Z$ in \mathcal{C} , such that $h = gf$. Similarly, a map $\Lambda^1[2] \rightarrow N(\mathcal{C})$ means taking two morphisms $f : X \rightarrow Y, g : Y \rightarrow Z$. Now the statement “any map $\Lambda^1[2] \rightarrow N(\mathcal{C})$, extends uniquely to a map $\Delta[2] \rightarrow N(\mathcal{C})$ ” simply means that every two morphisms that are tip-to-tail are uniquely composable. Similarly, the statement “any map $\Lambda^i[3] \rightarrow N(\mathcal{C})$, where $i = 1, 2$, extends uniquely to a map $\Delta[3] \rightarrow N(\mathcal{C})$ ” means that composition is strictly associative. Note that we do not include the cases $i = 0$ and $i = n$. This is because “any map $\Lambda^0[2] \rightarrow N(\mathcal{C})$, extends uniquely to a map $\Delta[2] \rightarrow N(\mathcal{C})$ ” will indicate that every map has a left inverse, which is only possible whenever \mathcal{C} is a groupoid.

We have seen that certain simplicial sets can be used to model ordinary categories. We may want to show that certain simplicial sets can be used to model $(\infty, 1)$ -categories.

Now, suppose K is a simplicial set, that is considered to be an $(\infty, 1)$ -category. What kind of properties would we like K to have? We would like to consider 0-simplices as objects, 1-simplices as morphisms, as usual. We would also like every two morphisms that are tip-to-tail are composable. This means that any map $\Lambda^1[2] \rightarrow K$, extends to a map $\Delta[2] \rightarrow K$. However, if we still require the composition to be unique, we are on our way back to ordinary categories. In fact, it is unreasonable, and unnatural, to require composition of two morphisms to be unique. For example, there exists multiple ways to join two paths to obtain a longer

path in any topological space X , meaning that there exists multiple choices of compositions in $\Pi(X)$. Nevertheless, these choices are homotopic. The condition that different compositions are homotopic, can be recovered by the statement “any map $\Lambda^i[3] \rightarrow K$, where $i = 1, 2$, extends to a map $\Delta[3] \rightarrow K$ ”. Similarly, higher homotopy relations can be recovered by the statement “any map $\Lambda^i[n] \rightarrow K$, where $0 < i < n$, extends to a map $\Delta[i] \rightarrow K$ ”. Again, we do not include the cases $i = 0$ and $i = n$, since they are only satisfied when every morphism in K is “invertible”.

We now make our discussion precise:

Definition 2.1.1.10. A **quasi-category**, is a simplicial set K , such that

$$(K \rightarrow *) \in \text{RLP}(\{\Lambda^i[n] \rightarrow \Delta[n] \mid 0 < i < n\}).$$

Since this is the mainly used model for $(\infty, 1)$ -categories, from now on, when we say an ∞ -category, we refer to a quasi-category, unless stated otherwise.

Example 2.1.1.11. The nerve of an ordinary category is an ∞ -category.

Example 2.1.1.12. Kan complexes are ∞ -categories. Moreover, they enjoy right lifting property with respect to $\Lambda^i[n] \rightarrow \Delta[n]$ even if $i = 0$ or $i = n$, meaning that every morphism in a Kan complex is “invertible”. For this reason, we simply define **∞ -groupoids** or **$(\infty, 0)$ -categories** to be Kan complexes, and say that two ∞ -groupoids are **equivalent** if they are weak equivalent. This give a nice realization of the homotopy hypothesis, and the construction Π taking a topological space to an ∞ -groupoid can be simply realized by Sing .

We now discuss the relations between ∞ -categories and simplicial categories. In order to show that these two models are equivalent (in an appropriate sense), we must from a pair of functors $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}}$ and $\underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}} \rightarrow \underline{\mathbf{SSet}}$. As shown in Proposition 1.3.1.23, a cosimplicial simplicial category will give rise to an adjunction $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}}$. In the case of ordinary categories, we gave the cosimplicial category $\Delta \rightarrow \underline{\mathbf{Cat}}, [n] \mapsto [n]$. This makes the composition of morphisms in a simplicial set strictly associative, which is what in the case of an ordinary category we want. Consequencely, we would like to “thicken the morphism space” to give rise to the homotopy-coherent idea of composing the morphisms. This gives rise to the following definition:

Definition 2.1.1.13. We define the functor $\mathfrak{C} : \underline{\mathbf{TOSet}} \rightarrow \underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}}$ as follows. For any totally ordered set J , define $\mathfrak{C}[J]$ to be the simplicial category with:

- Objects being all objects of J ;
- For any $i, j \in J$, $\text{Hom}_{\mathfrak{C}[J]}(i, j) = \emptyset$ if $i > j$, and $\text{Hom}_{\mathfrak{C}[J]}(i, j) = N(P_{i,j})$ if $i \leq j$, where $P_{i,j}$ is the partially ordered set $\{I \subseteq J \mid i, j \in I; \forall k \in I, i \leq k \leq j\}$;
- For $i \leq j \leq k$, the composition law $\text{Hom}_{\mathfrak{C}[J]}(i, j) \times \text{Hom}_{\mathfrak{C}[J]}(j, k) \rightarrow \text{Hom}_{\mathfrak{C}[J]}(i, k)$ is induced by the map of partially ordered sets $P_{i,j} \times P_{j,k} \rightarrow P_{i,k}, (I, I') \mapsto I \cup I'$;

and for any maps $f : J \rightarrow J'$ between totally ordered sets, define $\mathfrak{C}[f]$ to be the functor with:

- $\mathfrak{C}[f](i) = f(i)$ for any object $i \in J$;
- For any $i, j \in J$, the map $\text{Hom}_{\mathfrak{C}[J]}(i, j) \rightarrow \text{Hom}_{\mathfrak{C}[J']}(f(i), f(j))$ is induced by the map $P_{i,j} \rightarrow P_{f(i), f(j)}, I \mapsto f(I)$.

This functor induces a cosimplicial object in the category of simplicial categories, which further induces an adjunction $\underline{\mathbf{SSet}} \rightarrow \underline{\mathbf{Cat}}_{\underline{\mathbf{SSet}}}$. We denote this adjunction by (\mathfrak{C}, N) , and call the right adjoint the **simplicial nerve** functor.

This is really a complicated definition, so we shall pause again to figure out this definition. Take $\mathfrak{C}[\Delta[3]]$ as example. We have

$$\mathrm{Hom}_{\mathfrak{C}[\Delta[3]]} = \left(\begin{array}{ccc} 03 & \longrightarrow & 013 \\ \downarrow & \searrow & \downarrow \\ 023 & \longrightarrow & 0123 \end{array} \right).$$

(Notice that it also contains two nondegenerate 2-simplices that are not shown.) We notice that each vertex of this Hom-space represents a way from 0 to 3. In particular, there is a way from 0 to 3 that does not pass through any other vertices, which is the 03 vertex. We denote the morphism $0 \rightarrow 3$ represented by the vertex 03 as p_{03} . Now we notice that $p_{03} \neq p_{13}p_{01}$. However, the 1-simplex from 03 to 013 indicates that these two morphisms are homotopic to each other. Moreover, the fact that the morphism space is weakly contractible, means that all possible compositions are canonically homotopic. This corresponds to the point that $\mathfrak{C}[n]$ is the “thicken version” of $[n]$ that we have dropped the strict associative condition and replace it by the idea that compositions are commutative only up to coherent homotopy. Take another example, $\mathfrak{C}[\partial\Delta[3]]$. We have

$$\mathrm{Hom}_{\mathfrak{C}[\partial\Delta[3]]} = \left(\begin{array}{ccc} 03 & \longrightarrow & 013 \\ \downarrow & & \downarrow \\ 023 & \longrightarrow & 0123 \end{array} \right).$$

We may see that the path of maps $03 \rightarrow 013 \rightarrow 0123 \leftarrow 023 \leftarrow 03$ is not homotopic to the constant path, which corresponds well with the fact that $\partial\Delta[3]$ is not weakly contractible.

As an exercise, the reader can show that if $0 \leq i \leq j \leq n$, then the map $\mathrm{Hom}_{\mathfrak{C}[\partial\Delta[n]]}(i, j) \rightarrow \mathrm{Hom}_{\mathfrak{C}[\Delta[n]]}(i, j)$ is identity unless $i = 0, j = n$, in which case it is a cofibration; then the map $\mathrm{Hom}_{\mathfrak{C}[\Delta[n]]}(i, j) \rightarrow \mathrm{Hom}_{\mathfrak{C}[\partial\Delta[n]]}(i, j)$ is identity unless $i = 0, j = n$, in which case it is an anodyne extension.

Definition 2.1.1.14. The **topological nerve** of a topological category \mathcal{C} is defined to be $\mathrm{NSing} \mathcal{C}$.

Proposition 2.1.1.15. Suppose \mathcal{C} is a fibrant simplicial category (in the canonical model structure of the category of simplicial categories) (which is equivalent to all of its Hom-spaces are Kan complexes). Then $\mathrm{N}\mathcal{C}$ is an ∞ -category.

Corollary 2.1.1.16. The topological nerve of any topological category is an ∞ -category.

We now give the definition to equivalences between ∞ -categories.

Definition 2.1.1.17. Suppose S is a simplicial set. The **homotopy category** of S is $\mathrm{h}S := \mathrm{h}\mathfrak{C}[S]$. A map $f : S \rightarrow T$ between simplicial sets is called a **categorical equivalence** if $\mathrm{h}f$ is an equivalence of $\underline{\mathbf{H}}$ -enriched categories.

We now state the following theorem, which will be proved in Section [\[ERROR! SECTION NOT FOUND\]](#).

Theorem 2.1.1.18. There exists a model structure on $\underline{\mathbf{S}}\mathbf{Set}$, called the **Joyal model structure**, where weak equivalences are all the categorical equivalences and fibrant objects are the quasi-categories, such that the adjunction

$$\begin{array}{ccc} & \mathfrak{C} & \\ \text{(\underline{S}Set, Joyal)} & \xrightleftharpoons{\quad} & \text{(\underline{Cat}_{\underline{\mathbf{S}Set}}, std)} \\ & \mathbf{N} & \end{array}$$


is a Quillen equivalence.

This means that the theory of quasi-categories and the theory of simplicial categories are equivalent models of ∞ -categories.

Finally we give the following definition:

Definition 2.1.1.19. Suppose \mathcal{C} is a simplicial category, or a topological category, or a simplicial set. We define $\text{Ho } \mathcal{C} := \pi_0 h \mathcal{C}$.

To get a better view of what we have done in this section, it would be better if you have this graph in mind. Here all the arrows above the categories are left adjoints, where all the arrows below the categories are right adjoints, equivalences means Quillen equivalences.



2.1.2 The Language of Infinity Categories

We now describe how most aspects of ordinary categories extends to infinity categories.

Remark 2.1.2.1. As we have stated above, since the three models are equivalent models for $(\infty, 1)$ -categories, what we have defined for ∞ -categories (that are invariant under categorical equivalences) should extend naturally to topological categories and simplicial categories. The readers may feel free to use the results we have stated for ∞ -categories on topological categories and simplicial categories, if they prefer.

Definition 2.1.2.2. Suppose \mathcal{C} is an ∞ -category. An **object** in \mathcal{C} is a vertex in \mathcal{C} . A **morphism** in \mathcal{C} is a 1-simplex in \mathcal{C} . The **domain** and **codomain** of a morphism $\phi : \Delta[1] \rightarrow \mathcal{C}$ are $\phi(0)$ and $\phi(1)$, respectively. If a morphism ϕ has domain x and codomain y , we write $\phi : x \rightarrow y$. The **identity morphism** of an object x is $\mathbb{1}_x := s_0(x) : x \rightarrow x$.

Definition 2.1.2.3. Suppose S is a simplicial set. We define the **opposite** of S to be the simplicial set S^{op} , with $S_n^{\text{op}} = S_n$ but

$$(d_i : S_n^{\text{op}} \rightarrow S_{n-1}^{\text{op}}) = (d_{n-i} : S_n \rightarrow S_{n-1}), (s_i : S_n^{\text{op}} \rightarrow S_{n+1}^{\text{op}}) = (s_{n-i} : S_n \rightarrow S_{n+1}).$$

If \mathcal{C} is an ∞ -category, we call \mathcal{C}^{op} the **opposite category** of \mathcal{C} .

We next consider equivalences between morphisms.

Definition 2.1.2.4. Suppose \mathcal{C} is an ∞ -category. Two morphisms $f, g : x \rightarrow y$ in \mathcal{C} are said to be **homotopic**, written as $f \simeq g$, if one of the following equivalence relations hold:

- i) There exists a 2-simplex $\begin{array}{ccc} x & \xrightarrow{f} & y \\ \mathbb{1}_x \downarrow & \searrow & \nearrow \\ x & \xrightarrow{g} & y \end{array}$ in \mathcal{C} ;
- ii) There exists a 2-simplex $\begin{array}{ccc} x & \xrightarrow{g} & y \\ \mathbb{1}_x \downarrow & \searrow & \nearrow \\ x & \xrightarrow{f} & y \end{array}$ in \mathcal{C} ;
- iii) There exists a 2-simplex $\begin{array}{ccc} & & y \\ & \nearrow f & \downarrow \mathbb{1}_y \\ x & \xrightarrow{g} & y \end{array}$ in \mathcal{C} ;

- iv) There exists a 2-simplex $x \begin{array}{c} \nearrow^g y \\ \searrow_f y \end{array} \downarrow \mathbb{1}_y$ in \mathcal{C} .

Proposition 2.1.2.5. *The homotopic relation is an equivalence relation.*

Definition 2.1.2.6. Suppose \mathcal{C} is an ∞ -category. We define a category $\pi(\mathcal{C})$ as follows:

- i) Its objects are vertices of \mathcal{C} ;
- ii) For any $x, y \in \mathcal{C}_0$, define $\text{Hom}_{\pi(\mathcal{C})}(x, y) = \{f : x \rightarrow y \in \mathcal{C}_1\} / \simeq$;
- iii) For any $x \in \mathcal{C}_0$, define $\mathbb{1}_x = [s_0 x]$;
- iv) For any morphisms $[f] : x \rightarrow y, [g] : y \rightarrow z$ in $\pi(\mathcal{C})$, take a lift of the following diagram:

$$\begin{array}{ccc} \Lambda^1[2] & \xrightarrow[\text{\scriptsize } g \text{ on } \Delta\{1,2\}]{\text{\scriptsize } f \text{ on } \Delta\{0,1\}} & \mathcal{C} \\ \downarrow & \nearrow \text{\scriptsize } h & \\ \Delta[2] & & \end{array}$$

and define $[g] \circ [f] = [d_1 h]$.

Proposition 2.1.2.7. *The category $\pi(\mathcal{C})$ is isomorphic to the category $\text{Ho}(\mathcal{C})$, meaning that two morphisms $f \simeq g$ if and only if they are equal in $\text{Ho } \mathcal{C}$.*

Next we discuss the mapping spaces between objects.

Definition 2.1.2.8. Suppose S is a simplicial set, x, y are vertices in S . We define the **mapping space** from x to y to be $\text{Map}_S(x, y) := \text{Hom}_{\mathbf{h}S}(x, y) \in \mathbf{H}$.

So how shall we compute this mapping space? By definition, $\text{Map}_S(x, y)$ is the homotopy type of $\text{Hom}_{\mathcal{C}[S]}(x, y)$. The advantage of this definition is that it has a natural composition law, and it works for all simplicial sets, not just ∞ -categories. However, this construction is quite complicated, and it is usually not a Kan complex, which makes it difficult to verify its properties.

We shall now introduce another simplicial set that represents the homotopy type $\text{Map}_S(x, y)$, at least when S is an ∞ -category.

Definition 2.1.2.9. Suppose S is a simplicial set, x, y are vertices in S . We define

$$\text{Hom}_S^R(x, y) = \{\sigma \in S_{n+1} \mid \sigma|_{\Delta\{0, \dots, n\}} = x, \sigma(n+1) = y\},$$

and

$$\text{Hom}_S^L(x, y) = \{\sigma \in S_{n+1} \mid \sigma(0) = x, \sigma|_{\Delta\{1, \dots, n+1\}} = y\}.$$

Moreover we define

$$\text{Hom}_S(x, y) = \{\sigma : \Delta[n] \times \Delta[1] \rightarrow S \mid \sigma|_{\Delta[n] \times \{0\}} = x, \sigma|_{\Delta[n] \times \{1\}} = y\}.$$

The following proposition will be proved in Section [ERROR! SECTION NOT FOUND] and [ERROR! SECTION NOT FOUND].

Proposition 2.1.2.10. *If S is an ∞ -category, then for any objects x, y , the simplicial sets $\text{Hom}_S^R(x, y), \text{Hom}_S^L(x, y), \text{Hom}_S(x, y)$ are Kan complexes. There exists a canonical isomorphism*

$$\text{Hom}_S^R(x, y) \xrightarrow{\simeq} \text{Hom}_{\mathcal{C}[S]}(x, y)$$

in the category \mathbf{H} . The natural inclusions

$$\text{Hom}_S^R(x, y) \hookrightarrow \text{Hom}_S(x, y) \hookleftarrow \text{Hom}_S^L(x, y)$$

are weak homotopy equivalences.

We next discuss when can we say a morphism is an equivalence.

Definition 2.1.2.11. Suppose \mathcal{C} is a topological category, of a simplicial category, or an ∞ -category. A morphism f in \mathcal{C} is called an **equivalence**, or **invertible**, if it is an isomorphism in $\mathbf{h}\mathcal{C}$.

The following criterion is useful while we want to determine whether a morphism is an equivalence or not.

Proposition 2.1.2.12. Suppose \mathcal{C} is a topological category, $f : X \rightarrow Y$ is a morphism in \mathcal{C} . The following conditions are equivalent:

- i) f is an equivalence;
- ii) f has a homotopy inverse; that is, there exists a morphism $g : Y \rightarrow X$ such that $fg \simeq \mathbb{1}_Y$ and $gf \simeq \mathbb{1}_X$;
- iii) For any $Z \in \mathcal{C}$, composition with f induces a homotopy equivalence $\mathrm{Hom}_{\mathcal{C}}(Z, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(Z, Y)$;
- iv) For any $Z \in \mathcal{C}$, composition with f induces a weak homotopy equivalence $\mathrm{Hom}_{\mathcal{C}}(Z, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(Z, Y)$;
- v) For any $Z \in \mathcal{C}$, composition with f induces a homotopy equivalence $\mathrm{Hom}_{\mathcal{C}}(Y, Z) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, Z)$;
- vi) For any $Z \in \mathcal{C}$, composition with f induces a weak homotopy equivalence $\mathrm{Hom}_{\mathcal{C}}(Y, Z) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, Z)$.

The following proposition will be proved in Section [ERROR! SECTION NOT FOUND].

Proposition 2.1.2.13. Suppose \mathcal{C} is an ∞ -category, ϕ is a morphism in \mathcal{C} . ϕ is an equivalence, if and only if for every map $f : \Lambda^0[n] \rightarrow \mathcal{C}$ such that $f|_{\Delta\{0,1\}} = \phi$, f extends to a map $\Delta[n] \rightarrow \mathcal{C}$.

Using this proposition we obtain the following result:

Proposition 2.1.2.14. Suppose \mathcal{C} is a simplicial set. The following conditions are equivalent:

1. \mathcal{C} is an ∞ -groupoid;
2. \mathcal{C} is an ∞ -category and $\mathrm{Ho} \mathcal{C}$ is a groupoid;
3. \mathcal{C} is an ∞ -category and \mathcal{C} has extension property with respect to all inclusions $\Lambda^0[n] \rightarrow \Delta[n]$;
4. \mathcal{C} is an ∞ -category and \mathcal{C} has extension property with respect to all inclusions $\Lambda^n[n] \rightarrow \Delta[n]$.

Next we talk about the “underlying ∞ -groupoid of an ∞ -category”.

Proposition 2.1.2.15. Suppose that \mathcal{C} is an ∞ -category. Let $\mathcal{C}' \subseteq \mathcal{C}$ be the largest simplicial subset of \mathcal{C} such that all edges of \mathcal{C}' are invertible in \mathcal{C} . Then \mathcal{C} is a Kan complex. Moreover for any Kan complex K , the induced map $\mathrm{Hom}_{\mathbf{SSet}}(K, \mathcal{C}') \rightarrow \mathrm{Hom}_{\mathbf{SSet}}(K, \mathcal{C})$ is an isomorphism. We call \mathcal{C}' the **largest Kan complex** contained in \mathcal{C} . If $f : \mathcal{C} \rightarrow \mathcal{D}$ is a categorical equivalence between ∞ -categories, then the induced map $f' : \mathcal{C}' \rightarrow \mathcal{D}'$ between the largest Kan complexes is a homotopy equivalence. ⁴

⁴The lecturer cannot figure out a proof to the last assertion while not using the Joyal model structure of simplicial sets, although the book said that it can be proved directly.

We next consider diagrams on ∞ -categories. This gives rise to the difference between homotopy-commutative diagrams and homotopy-coherent diagrams, as it has been promised that we will discuss it in this section.

The issue is as follows. Suppose \mathcal{C} is an ∞ -category (or a topological category, simplicial category). To a first approximation, it is very similar to work on \mathcal{C} and $\mathbf{h}\mathcal{C}$, since they have the same objects and morphisms. However, in \mathcal{C} , we should ask whether or not two morphisms are equivalent or homotopic, but not equal. In this case, the homotopy between the morphisms need to be taken in consider, since they may as well affect the higher-dimensional homotopies. However, a commutative diagram in $\mathbf{h}\mathcal{C}$, does not contain the information on the homotopies between the morphisms. Consequently, commutative diagrams on $\mathbf{h}\mathcal{C}$, which corresponds to the notion of **homotopy-commutative diagrams** on \mathcal{C} , is quite unnatural and need to be refined by the notion of **homotopy-coherent diagrams** on \mathcal{C} .

Let us give two examples to show their differences.

Example 2.1.2.16. Consider a monoid object G in $\underline{\mathbf{H}}$, and consider the $\underline{\mathbf{H}}$ -enriched category $\underline{\mathbf{B}}G$. Then a $\underline{\mathbf{B}}G$ -valued diagram in a topological category is a homotopy-commutative diagram. On the other hand, the multiplication on G is homotopy commutative, i.e. the maps $(xy)z, x(yz) : G \times G \times G \rightarrow G$ are homotopic, which means that G is an A_3 -algebra. However, in practice we would like to require that G is not just an A_3 -algebra, but also satisfies higher homotopy-coherent properties. For example, we would like the following diagram of homotopies between morphisms $G \times G \times G \times G \rightarrow G$ to be commutative up to a homotopy:

$$\begin{array}{ccc}
 (x(yz))w & \text{---} & x((yz)w) \\
 & \swarrow \quad \searrow & \\
 ((xy)z)w & & x(y(zw)) \\
 & \swarrow \quad \searrow & \\
 & (xy)(zw); &
 \end{array}$$


and also the following diagram of homotopies between homotopies between maps $G \times G \times G \times G \times G \rightarrow G$ to be commutative up to a homotopy:

$$\begin{array}{ccccccc}
 & & & & (((ab)c)d)e & \text{---} & ((ab)(cd))e \\
 & & & & \downarrow & & \downarrow \\
 & & ((a(bc))d)e & & & & \\
 & & \downarrow & & & & \\
 (a(((bc)d)e)) & \text{---} & & \text{---} & (a(b(cd)))e & & \\
 & \downarrow & & \downarrow & \downarrow & & \\
 a(((bc)d)e) & \text{---} & & \text{---} & a((b(cd))e) & & \\
 & \downarrow & & \downarrow & \downarrow & & \\
 & & ((ab)c)(de) & \text{---} & (ab)(c(de)) & & \\
 & & \downarrow & & \downarrow & & \\
 (a(bc))(de) & & & & a(b((cd)e)) & & \\
 & \downarrow & & \downarrow & \downarrow & & \\
 a((bc)(de)) & \text{---} & & \text{---} & a(b(c(de))) & & \\
 & & & & \downarrow & & \\
 & & & & a(b(c(de))) & &
 \end{array}$$

This requirement is equivalent to assuming that G is an A_∞ -algebra. But there exists A_3 algebras that are not A_∞ -algebras. For such G , $\underline{\mathbf{B}}G$ -valued diagrams are “bad” since they cannot recover

all the homotopy relations that we want. Also, because of the lack of higher homotopies, the $\underline{\mathbf{H}}$ -enriched category $\underline{\mathbf{B}}G$ should not be viewed as an ∞ -category, which corresponds to the fact that not all $\underline{\mathbf{H}}$ -enriched categories are $(\infty, 1)$ -categories, as it has been mentioned in the previous section.

Example 2.1.2.17. Now suppose that we have an $[n]$ -valued diagram in the simplicial category \mathbf{SSet} , that we wish to be homotopy-coherent. We denote this diagram by F . For simplicity, we suppose that $F(i) = *$ for every $0 \leq i < n$. (The reader should also think of what will happen if $F(i)$ are arbitrary simplicial sets, as we will come back to this in several sections.) We define e_i to be the image of $F(i)$ in $F(n)$. Then, for every $0 \leq i \leq j < n$, since the maps $i \rightarrow j \rightarrow n$ and $i \rightarrow n$ are equal, we would like to assume that there exists a 1-simplex e_{ij} from e_i to e_j in $F(n)$. Also, for every $0 \leq i \leq j \leq k < n$, since the diagram



is commutative, we would like to assume that there exists a 2-simplex e_{ijk} in $F(n)$ filling the homotopies e_{ij}, e_{ik}, e_{jk} ; and for every $0 \leq i \leq j \leq k \leq l < n$, we would like to assume that there exists a 3-simplex e_{ijkl} in $F(n)$ filling the homotopies $e_{ijk}, e_{ijl}, e_{ikl}, e_{jkl}$; and homotopies of higher degrees as well. Such a diagram contains much more information than a homotopy-commutative diagram, where the latter only contains the information of e_{ij} 's, and losing the informations of higher homotopies.

So how shall we define a homotopy-coherent diagram? We could define this by working degree by degree in topological categories or simplicial categories. However, this is a great amount of data and is really difficult to actually work with. But in ∞ -categories, things are really easy to formulate: Suppose \mathcal{I} is an ordinary category and \mathcal{C} is an ∞ -category, we may simply define an \mathcal{I} -valued (homotopy-coherent) diagram in \mathcal{C} to be a map $N(\mathcal{I}) \rightarrow \mathcal{C}$ of simplicial sets. (This is, in fact, one of the reasons that quasi-categories are one of the most widely-used model for $(\infty, 1)$ -categories.)

The notion of homotopy-coherent diagrams in higher categories is a special case of functors between higher categories. Again, this is very easy to define in the settings of ∞ -categories:

Definition 2.1.2.18. Suppose \mathcal{C}, \mathcal{D} are ∞ -categories. A **functor** from \mathcal{C} to \mathcal{D} is simply a map $F : \mathcal{C} \rightarrow \mathcal{D}$ of simplicial sets. We define $\mathbf{Fun}(K, \mathcal{D}) := \mathcal{D}^K$, for every simplicial set K . (In this notation, K will often, but not always, be a simplicial set.)

The following proposition will be proved in Section [ERROR! SECTION NOT FOUND].

Proposition 2.1.2.19. Suppose K, K' are simplicial sets, \mathcal{C}, \mathcal{D} are ∞ -categories. Then:

- i) $\mathbf{Fun}(K, \mathcal{C})$ is an ∞ -category;
- ii) If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a categorical equivalence, then the induced map $\mathbf{Fun}(K, \mathcal{C}) \rightarrow \mathbf{Fun}(K, \mathcal{D})$ is a categorical equivalence;
- iii) If $f : K \rightarrow K'$ is a categorical equivalence, then the induced map $\mathbf{Fun}(K', \mathcal{C}) \rightarrow \mathbf{Fun}(K, \mathcal{C})$ is a categorical equivalence.

Definition 2.1.2.20. Suppose \mathcal{C}, \mathcal{D} are ∞ -categories. We define the ∞ -category of functors from \mathcal{C} to \mathcal{D} to be $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$. We define **natural transformations** between functors from \mathcal{C} to \mathcal{D} to be morphisms in $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$, and call a natural transformation a **natural equivalence** if it is invertible in $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$.