


Divide-and-Conquer :

 $4 - 1x$

a table S

* $C_2^n = O(n^2)$ possible $S[i, j]$

- * compute all $S[i, j]$ in $O(n^3)$ time

1	2	3	4	5	6	7	8	9	10	11	12
-7	8	-5	20	-3	-8	-23	18	20	-7	12	-5

take $O(j - i + 1) = O(n)$ time

 $4 - 2x$

i $S[i, j]$ j

	1	2	3	4	5	6	7	8	9	10	11	12
A	-7	8	-5	20	-3	-8	-23	18	20	-7	12	-5

take $O(j - i + 1)$ time = $O(n)$ time

len l # of pairs

1 n
 2 $n-1$
 3 $n-2$
 4 $n-3$
 ⋮
 n 1

$$\begin{aligned}
 T(n) &= \sum_{1 \leq l \leq n} (n - l + 1) \times l \\
 &= (n+1) \sum l - \sum l^2 \\
 &\cong (1/6) n^3 \\
 &= O(n^3)
 \end{aligned}$$

4-2y

(i_c, j_c, s_c)
 $A[2, 11] \quad (32)$

$p = 1$ $q = 6$ $r = 12$

	1	2	3	4	5	6	7	8	9	10	11	12
A	-7	8	-5	20	-3	-8	-23	18	20	-7	12	-5

$A[2, 4]$
(23)
 $(i_1, j_1, s_1) = (2, 4, 23)$
 $A[8, 11]$
(43)
 $(i_2, j_2, s_2) = (8, 11, 43)$

⇒ $A[8, 11]$ is the answer !

4-3x

Addition of n -bit numbers

$$\begin{array}{r}
 \overbrace{1011010110011}^n \\
 + 0010100010110 \\
 \hline
 \end{array}$$

$\begin{array}{c} 1 \\ \swarrow \\ 0 \\ \swarrow \\ 1 \end{array}$

⇒ $O(n)$ time

Multiplication of n -bit numbers

$$\begin{array}{r}
 \overbrace{1011}^n \\
 \times 1001 \\
 \hline
 1011 \\
 0000 \\
 0000 \\
 + 1011 \\
 \hline
 \text{sum of } n \text{ numbers}
 \end{array}$$

$\left. \begin{array}{c} \\ \\ \\ \end{array} \right\} n$

⇒ $O(n^2)$ time

$$X = 123456$$

$$Y = 789555$$

$$X \cdot Y = (123 \times 10^3 + 456) \cdot (789 \times 10^3 + 555)$$

$$= \underbrace{123 \times 789}_{①} \cdot \underbrace{10^6}_{①} + \underbrace{(123 \times 555 + 456 \times 789)}_{②} \cdot \underbrace{10^3}_{③} + \underbrace{456 \times 555}_{④}$$

$$= 96678 \cdot 10^6 + (68265 + 359784) \cdot 10^3 + 253080$$

$$= 96678 \cdot 10^6 + 428049 \cdot 10^3 + 253080$$

$$= 96678000000 + 428049000 + 253080$$


$$= 97475302080$$

One big * can be replaced by $\left\{ \begin{array}{l} 4 \text{ smaller } * \text{ and} \\ 3 + (\text{and } 2 \text{ shift}) \end{array} \right.$

$$(1) X \cdot Y = (A \cdot 2^{n/2} + B) \cdot (C \cdot 2^{n/2} + D)$$

$$= \underbrace{AC}_{①} \cdot 2^n + \underbrace{(AD + BC)}_{②} \cdot 2^{n/2} + \underbrace{BD}_{④}$$

n bits



$$T(n) = 4 T(n/2) + O(n) = O(n^2)$$

(3+, 2 shift)

$$(2) \text{ Let } P = AC, Q = BD, R = (A + B)(C + D)$$

$$X \cdot Y = \underbrace{P}_{①} \cdot 2^n + \underbrace{(R - P - Q)}_{② + ③} \cdot 2^{n/2} + \underbrace{Q}_{④}$$

$$T(n) = 3 T(n/2) + O(n) = O(n^{\log_2 3})$$

(6+, 2 shift)

Matrix multiplication

$$C = A \times B$$

$$c_{ij} = \sum_k \{ a_{ik} \times b_{kj} \}$$

$$i \begin{bmatrix} \boxed{8} \\ \vdots \\ j \end{bmatrix} = i \begin{bmatrix} 1 & 2 & 3 \\ \vdots & \vdots & \vdots \\ j \end{bmatrix} \times \begin{bmatrix} 1 \\ 2 \\ 1 \\ \vdots \\ j \end{bmatrix}$$

Time: $O(n^3)$

Matrix addition

$$C = A + B$$

$$c_{ij} = a_{ik} + b_{kj}$$

Time: $O(n^2)$

$$C = A \times B$$

$$\begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{k2} & C_{33} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{k2} & A_{33} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{21} & B_{22} & B_{23} \\ B_{31} & B_{32} & B_{33} \end{bmatrix}$$

$\frac{n}{3} * \frac{n}{3}$

$$C_{ij} = \sum_{1 \leq s \leq 3} A_{is} B_{sj} \Rightarrow T(n) = 3^3 T(n/3) + O(n^2) = O(n^3)$$

$$\Rightarrow T(n) = ? q T(n/3) + O(n^2) \quad (q < 3^3)$$

$$\Rightarrow T(n) = O(n^{\log_3 q}) \text{ by Master Thm.}$$

4-9x

$$C = A \times B \quad 4-9a$$

$$\begin{bmatrix} C_{11} & C_{12} & & C_{1k} \\ C_{21} & C_{22} & & C_{2k} \\ & & \dots & \\ C_{k1} & C_{k2} & & C_{kk} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} & & A_{1k} \\ A_{21} & A_{22} & & A_{2k} \\ & & \dots & \\ A_{k1} & A_{k2} & & A_{kk} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} & & B_{1k} \\ B_{21} & B_{22} & & B_{2k} \\ & & \dots & \\ B_{k1} & B_{k2} & & B_{kk} \end{bmatrix}$$

$\frac{n}{k} * \frac{n}{k}$

$$C_{ij} = \sum_{1 \leq s \leq k} A_{is} B_{sj} \Rightarrow T(n) = k^3 T(n/k) + O(n^2)$$

$$\Rightarrow T(n) = ? q T(n/k) + O(n^2) \quad (q < k^3)$$

$$\Rightarrow T(n) = O(n^{\log_k q}) \text{ by Master Thm.}$$

" \leq " (upper bound)

* $f(n) = O(g(n))$ if we can find positive constants c and n_0
s.t. $0 \leq f(n) \leq cg(n)$ for all $n \geq n_0$

* Prove that $3n^2 + 6n = O(n^2)$
 f g

Find c and n_0 such that

$$3n^2 + 6n \leq cn^2 \text{ for all } n \geq n_0$$

f cg

$$\Rightarrow \text{Choose } \begin{cases} c = 9 \\ n_0 = 1 \end{cases} \text{ or } \begin{cases} c = 4 \\ n_0 = 6 \end{cases}$$

4-10x

$T(n) = 2T(\lfloor n/2 \rfloor) + n$ (with $T(1) = 1$), find an upper bound

Claim: $T(n) = O(n \lg n)$
($\exists c$ and n_0 such that $T(n) \leq cn \lg n$ for all $n \geq n_0$)

Claim: $T(n) = O(n \lg n)$ since $T(n) \leq 3n \lg n$ for all $n \geq 2$
 c n_0

Problem: How can we verify the solution of a recurrence?

\Rightarrow By induction

Outline of Proof

Basis: ($n = n_0 = 2$) Show that $T(n) \leq 3n \lg n$ is true for $n = n_0 = 2$.

Induction: ($n > n_0$) Assume $T(x) \leq 3x \lg x$ for $x = 2, 3, \dots, n-1$.
Show that $T(n) \leq 3n \lg n$ is true for n as well.

4-10y

$$T(n) = 2T(\lfloor n/2 \rfloor) + n \quad (\text{with } T(1) = 1)$$

4-10a

Claim:

$$T(n) = O(n \lg n) \text{ since}$$

$$T(n) \leq 3n \lg n \text{ for all } n \geq 2$$

$\nwarrow c$
 $\nearrow n_0$

Basis: ($n = n_0 = 2$)

$$T(n_0) = T(2) = 2T(1) + 2 = 4$$

$$3n_0 \lg n_0 = 3 \cdot 2 \lg 2 = 6$$

$$T(2) = 4 \leq 6$$

\Rightarrow OK!

4-10b

Induction: ($n > 2$)

Assume $T(x) \leq 3x \lg x$
for $x = 2, 3, \dots, n-1$.

$$\begin{aligned}
 T(n) & \overset{\text{wavy arrow}}{\sim} x = \lfloor n/2 \rfloor \leq n-1 \\
 &= 2T(\lfloor n/2 \rfloor) + n \\
 &\leq 2(3 \lfloor n/2 \rfloor \lg \lfloor n/2 \rfloor) + n \\
 &\leq 3n \lg(n/2) + n \\
 &\leq 3n \lg n - 3n \lg 2 + n \\
 &\leq 3n \lg n - 3n + n \\
 &\leq 3n \lg n - 2n \\
 &\text{-----} \\
 &\leq 3n \lg n \text{ (goal !)} \quad \text{OK!}
 \end{aligned}$$

\Rightarrow Done !

$\Rightarrow T(n) = O(n \lg n)$

$$A(1) = 1, A(2) = 1, A(n) = A(n-1) + A(n-2) \text{ for } n \geq 3$$

Prove $A(n) \leq 2^{n-2}$ for $n \geq 2$

By induction

Basis: $n_0 = 2, 3$

$$A(2) = 1 \leq 2^{2-2} = 1$$

OK!

$$A(3) = A(1) + A(2) = 2 \leq 2^{3-2} = 2 \quad \text{OK!}$$

multi-value !!!

Induction: ~~(for $n > 2$)~~ (for $n > 3$)

Assume $T(x) \leq 2^{x-2}$ for $x = 2, 3, \dots, n-1$

$$A(n) = A(n-1) + A(n-2) \quad /* \underline{n-1, n-2 \in 2, 3, \dots, n-1}$$

$$\leq 2^{n-3} + 2^{n-4}$$

$$\leq 2 \times 2^{n-3}$$

$$\leq 2^{n-2}$$

Done!

Wrong for $n = 3$!!!

4-10z

4-10b

Induction: ~~($n > 2$)~~ (for $n > 3$)

Assume $T(x) \leq 3 \times \lg x$

for $x = 2, 3, \dots, n-1$.

$$T(n) \quad \rightsquigarrow \quad x = \lfloor n/2 \rfloor \leq n-1$$

$$= 2T(\lfloor n/2 \rfloor) + n \quad \lfloor n/2 \rfloor \geq 2$$

$$\leq 2(3 \lfloor n/2 \rfloor \lg \lfloor n/2 \rfloor) + n$$

$$\leq 3n \lg(n/2) + n$$

$$\leq 3n \lg n - 3n \lg 2 + n$$

$$\leq 3n \lg n - 3n + n$$

$$\leq 3n \lg n - 2n$$

$$\leq 3n \lg n \quad (\text{goal!}) \quad \text{OK!}$$

⇒ Done!

$$T(n) = 2T(\lfloor n/2 \rfloor) + n \quad (\text{with } T(1) = 1)$$

4-10a

Claim:

$$T(n) = O(n \lg n) \text{ since}$$

$$T(n) \leq c n \lg n \text{ for all } n \geq n_0$$

Basis: ~~$(n = n_0 = 2)$~~ $(n = 2, 3)$

$$T(n_0) = T(2) = 2T(1) + 2 = 4$$

$$c n_0 \lg n_0 = c \cdot 2 \lg 2 = 2c$$

$$T(2) = 4 \leq 2c$$

⇒ OK!

$$c \cdot 3 \lg 3 \sim 13.5c$$

$$T(3) = 5 \leq 13.5c$$

⇒ OK!

$$T(n) = 2T(\lfloor n/2 \rfloor) + n \quad (\text{with } T(1) = 1)$$

4-10a

Claim:

$$T(n) = O(n \lg n) \text{ since}$$

$$T(n) \leq c n \lg n \text{ for all } n \geq n_0$$

Basis: ~~$(n = n_0 = 2)$~~ $(n = 2, 3)$

$$T(n_0) = T(2) = 2T(1) + 2 = 4$$

$$c n_0 \lg n_0 = c \cdot 2 \lg 2 = 2c$$

$$T(2) = 4 \leq 2c$$

⇒ OK!

$$c \cdot 3 \lg 3 \sim 13.5c$$

$$T(3) = 5 \leq 13.5c$$

⇒ OK!

Claim:

$$T(n) = O(n \lg n) \quad (\exists c \text{ and } n_0 \text{ s.t.})$$

$$T(n) \leq c n \lg n \text{ for all } n \geq n_0$$

Basis: $(n = n_0)$

$$n_0 = 1? \quad T(1) = 1 \leq c \cdot 1 \lg 1? \quad (\times)$$

$$n_0 = 2? \quad T(2) = 4 \leq c \cdot 2 \lg 2?$$

$$\text{OK for } c \geq 4/(2 \lg 2) = 2$$

$$n_0 = 3? \quad T(3) = 5 \leq c \cdot 3 \lg 3?$$

$$\text{OK for } c \geq 5/(3 \lg 3)$$

$$T(n_0) \leq c n_0 \lg n_0$$

$$\Rightarrow \text{OK for } \begin{cases} n_0 \geq 2 \\ c \geq T(n_0)/n_0 \lg n_0 \end{cases} \quad \textcircled{1}$$

Induction: ~~$(n > n_0)$~~ (for $n > 3$)

Assume $T(x) \leq 3x \lg x$

for $x = 2, 3, \dots, n-1$.

$$\begin{aligned}
 T(n) & \xrightarrow{x = \lfloor n/2 \rfloor \leq n-1} 2T(\lfloor n/2 \rfloor) + n \quad \lfloor n/2 \rfloor \geq 2 \\
 & \leq 2(3\lfloor n/2 \rfloor \lg \lfloor n/2 \rfloor) + n \\
 & \leq 3n \lg(n/2) + n \\
 & \leq 3n \lg n - 3n \lg 2 + n \\
 & \leq 3n \lg n - 3n + n \\
 & \leq 3n \lg n - 2n \\
 & \text{-----} \\
 & \leq 3n \lg n \quad (\text{goal!}) \quad \text{OK!}
 \end{aligned}$$

⇒ Done!

Induction: $(n > n_0)$

Assume $T(x) \leq c x \lg x$

for $x = n_0, n_0+1, \dots, n-1$.

$$\begin{aligned}
 T(n) & \xrightarrow{\textcircled{1} n_0 \leq \lfloor n/2 \rfloor \leq n-1} 2T(\lfloor n/2 \rfloor) + n \\
 & \leq 2(c\lfloor n/2 \rfloor \lg \lfloor n/2 \rfloor) + n \\
 & \leq c n \lg(n/2) + n \\
 & \leq c n \lg n - c n \lg 2 + n \\
 & \leq c n \lg n - c n + n \\
 & \leq c n \lg n - (c-1)n \\
 & \text{-----} \\
 & \leq c n \lg n \quad (\text{goal!})
 \end{aligned}$$

⇒ OK for $c \geq 1$ ②

① ∩ ① ∩ ② ≠ ∅ ⇒ Done!

Basis: $(n = n_0)$

$$n_0 = 1? \quad T(1) = 1 \leq c \cdot 1 \lg 1? \quad (*)$$

$$n_0 = 2? \quad T(2) = 4 \leq c \cdot 2 \lg 2?$$

OK for $c \geq 4/(2 \lg 2) = 2$

$$n_0 = 3? \quad T(3) = 5 \leq c \cdot 3 \lg 3?$$

OK for $c \geq 5/(3 \lg 3)$

$$T(n_0) \leq c n_0 \lg n_0$$

⇒ OK for $\begin{cases} n_0 \geq 2 \\ c \geq T(n_0)/n_0 \lg n_0 \end{cases}$ ①

Induction: $(n > n_0)$

⇒ OK for $c \geq 1$ ②

$$\textcircled{1} n_0 \leq \lfloor n/2 \rfloor \leq n-1$$

$$\textcircled{1} n_0 = 2 \Rightarrow \begin{aligned} & n \geq 4 \\ & \text{basis: } 2, 3 \end{aligned}$$

$$\textcircled{1} \begin{cases} 2 \text{ ok for } c \geq 2 \\ 3 \text{ ok for } c \geq 5/(3 \lg 3) \sim 1.1 \end{cases} \Rightarrow c \geq 2$$

$$\textcircled{1} \cap \textcircled{1} \cap \textcircled{2} \Rightarrow n_0 = 2, c = 2$$

$$\textcircled{1} n_0 = 3 \Rightarrow \begin{aligned} & n \geq 6 \\ & \text{basis: } 3, 4, 5 \end{aligned}$$

$$\textcircled{1} \begin{cases} 3 \text{ ok for } c \geq 1.1 \\ 4 \text{ ok for } c \geq 1.08 \\ 5 \text{ ok for } c \geq 1.07 \end{cases}$$

$$\Rightarrow c \geq 1.1$$

$$\textcircled{1} \cap \textcircled{1} \cap \textcircled{2} \Rightarrow n_0 = 3, c = 1.5$$

$$\textcircled{1} n_0 = 4 \Rightarrow \begin{aligned} & n \geq 8 \\ & \text{basis: } 4, 5, 6, 7 \end{aligned}$$

$$T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \text{ (with } T(1) = 1)$$

4-12a

Prove $T(n) = 2n - 1$ (By induction)

Basis: $n = 1$

$$T(1) = 1 = 2 * 1 - 1 \text{ OK!}$$

Induction:

Assume $T(x) = 2x - 1$ for $x < n$.

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \\ &= (2 \times \lfloor n/2 \rfloor - 1) + (2 \times \lceil n/2 \rceil - 1) + 1 \\ &= 2n - 1 \text{ (goal!) OK!} \end{aligned}$$

⇒ Done!

Prove $T(n) \leq 2n$

Basis: $n = 1$

$$T(1) = 1 \leq 2 * 1 \text{ OK!}$$

Induction:

Assume $T(x) \leq 2x$ for $x < n$.

$$\begin{aligned} T(n) &= T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + 1 \\ &\leq 2 \times \lfloor n/2 \rfloor + 2 \times \lceil n/2 \rceil + 1 \\ &\leq 2n + 1 \\ &\quad \swarrow \text{---} \\ &\leq 2n \text{ (goal!)} \end{aligned}$$

⇒ Fail! Why???

$$T(n) = T(n-1) + 1 \text{ (with } T(1) = 1) \quad * \text{ Answr: } T(n) = n$$

Prove $T(n) = 1$ (By induction)

Basis: $n = 1$

$$T(1) = 1 \text{ OK!}$$

Induction:

Assume $T(x) = 1$ for $x = 1, 2, \dots, n - 1$.

$$\begin{aligned} T(n) &= T(n-1) + 1 \\ &= 1 + 1 \\ &= 2 \end{aligned}$$

⇒ We can not prove $T(n) = 1$.
But, we conclude that $T(n) = 2$. ???!!!

4-13x

Find upper bound of $T(n) = T(n - 1) + 2n$ (with $T(1) = 1$)

$$T(x) = 2x + T(x - 1)$$

$$\begin{aligned}
 T(n) &= 2n + \underline{T(n - 1)} \\
 &\leq 2n + 2(n-1) + \underline{T(n - 2)} \\
 &\leq 2n + 2(n-1) + 2(n-2) + \underline{T(n - 3)} \\
 &\quad \vdots \\
 &\leq 2n + 2(n-1) + 2(n-2) + \dots + 2(2) + T(1) \\
 &\leq 2(2 + 3 + \dots + n) + 1 \\
 &\leq 2(1 + 2 + 3 + \dots + n) - 1 \\
 &\leq n(n+1) - 1 \\
 &= O(n^2)
 \end{aligned}$$

4-14x


Find upper bound of $T(n) = 3T(\lfloor n/4 \rfloor) + n$ (with $T(1/0) = \theta(1)$)

$$\begin{aligned}
 T(n) &= n + 3\underline{T(\lfloor n/4 \rfloor)} \quad T(x) = x + 3T(\lfloor x/4 \rfloor) \\
 &\leq n + 3\{ \lfloor n/4 \rfloor + 3T(\lfloor \lfloor n/4 \rfloor / 4 \rfloor) \} \\
 &\leq n + 3\lfloor n/4 \rfloor + 3^2 \underline{T(\lfloor n/4^2 \rfloor)} \\
 &\leq n + 3\lfloor n/4 \rfloor + 3^2\{ \lfloor n/4^2 \rfloor + 3T(\lfloor \lfloor n/4^2 \rfloor / 4 \rfloor) \} \\
 &\leq n + 3\lfloor n/4 \rfloor + 3^2\lfloor n/4^2 \rfloor + 3^3 \underline{T(\lfloor n/4^3 \rfloor)} \\
 &\quad \vdots \\
 &\leq n + 3\lfloor n/4 \rfloor + 3^2\lfloor n/4^2 \rfloor + 3^3\lfloor n/4^3 \rfloor + \dots + 3^k \underline{T(\lfloor n/4^k \rfloor)} \\
 &\quad T(1) \text{ or } T(0) \\
 &\leq n + 3n/4 + 3^2n/4^2 + 3^3n/4^3 + \dots + 3^k \theta(1) \\
 &\leq O(n) + 3^k \theta(1)
 \end{aligned}$$

$$\frac{n}{4^k} \leq 1 \Rightarrow k \geq \lg_4 n$$

等 比 會 收 斂 !

4-14y



4-14z

Assume that we do not take ceiling for both.

4-15a

- * cost(**internal-nodes**) of a level is at most n .
 (Prove that $n/3 + 2n/3 \leq n$ for $\square\square\square$, $\square\square\square$, and $\square\square\square$.)

* $n/3 + 2n/3 \leq n$ for $\lfloor \rfloor \lfloor \rfloor$, $\lfloor \rfloor \lceil \rceil$, and $\lceil \rceil \lfloor \rfloor$

Proof for $\lfloor \rfloor \lceil \rceil$

Case 1. $n = 3k$

$$\lfloor n/3 \rfloor + \lceil 2n/3 \rceil = k + 2k = 3k = n$$

Case 2. $n = 3k + 1$

$$\lfloor n/3 \rfloor + \lceil 2n/3 \rceil = k + \lceil 2k + 2/3 \rceil = k + 2k + 1 = 3k + 1 = n$$

Case 3. $n = 3k + 2$

$$\lfloor n/3 \rfloor + \lceil 2n/3 \rceil = k + \lceil 2k + 1 + 1/3 \rceil = k + 2k + 2 = 3k + 2 = n$$

4-15x

Assume that we do not take ceiling for both.

4-15a

* cost(internal-nodes) of a level is at most n .
(Prove that $n/3 + 2n/3 \leq n$ for $\lfloor \rfloor \lfloor \rfloor$, $\lfloor \rfloor \lceil \rceil$, and $\lceil \rceil \lfloor \rfloor$.)

How to compute the number of leaves L ?

* From $L \leq 2^{\lg(3/2)n}$, we have $L = \omega(\lg n)$, which is larger than $O(\lg n)$. (太悲觀)

$$2^{\lg(3/2)n} = n^{\lg(3/2)} = O(n^{1.585})$$

* Avoid the computation of L . (text book)
Prove $O(\lg n)$ is correct by substitution method.

* Prove by induction that $L \leq n$.
(Try it!)

* If we take ceiling for both, $\lceil n/3 \rceil + \lceil 2n/3 \rceil$ may be larger than n . For example, 11 is partitioned into 4 and 8. Then,

(a) cost(internal-nodes) of a level may be larger than n .


(b) The number of leaves, L , is unknown.

(c) Using substitution method, $T(n) = O(n \lg n)$ still can be proved. (with some effort!)

* Usually, cost(leaves) can be ignored; however, we should consider it in a formal proof.

4-15y

4-16a



$O: \leq$ (upper bound) $\Theta: =$ (tight bound) $\Omega: \geq$ (lower bound)



Prove upper bound

Prove tight bound

Prove lower bound

$$\begin{aligned} & T(n) \\ & \leq A1 \\ & \leq A2 \\ & \leq A3 \\ & = O(A3) \end{aligned}$$

(1) prove upper bound
(2) prove lower bound

$$\begin{aligned} & T(n) \\ & \geq A1 \\ & \geq A2 \\ & \geq A3 \\ & = \Omega(A3) \end{aligned}$$

過程中可多算不可少算

過程中可少算不可多算

Find **upper bound** of $T(n) = 3T(\lfloor n/4 \rfloor) + n$

(page 4-14)

$$T(n) = n + 3T(\lfloor n/4 \rfloor) \quad T(x) = x + 3T(\lfloor x/4 \rfloor)$$

$$\begin{aligned}
 & \leq n + 3\{ \lfloor n/4 \rfloor + 3T(\lfloor \lfloor n/4 \rfloor / 4 \rfloor) \} \\
 & \leq n + 3\lfloor n/4 \rfloor + 3^2 T(\lfloor n/4^2 \rfloor) \\
 & \leq n + 3\lfloor n/4 \rfloor + 3^2 \{ \lfloor n/4^2 \rfloor + 3T(\lfloor \lfloor n/4^2 \rfloor / 4 \rfloor) \} \\
 & \vdots \\
 & \leq n + 3\lfloor n/4 \rfloor + 3^2 \lfloor n/4^2 \rfloor + 3^3 \lfloor n/4^3 \rfloor + \dots + 3^k T(\lfloor n/4^k \rfloor) \\
 & \quad T(1) \text{ or } T(0) \\
 & \leq n + 3n/4 + 3^2 n/4^2 + 3^3 n/4^3 + \dots + 3^k \theta(1) \\
 & \leq n(1 - (1/4)^k) / (1 - 1/4) + 3^k \theta(1) \\
 & \leq 4n + o(n) \\
 & = O(n)
 \end{aligned}$$

$$\frac{n}{4^k} \leq 1 \Rightarrow k \geq \lg_4 n$$

4-14Q

Find **lower bound** of $T(n) = 3T(\lfloor n/4 \rfloor) + n$

$$T(n) = n + 3T(\lfloor n/4 \rfloor)$$

$$\begin{aligned}
 & \geq n + 3\{ \lfloor n/4 \rfloor + 3T(\lfloor \lfloor n/4 \rfloor / 4 \rfloor) \} \\
 & \geq n + 3\lfloor n/4 \rfloor + 3^2 T(\lfloor n/4^2 \rfloor) \\
 & \geq n + 3\lfloor n/4 \rfloor + 3^2 \{ \lfloor n/4^2 \rfloor + 3T(\lfloor \lfloor n/4^2 \rfloor / 4 \rfloor) \} \\
 & \vdots \\
 & \geq n + 3\lfloor n/4 \rfloor + 3^2 \lfloor n/4^2 \rfloor + 3^3 \lfloor n/4^3 \rfloor + \dots + 3^k T(\lfloor n/4^k \rfloor) \\
 & \quad T(1) \text{ or } T(0) \\
 & \geq n + 3(n/4 - 1) + 3^2(n/4^2 - 1) + 3^3(n/4^3 - 1) + \dots + 3^k \theta(1) \\
 & \geq n(1 - (1/4)^k) / (1 - 1/4) - (3 + 3^2 + \dots + 3^{k-1}) + 3^k \theta(1) \\
 & \geq n - 3^k + 3^k \theta(1) \\
 & \geq n - o(n) + o(n) \\
 & = \Omega(n)
 \end{aligned}$$

$$\frac{n}{4^k} \leq 1 \Rightarrow k \geq \lg_4 n$$

4-14Q

$O: \leq$ (upper bound)

Prove upper bound

$$\begin{aligned} & \stackrel{Q1}{=} T(n) \\ & \stackrel{Q2}{\leq} A1 \\ & \leq A2 \\ & \leq 5n^5 \\ & = O(n^5) \end{aligned}$$

$\xrightarrow{Q3} \Theta(n^5) ???$

$$\begin{aligned} & T(n) \\ & = 3n^4 + 2n \\ & \leq 5n^4 \\ & \leq 5n^5 \\ & = O(n^5) \end{aligned}$$

$\xrightarrow{Q3} \Theta(n^5) ???$

4-14Q

Source: <http://www.csd.uwo.ca/~moreno/CS433-CS9624/Resources/master.pdf>

22. $T(n) = T(n/2) + n(2 - \cos n)$

We are in Case 3, but the **regularity condition** is violated. (Consider $n = 2\pi k$, where k is odd and arbitrarily large. For any such choice of n , you can show that $c \geq 3/2$, thereby violating the regularity condition.)

4-16Q