
### Introduction to Audio Content Analysis

### Module 3.1: Feature Extraction — Statistical Features

#### alexander lerch





### introduction

overview



#### corresponding textbook section

Chapter 3 — Instantaneous Features: pp. 35-41

- lecture content
  - introduction to statistical features
    - mean
    - variance and standard deviation
    - quantiles
- learning objectives
  - give examples of where statistical features can be used
  - describe the meaning of the introduced statistical features



### introduction

overview



### corresponding textbook section

Chapter 3 — Instantaneous Features: pp. 35-41

#### lecture content

- introduction to statistical features
  - mean
  - variance and standard deviation
  - quantiles

#### learning objectives

- give examples of where statistical features can be used
- describe the meaning of the introduced statistical features



# statistical features introduction



- statistical features are...
  - numerical descriptors of statistical properties of the signal or its PDF
  - general features that are not audio-related

#### usage

often not directly applied to audio signal but to feature representations of it

# Georgia Center for Music Tech College of Design

• from signal x:


$$\mu_{\mathsf{x}}(n) = rac{1}{\mathcal{K}} \sum_{i=i_{\mathsf{s}}(n)}^{i_{\mathsf{e}}(n)} \mathsf{x}(i)$$

• from distribution p

$$\mu_{\mathsf{X}}(n) = \sum_{n=1}^{\infty} x \cdot p_{\mathsf{X}}(x)$$

from unnormalized distribution d

$$\mu_{x}(n) = \frac{\sum_{x=-\infty}^{\infty} x \cdot d_{x}(x)}{\sum_{x=-\infty}^{\infty} d_{x}(x)}$$



# Georgia Center for Music Technology

• from signal x:


$$\mu_{\mathsf{x}}(n) = \frac{1}{\mathcal{K}} \sum_{i=i_{\mathsf{s}}(n)}^{i_{\mathsf{e}}(n)} \mathsf{x}(i)$$

• from distribution  $p_x$ :

$$\mu_{\mathsf{x}}(\mathsf{n}) = \sum_{\mathsf{x} = -\infty}^{\infty} \mathsf{x} \cdot \mathsf{p}_{\mathsf{x}}(\mathsf{x})$$

• from unnormalized distribution d

$$\mu_{x}(n) = \frac{\sum_{x=-\infty}^{\infty} x \cdot d_{x}(x)}{\sum_{x=-\infty}^{\infty} d_{x}(x)}$$



## Georgia Center for Music Tech College of Design

from signal x:


$$\mu_{\mathsf{x}}(n) = rac{1}{\mathcal{K}} \sum_{i=i_{\mathsf{s}}(n)}^{i_{\mathsf{e}}(n)} \mathsf{x}(i)$$

• from distribution  $p_x$ :

$$\mu_{\mathsf{x}}(\mathsf{n}) = \sum_{-\infty}^{\infty} x \cdot \mathsf{p}_{\mathsf{x}}(\mathsf{x})$$

• from unnormalized distribution  $d_x$ :

$$\mu_{x}(n) = \frac{\sum_{x=-\infty}^{\infty} x \cdot d_{x}(x)}{\sum_{x=-\infty}^{\infty} d_{x}(x)}$$



#### geometric mean

$$M_{x}(0,n) = \sqrt[\kappa]{\prod_{i=i_{s}(n)}^{i_{e}(n)} x(i)}$$

$$= \exp\left(\frac{1}{\mathcal{K}} \sum_{i=i_{s}(n)}^{i_{e}(n)} \log[x(i)]\right)$$

harmonic mean

$$M_{x}(-1,n) = \frac{\mathcal{K}}{\sum_{i=i_{x}(n)}^{i_{e}(n)} \frac{1}{x} \times (i)}$$

#### geometric mean

$$M_{x}(0,n) = \sqrt[K]{\prod_{i=i_{s}(n)}^{i_{e}(n)} x(i)}$$

$$= \exp\left(\frac{1}{K} \sum_{i=i_{s}(n)}^{i_{e}(n)} \log[x(i)]\right)$$

harmonic mean

$$M_{x}(-1,n) = \frac{\mathcal{K}}{\sum_{i=i_{s}(n)} 1/x(i)}$$

generalized mean

#### generalized mean

$$M_{x}(\beta, n) = \sqrt[\beta]{rac{1}{\mathcal{K}}\sum_{i=i_{s}(n)}^{i_{e}(n)}x^{\beta}(i)}$$

- ullet eta=1: arithmetic mear
- $\beta = 2$ : quadratic mean
- $\beta = -1$ : harmonic mean
- $\beta \rightarrow 0$ : geometric mean
- $\beta \to -\infty$ : minimum
- $\beta \to \infty$ : maximum

generalized mean

### generalized mean

$$M_{x}(\beta, n) = \sqrt{\frac{1}{\mathcal{K}} \sum_{i=i_{s}(n)}^{i_{e}(n)} x^{\beta}(i)}$$

- $\beta = 1$ : arithmetic mean
- $\beta = 2$ : quadratic mean
- $\beta = -1$ : harmonic mean
- $\beta \rightarrow 0$ : geometric mean
- $\beta \to -\infty$ : minimum
- $\beta \to \infty$ : maximum

centroid: center of gravity

$$v_{\mathrm{C}}(n) = rac{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} \left(i-i_{\mathrm{s}}(n)
ight) \cdot x(i)}{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} x(i)}$$

statistical features — centroid

centroid: center of gravity

$$v_{\mathrm{C}}(n) = rac{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} \left(i-i_{\mathrm{s}}(n)
ight) \cdot x(i)}{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} x(i)}$$

Why does this look familiar?



statistical features — centroid



centroid: center of gravity

$$v_{\mathrm{C}}(n) = rac{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} \left(i-i_{\mathrm{s}}(n)
ight) \cdot x(i)}{\sum\limits_{i=i_{\mathrm{s}}(n)}^{i_{\mathrm{e}}(n)} x(i)}$$

#### Why does this look familiar?

ightarrow compare arithmetic mean



#### variance & standard deviation

measure of *spread* of the signal around its mean

- variance
  - from signal block:


$$\sigma_{\scriptscriptstyle X}^2(n) = rac{1}{\mathcal{K}} \sum_{i=i_{\scriptscriptstyle S}(n)}^{i_{\scriptscriptstyle E}(n)} ig(x(i) - \mu_{\scriptscriptstyle X}(n)ig)^2$$

from distribution

$$\sigma_x^2(n) = \sum_{x=0}^{\infty} (x - \mu_x)^2 \cdot p_x(x)$$

standard deviation

$$\sigma_{X}(n) = \sqrt{\sigma_{X}^{2}(n)}$$



### statistical features

## Georgia Center for Music Tech College of Design

variance & standard deviation

measure of spread of the signal around its mean

- variance
  - from signal block:


$$\sigma_{\scriptscriptstyle X}^2(n) = rac{1}{\mathcal{K}} \sum_{i=i_{\scriptscriptstyle S}(n)}^{i_{\scriptscriptstyle E}(n)} ig(x(i) - \mu_{\scriptscriptstyle X}(n)ig)^2$$

• from distribution:

$$\sigma_x^2(n) = \sum_{x=0}^{\infty} (x - \mu_x)^2 \cdot p_x(x)$$

standard deviation

$$\sigma_{\mathsf{x}}(n) = \sqrt{\sigma_{\mathsf{x}}^2(n)}$$



### statistical features

Georgia Center for Music Tech Techology

variance & standard deviation

measure of spread of the signal around its mean

- variance
  - from signal block:


$$\sigma_{\scriptscriptstyle X}^2(n) = rac{1}{\mathcal{K}} \sum_{i=i_{\scriptscriptstyle \mathrm{S}}(n)}^{i_{\scriptscriptstyle \mathrm{E}}(n)} ig( x(i) - \mu_{\scriptscriptstyle X}(n) ig)^2$$

• from distribution:

$$\sigma_x^2(n) = \sum_{x=0}^{\infty} (x - \mu_x)^2 \cdot p_x(x)$$

standard deviation

$$\sigma_{x}(n) = \sqrt{\sigma_{x}^{2}(n)}$$




# statistical features quantiles & quantile ranges



dividing the PDF into (equal sized) subsets

$$Q_{\mathrm{X}}(c_{p}) = \operatorname{argmin}\left(F_{\mathrm{X}}(x) \leq c_{p}\right)$$
 with  $F_{\mathrm{X}}(x) = \int\limits_{-\infty}^{x} p_{\mathrm{x}}(y) \, dy$ 



median

$$Q_{\rm X}(0.5) = \operatorname{argmin} \left( F_{\rm X}(x) \le 0.5 \right)$$

- quartiles:  $Q_X(0.25)$ ,  $Q_X(0.5)$ , and  $Q_X x(0.75)$
- quantile range, e.g.

$$\Delta Q_{\rm X}(0.9) = Q_{\rm X}(0.95) - Q_{\rm X}(0.05)$$

## Georgia Center for Music Tech Technology

#### statistical features

- summarize technical signal characteristics in few numerical values
- may be used on a time domain, frequency domain, or feature domain signal

#### feature description

- mean: average value
- variance and standard deviation: measure of expected deviation from the mean
- quantiles: numerical pdf shape description

