

Some contributions to the theory of distances

Frank Nielsen

Sony Computer Science Laboratories Inc., Tokyo, Japan

E-mail: `Frank.Nielsen@acm.org`

November 11, 2020

1 Calculating statistical distances, relative entropies, cross-entropies and entropies

- Cumulant-free closed-form formulas for some common (dis)similarities between densities of an exponential family (<https://arxiv.org/abs/2003.02469>)

The Bregman and Jensen divergences are defined for a strictly convex generator F by:

$$B_F(\theta_1 : \theta_2) := F(\theta_1) - F(\theta_2) - (\theta_1 - \theta_2)^\top \nabla F(\theta_2) \quad (1)$$

$$J_F(\theta_1 : \theta_2) := \frac{F(\theta_1) + F(\theta_2)}{2} - F\left(\frac{\theta_1 + \theta_2}{2}\right). \quad (2)$$

Since the Jensen and Bregman convex generators $F(\theta)$ are defined modulo an *affine term* $\langle a, \theta \rangle + b$ (i.e., $J_F(\theta_1 : \theta_2) = J_G(\theta_1 : \theta_2)$ and $B_F(\theta_1 : \theta_2) = B_G(\theta_1 : \theta_2)$ with $G(\theta) = F(\theta) + \langle a, \theta \rangle + b$), we can choose the equivalent generator $G(\theta) := -\log p_\theta(x) = F(\theta) - \langle t(x), \theta \rangle - k(x)$ (i.e., $a = -t(x)$) and $b = -k(x)$, and express the Kullback-Leibler divergence, the skewed Bhattacharyya divergences, the α -divergences and many other statistical distances between densities of a natural exponential family $\{p_\theta(x) = 1_{\mathcal{X}}(x) \exp(\langle t(x), \theta \rangle - F(\theta) + k(x))\}$ without *explicitly* using the log-normalizer $F(\theta) = \log(\int_{x \in \mathcal{X}} \exp(\langle t(x), \theta \rangle + k(x)) d\mu(x))$ of the exponential family (also called cumulant function or log-partition function).

For example, the *Bhattacharyya similarity coefficient* is expressed as:

$$\begin{aligned} \rho[p_{\theta_1}, p_{\theta_2}] &:= \int_{x \in \mathcal{X}} \sqrt{p_{\theta_1}(x) p_{\theta_2}(x)} d\mu(x), \\ &= \exp(-J_F(\theta_1 : \theta_2)) = \exp(-J_{-\log p_\theta(\omega)}(\theta_1 : \theta_2)), \quad \forall \omega \in \mathcal{X}, \\ &= \frac{p_{\bar{\theta}}(\omega)}{\sqrt{p_{\theta_1}(\omega) p_{\theta_2}(\omega)}}, \quad \forall \omega \in \mathcal{X}, \end{aligned}$$

where $\bar{\theta} := \frac{\theta_1 + \theta_2}{2}$. For generic exponential families parameterized by $\lambda(\theta)$ (i.e., not in natural form), we need to explicit the mid-parameter $\bar{\lambda} := \lambda(\bar{\theta})$ from the *partial* factorization of the exponential family (the λ -mean corresponding to the θ -mean).

For the Kullback-Leibler divergence, using the fact that $D_{\text{KL}}[p_{\theta_1} : p_{\theta_2}] = B_F[\theta_2 : \theta_1] = B_G[\theta_2 : \theta_1]$ (better written as $D_{\text{KL}}^*[p_{\theta_2} : p_{\theta_1}] = B_F[\theta_2 : \theta_1]$ where D_{KL}^* is the reverse divergence) with the equivalent generator $G(\theta) = -\log p_\theta(x)$, we get

$$D_{\text{KL}}[p_{\theta_1} : p_{\theta_2}] = \log\left(\frac{p_{\theta_1}(\omega)}{p_{\theta_2}(\omega)}\right) + (\theta_2 - \theta_1)^\top (t(\omega) - \nabla F(\theta_1)), \quad \forall \omega \in \mathcal{X}.$$

Choosing ω such that $t(\omega) = \nabla F(\theta_1) = E_{p_{\theta_1}}[t(x)] =: \eta_1$, we express the KLD as a log density ratio: $D_{\text{KL}}[p_{\theta_1} : p_{\theta_2}] = \log\left(\frac{p_{\theta_1}(\omega)}{p_{\theta_2}(\omega)}\right)$. In general we may need several ω_i 's so that $\frac{1}{s} \sum_i t(\omega_i) = \nabla F(\theta_1) = \eta_1$. Thus we get the three equivalent formula for the KLD between densities of an exponential family:

$$D_{\text{KL}}[p_{\lambda_1} : p_{\lambda_2}] = \int_{x \in \mathcal{X}} p_{\lambda_1}(x) \log\left(\frac{p_{\lambda_1}(x)}{p_{\lambda_2}(x)}\right) d\mu(x) \quad (3)$$

$$= B_F(\theta(\lambda_2) : \theta(\lambda_1)) \quad (\text{require } F(\theta), \nabla F(\theta)) \quad (4)$$

$$= \log\left(\frac{p_{\lambda_1}(\omega)}{p_{\lambda_2}(\omega)}\right) + (\theta(\lambda_2) - \theta(\lambda_1))^\top (t(\omega) - E_{p_{\lambda_1}}[t(x)]), \quad \forall \omega \in \mathcal{X} \quad (\text{require } E_{p_{\lambda_1}}[t(x)])$$

$$= \frac{1}{s} \sum_{i=1}^s \log\left(\frac{p_{\lambda_1}(\omega_i)}{p_{\lambda_2}(\omega_i)}\right), \quad (\text{require } \frac{1}{s} \sum_{i=1}^s t(\omega_i) = E_{p_{\lambda_1}}[t(x)]) \quad (6)$$

The last formula bears some similarity with the Monte-Carlo stochastic approximation of the Kullback-Leibler divergence:

$$\begin{aligned} x_1, \dots, x_n &\sim_{\text{iid}} p_{\lambda_1} \\ \tilde{D}_{\text{KL},n}[p_{\lambda_1} : p_{\lambda_2}] &:= \frac{1}{n} \sum_{i=1}^n \log\left(\frac{p_{\lambda_1}(x_i)}{p_{\lambda_2}(x_i)}\right) \\ \lim_{n \rightarrow \infty} \tilde{D}_{\text{KL},n}[p_{\lambda_1} : p_{\lambda_2}] &= D_{\text{KL}}[p_{\lambda_1} : p_{\lambda_2}] \\ \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n t(x_i) &= E_{p_{\lambda_1}}[t(x)] \end{aligned}$$

For example, we can write the KLD between two multivariate normal distributions as