

The α -representations of the Fisher Information Matrix

— On gauge freedom of the FIM —

Frank Nielsen
`Frank.Nielsen@acm.org`

19 September 2017
 Revised September 2020

The *Fisher Information Matrix* [1] (FIM) for a family of *parametric* probability models $\{p(x; \theta)\}_{\theta \in \Theta}$ (densities $p(x; \theta)$ expressed with respect to a positive base measure ν) indexed by a D -dimensional parameter vector $\theta := (\theta^1, \dots, \theta^D)$ is historically defined by

$$I(\theta) := [I_{ij}(\theta)], \quad I_{ij}(\theta) := E_{p(x; \theta)} [\partial_i l(x; \theta) \partial_j l(x; \theta)], \quad (1)$$

where $l(x; \theta) := \log p(x; \theta)$ is the *log-likelihood function*, and $\partial_i := \frac{\partial}{\partial \theta^i}$ (by notational convention). The FIM is a $D \times D$ positive semi-definite matrix for a D -order parametric family.

The FIM is a cornerstone in statistics and occurs in many places, like for example the celebrated *Cramér-Rao lower bound* [3] for an unbiased estimator $\hat{\theta}$:

$$\text{Var}_{p(x; \theta)}[\hat{\theta}] \succeq I^{-1}(\theta),$$

where \succeq denotes the Löwner @artial ordering of positive semi-definite matrices: $A \succeq B$ iff. $A - B \succ 0$ is positive semi-definite. Another use of the FIM is in gradient descent method using the *natural gradient* (see [6] for its use in deep learning).

Yet, it is common to encounter another equivalent expression of the FIM in the literature [3, 1]:

$$I'_{ij}(\theta) := 4 \int \partial_i \sqrt{p(x; \theta)} \partial_j \sqrt{p(x; \theta)} d\nu(x) \quad (2)$$

This form of the FIM is well-suited to prove that the FIM is always positive semi-definite matrix [1]: $I(\theta) \succeq 0$.

It turns out that one can define a family of equivalent representations of the FIM using the α -embeddings of the parametric family. We define the α -representation of densities $l^{(\alpha)}(x; \theta) := k_\alpha(p(x; \theta))$ with

$$k_\alpha(u) := \begin{cases} \frac{2}{1-\alpha} u^{\frac{1-\alpha}{2}}, & \text{if } \alpha \neq 1 \\ \log u, & \text{if } \alpha = 1. \end{cases} \quad (3)$$

The function $l^{(\alpha)}(x; \theta)$ is called the α -likelihood function.

The α -representation of the FIM (or α -FIM for short) is

$$I_{ij}^{(\alpha)}(\theta) := \int \partial_i l^{(\alpha)}(x; \theta) \partial_j l^{(-\alpha)}(x; \theta) d\nu(x) \quad (4)$$

In compact notation, we have $I_{ij}^{(\alpha)}(\theta) = \int \partial_i l^{(\alpha)} \partial_j l^{(-\alpha)} d\nu(x)$ (this is the α -FIM). We can expand the α -FIM expressions as follows

$$I_{ij}^{(\alpha)}(\theta) = \begin{cases} \frac{1}{1-\alpha^2} \int \partial_i p(x; \theta)^{\frac{1-\alpha}{2}} \partial_j p(x; \theta)^{\frac{1+\alpha}{2}} d\nu(x) & \text{for } \alpha \neq \pm 1 \\ \int \partial_i \log p(x; \theta) \partial_j p(x; \theta) d\nu(x) & \text{for } \alpha \in \{-1, 1\} \end{cases}$$

The proof that $I_{ij}^{(\alpha)}(\theta) = I_{ij}(\theta)$ follows from the fact that

$$\partial_i l^\alpha = p^{-\frac{\alpha+1}{2}} \partial_i p = p^{\frac{1-\alpha}{2}} \partial_i l,$$

since $\partial_i l = \frac{\partial_i p}{p}$.

Therefore we get

$$\partial_i l^{(\alpha)} \partial_j l^{(-\alpha)} = p \partial_i l \partial_j l,$$

and $I_{ij}^{(\alpha)}(\theta) = E[\partial_i l \partial_j l] = I_{ij}(\theta)$.

Thus Eq. 1 and Eq. 2 where two examples of the α -representation, namely the 1-representation and the 0-representation, respectively. The 1-representation of Eq. 1 is called the logarithmic representation, and the 0-representation of Eq. 2 is called the square root representation.

Note that $I_{ij}(\theta) = E[\partial_i l \partial_j l] = \int p \partial_i l \partial_j l d\nu(x) = \int \partial_i p \partial_j l d\nu(x) = I_{ij}^{(1)}(\theta)$ since $\partial_i l = \frac{\partial_i p}{p}$

In information geometry [1], $\{\partial_i l^{(\alpha)}\}_i$ plays the role of tangent vectors, the α -scores. Geometrically speaking, the tangent plane $T_{p(x; \theta)}$ can be described using any α -base. The statistical manifold $M = \{p(x; \theta)\}_\theta$ is imbedded into the function space $\mathbb{R}^{\mathcal{X}}$, where \mathcal{X} denotes the support of the densities.

Under regular conditions [3, 1], the α -representation of the FIM for $\alpha \neq -1$ can further be rewritten as

$$I_{ij}^{(\alpha)}(\theta) = -\frac{2}{1+\alpha} \int p(x; \theta)^{\frac{1+\alpha}{2}} \partial_i \partial_j l^{(\alpha)}(x; \theta) d\nu(x). \quad (5)$$

Since we have

$$\partial_i \partial_j l^{(\alpha)}(x; \theta) = p^{\frac{1-\alpha}{2}} \left(\partial_i \partial_j l + \frac{1-\alpha}{2} \partial_i l \partial_j l \right),$$

it follows that

$$I_{ij}^{(\alpha)}(\theta) = -\frac{2}{1+\alpha} \left(-I_{ij}(\theta) + \frac{1-\alpha}{2} I_{ij}(\theta) \right) = I_{ij}(\theta).$$

Notice that when $\alpha = 1$, we recover the equivalent expression of the FIM (under mild conditions)

$$I_{ij}^{(1)}(\theta) = -E[\nabla^2 \log p(x; \theta)].$$

In particular, when the family is an exponential family [5] with cumulant function $F(\theta)$, we have

$$I(\theta) = \nabla^2 F(\theta) \succ 0.$$

Similarly, the coefficients of the α -connection can be expressed using the α -representation as

$$\Gamma_{ij,k}^{(\alpha)} = \int \partial_i \partial_j l^{(\alpha)} \partial_k^{(-\alpha)} d\nu(x).$$

The Riemannian metric tensor g_{ij} (a geometric object) can be expressed in matrix form $I_{ij}^{(\alpha)}(\theta)$ using the α -base, and this tensor is called the Fisher metric tensor.

The FIM may further be represented using the more general (ρ, τ) -monotone embeddings [2]: Let ρ and τ be two strictly increasing functions, and f a strictly convex function such that $f'(\rho(u)) = \tau(u)$ (with f^* denoting its convex conjugate). Let us write $p_\theta(x) = p(x; \theta)$. Then we have ${}^{\rho, \tau}g(\theta) = [{}^{\rho, \tau}g_{ij}(\theta)]_{ij}$ with

$${}^{\rho, \tau}g_{ij}(\theta) = \int (\partial_i \rho(p_\theta(x))) (\partial_j \tau(p_\theta(x))) d\nu(x), \quad (6)$$

$$= \int f''(\rho(p_\theta(x))) (\partial_i \rho(p_\theta(x))) (\partial_j \rho(p_\theta(x))) d\nu(x), \quad (7)$$

$$= \int (f^*)''(\tau(p_\theta(x))) (\partial_i \tau(p_\theta(x))) (\partial_j \tau(p_\theta(x))) d\nu(x), \quad (8)$$

$$= \int \frac{1}{\rho'(p_\theta(x))\tau'(p_\theta(x))} (\partial_i p_\theta(x)) (\partial_j p_\theta(x)) d\nu(x). \quad (9)$$

This last equation shows that there is a gauge function freedom $\Psi(u) := \frac{1}{\rho'(u)\tau'(u)}$ when calculating the FIM.

The metric tensor can be derived [4] from the (ρ, τ) -divergence:

$$D_{\rho, \tau}(p : q) = \int (f(\rho(p(x))) + f^*(\tau(q(x))) - \rho(p(x))\tau(q(x))) d\nu(x) \quad (10)$$

Initially created 19th September 2017 (last updated September 2, 2020).

References

- [1] O. Calin and C. Udrişte. *Geometric Modeling in Probability and Statistics*. Mathematics and Statistics. Springer International Publishing, 2014.
- [2] Jan Naudts and Jun Zhang. Rho–tau embedding and gauge freedom in information geometry. *Information geometry*, 1(1):79–115, 2018.
- [3] Frank Nielsen. Cramér-Rao lower bound and information geometry. *arXiv preprint arXiv:1301.3578*, 2013.
- [4] Frank Nielsen. An elementary introduction to information geometry. *arXiv preprint arXiv:1808.08271*, 2018.
- [5] Frank Nielsen and Vincent Garcia. Statistical exponential families: A digest with flash cards. *arXiv preprint arXiv:0911.4863*, 2009.
- [6] Ke Sun and Frank Nielsen. Relative Fisher information and natural gradient for learning large modular models. In Doina Precup and Yee Whye Teh, editors, *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pages 3289–3298, International Convention Centre, Sydney, Australia, 06–11 Aug 2017. PMLR.