

Finance Data Science
Lecture 3: Covariance Matrix Estimation

Laurent El Ghaoui

MFE 230P Summer 2017
MFE Program
Haas School of Business
UC Berkeley

6/12/2017

Finance Data Science
3. Covariance Matrix
MFE 230P Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Outline

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage models

Sparse graphical models

Conditional independence

Penalized maximum-likelihood

Examples

References

Finance Data Science
3. Covariance Matrix
MFE 230P Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Motivations and goals

Covariance matrices are widely used in finance:

- Exploratory data analysis (see lecture 4).
- Risk analysis.
- Portfolio optimization.

In practice the number of data points n may be less than the number of dimensions p (assets).

This lecture: examine three estimation methods: one naïve (sample estimate), the other classical (factor model), the last modern.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Eigenvalue decomposition for symmetric matrices

Theorem (EVD of symmetric matrices)
We can decompose any symmetric $p \times p$ matrix S as

$$S = U \Lambda U^T = \sum_{i=1}^p \lambda_i u_i u_i^T,$$

where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$, with $\lambda_1 \geq \dots \geq \lambda_p$ the eigenvalues, and $U = [u_1, \dots, u_p]$ is a $p \times p$ orthogonal matrix ($U^T U = I_p$) that contains the eigenvectors u_i of S , that is:

$$S u_i = \lambda_i u_i, \quad i = 1, \dots, p.$$

Corollary: If S is square, symmetric:

$$\lambda_{\max}(S) = \max_{x : \|x\|_2=1} x^T S x. \tag{1}$$

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Positive semi-definite (PSD) matrices

A (square) symmetric matrix S is said to be *positive semi-definite* (PSD) if

$$\forall x, \quad x^T S x \geq 0.$$

In this case, we write $S \succeq 0$.

From EVD theorem: for any square, symmetric matrix S :

$$S \succeq 0 \iff \text{every eigenvalue of } S \text{ is non-negative.}$$

Hence we can numerically (via EVD) check positive semi-definiteness.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

The sample covariance matrix

Motivation

We can easily define the variance of a collection of numbers

z_1, \dots, z_m :

$$\sigma^2 = \frac{1}{m} \sum_{i=1}^m (z_i - \hat{z})^2,$$

where $\hat{z} = (1/m)(z_1 + \dots + z_m)$ is the average of the z_i 's.

- How can we extend this notion to higher dimensions (with z_i 's as vectors)?
- Why would we want to do that?

Note: for technical reasons the factor $1/m$ is often replaced with $1/(m-1)$, with little effect when m is large.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

The sample covariance matrix

Definition

Given a $p \times m$ data matrix $A = [a_1, \dots, a_m]$ (each row representing say a log-return time-series over m time periods), the *sample covariance matrix* is defined as the $p \times p$ matrix

$$S = \frac{1}{m} \sum_{i=1}^m (a_i - \hat{a})(a_i - \hat{a})^T, \quad \hat{a} := \frac{1}{m} \sum_{i=1}^m a_i.$$

We can express S as

$$S = \frac{1}{m} A_c A_c^T,$$

where A_c is the *centered data matrix* :

$$A_c = \begin{pmatrix} a_1 - \hat{a} & \dots & a_m - \hat{a} \end{pmatrix}$$

Note: for technical reasons the factor $1/m$ is often replaced with $1/(m - 1)$, with little effect when m is large.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

The sample covariance matrix

Link with directional variance

The (sample) variance along direction x is

$$\text{var}(x) = \frac{1}{m} \sum_{i=1}^m [x^T (a_i - \hat{a})]^2 = x^T S x = \frac{1}{m} \|A_c x\|_2^2.$$

where A_c is the centered data matrix:

Hence:

- ▶ the covariance matrix gives information about variance along **any** direction, via the quadratic function $x \rightarrow x^T S x$;
- ▶ the covariance matrix is always symmetric ($S = S^T$);
- ▶ It is also positive-semidefinite (PSD), since $x^T S x = \text{var}(x) \geq 0$ for every x .

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Application: portfolio risk

- **Data:** Consider n assets with returns over one period (e.g., day) $r_i, i = 1, \dots, n$. In general not known in advance.
- **Portfolio:** described by a vector $x \in \mathbf{R}^n$, with $x_i \geq 0$ the proportion of a total wealth invested in asset i .
- **Portfolio return:** $r^T x$; in general not known.
- **Expected return:** mean value of portfolio return, given by

$$\mathbf{E} r^T x = \hat{r}^T x,$$

- with $\hat{r} = (\hat{r}_1, \dots, \hat{r}_n)$ the vector of mean returns.
- **Portfolio risk:** Assuming return vector r is random, with mean \hat{r} and covariance matrix S , the variance of the portfolio is

$$\sigma^2(x) := \mathbf{E}_r(r^T x - \hat{r}^T x)^2 = x^T S x.$$

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

The sample covariance matrix

Total variance

For a given sample covariance matrix, we define the **total variance** to be the sum of the variances along the unit vectors

$$e_i = (0, \dots, 1, \dots, 0) \text{ (with 1 in } i\text{-th position, 0 otherwise).}$$

Total variance writes:

$$\sum_{i=1}^p \mathbf{var}(e_i) = \sum_{i=1}^p e_i^T S e_i = \sum_{i=1}^p S_{ii} := \mathbf{Tr} S,$$

where the symbol **Tr** (trace) denotes the sum of the diagonal elements of its matrix argument.


Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

What is wrong with the sample covariance?

Assume we draw random data with zero mean and true covariance $S = I_p$, and look at eigenvalues of the sample estimate, when both p, n are large.



- ▶ Eigenvalues should be all close to 1!
- ▶ This becomes true only when p is fixed and number of samples $n \rightarrow +\infty$.
- ▶ Red curve shows theoretical result from "random matrix theory" [2], which works for "large p , large n " case (see later).

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Estimation problem

In practice, the sample estimate might not work well in high dimensions; so we need to look for better estimates.

Problem: Given data points $x_1, \dots, x_n \in \mathbf{R}^d$, find an estimate of the covariance \hat{C} .

- ▶ Many methods start with the sample estimate ...
- ▶ ... and remove "noise" from it.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Measuring estimation quality

- Cross-validation principle:
- ▶ Remove 10 % of data points.
 - ▶ Record new estimate.
 - ▶ Measure average "error" between estimates.

How do we measure errors? We need a concept of distance between matrices:

- ▶ Frobenius norm (square-root of sum of squares of entries).
- ▶ If using a generative model (e.g., Gaussian), we can use Kullback-Leibler divergence (not quite a distance).

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Gaussian assumption

Let us assume that the data points are zero-mean, and follow a multi-variate Gaussian distribution: $x \simeq \mathcal{N}(0, \Sigma)$, with Σ a $p \times p$ covariance matrix. Assume Σ is positive definite.

The Gaussian probability density function is

$$p(\Sigma, x) := \frac{1}{(2\pi \det \Sigma)^{p/2}} \exp((1/2)x^T \Sigma^{-1} x).$$

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Maximum-likelihood

How can we find an estimate $\hat{\Sigma}$ of the true Σ , based on data points x_1, \dots, x_n ?

Maximum-likelihood principle: maximize the likelihood

$$L(\Sigma) := \prod_{i=1}^n p(\Sigma, x_i)$$

over the variable Σ .

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Solution

Changing variables ($X := \Sigma^{-1}$), and taking the log of the likelihood, the problem can be written as

$$\max_X \log \det X - \text{Tr } \hat{C} X$$

where \hat{C} is the sample covariance matrix. In this form, the maximum-likelihood problem is **convex**.

Solution: $X = \hat{C}^{-1}$, where \hat{C} is the sample covariance matrix!

Caveat: approach fails when \hat{C} is not positive-definite (e.g., when $p > n$).

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Issues

- What is wrong with the sample (i.e., ML) estimate?
- ▶ Fails in (interesting) case when $p > n$.
 - ▶ Does not handle missing data.
 - ▶ High sensitivity to outliers.
 - ▶ Can come up with better estimates (see next).
 - ▶ Gaussian assumption is not very good with finance data.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Data generative model

$y = Lz + \sigma e$

- ▶ y is the observation (data points).
- ▶ e is a noise vector (assume $E e = 0, E e e^T = \sigma^2 I$).
- ▶ z contains “factors” (assume $E e = 0, E e e^T = I$).
- ▶ L is a $p \times k$ loading matrix (usually, $k \ll p$).

This corresponds to a covariance matrix $\Sigma = \sigma^2 I + LL^T$.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Fitting factor models

Given sample covariance matrix $\hat{C} \succeq 0$, we can find L and α by solving

$$\min_{\alpha \succeq 0, L} \|\hat{C} - \alpha I - LL^T\|_F.$$

Solution: via EVD of \hat{C} .

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Scaled version

In practice, we may assume that each random variable has its own noise variance.

Modified problem:

$$\min_{D, L} \|\hat{C} - D - LL^T\|_F : D \text{ diagonal}, D \succeq 0.$$

This time, no obvious solution . . .

Can alternate optimization over D (easy) and L (EVD). Results in **local** optimum.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Computational benefits of factor models

A simple portfolio optimization problem

Risk-return trade-off:

min_x f(x) := x^T C x - lambda r^T x

- ▶ r in R^p (estimate) of returns.
- ▶ x in R^p portfolio vector (shorting allowed).
- ▶ C (estimate of) covariance matrix.
- ▶ Parameter lambda > 0 allows to choose trade-off.

The above problem is convex .

Assuming C > 0, optimal point found via grad f(x) = 0:

x* = lambda C^-1 r.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Computational benefits of factor models

Direct approach

Assume C = D + LL^T, with D > 0, diagonal, and F in R^{p x k}, with k << p: we need to solve

x* = (D + LL^T)^-1 y

with y := lambda r.

Direct approach: solve the p x p linear system

(D + LL^T)x = y,

without further exploiting structure. Cost: O(p^3).

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Computational benefits of factor models

Exploiting structure

Define $z := L^T x$, and rewrite $(D + LL^T)x = y$ as

$$\begin{pmatrix} D & L \\ L^T & -I_k \end{pmatrix} \begin{pmatrix} x \\ z \end{pmatrix} = \begin{pmatrix} y \\ 0 \end{pmatrix}$$

- ▶ Eliminate $x = D^{-1}(y - L^T z)$ and get teh $k \times k$ system in z :

$$(I + L^T D^{-1} L)z = D^{-1} Ly.$$

- ▶ Then solve for x via $Dx = (y - L^T z)$.

Cost: linear in p !

- ▶ Invert diagonal D : $O(p)$.
- ▶ Form $I + L^T D^{-1} L$ and solve for z : $O(k^3 + pk^2)$.
- ▶ Get x from z : $O(p)$.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized
maximum likelihood
Examples
References

Notes

The need for shrinkage

- ▶ Maximum-likelihood approach fails when $\hat{C} \not\succeq 0$ (e.g., $p > n$).
- ▶ Well-conditioned estimate is often needed for subsequent use (e.g., portfolio optimization).
(Condition number of \hat{C} is $\lambda_{\max}(\hat{C})/\lambda_{\min}(\hat{C})$.)

Basic idea: Modify \hat{C} by adding a diagonal, positive-definite term.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized
maximum likelihood
Examples
References

Notes

Ledoit and Wolf's model [6]

Estimate computed as a convex combination:

$$\hat{\Sigma} = \lambda I + (1 - \lambda)\hat{C},$$

where $\lambda \in (0, 1)$ is a *shrinkage* factor.

- ▶ A formula for λ is provided in [7] (has some nice statistical properties).
- ▶ Alternatively, choose λ based on cross-validation.
- ▶ Can replace the identity with another positive-definite matrix (allows to mix heterogeneous views on markets, such as news-based and price-based).
- ▶ Authors show improvements in the context of portfolio optimization.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Motivation

Assume we are given prices corresponding to many assets. We'd like to draw a graph that describes the links between the prices.

- ▶ Edges in the graph should exist when some strong, natural metric of similarity exist between assets.
- ▶ For better interpretability, a *sparse* graph is desirable.
- ▶ Various motivations: portfolio optimization (with sparse risk term), clustering, etc.

Here we focus on exploring *conditional independence* within nodes.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Gaussian assumption

Let us assume that the data points are zero-mean, and follow a multi-variate Gaussian distribution: $x \simeq \mathcal{N}(0, \Sigma)$, with Σ a $p \times p$ covariance matrix. Assume Σ is positive definite.

Gaussian probability density function:

$$p(x) = \frac{1}{(2\pi \det \Sigma)^{p/2}} \exp(-(1/2)x^T \Sigma^{-1} x).$$

where $X := \Sigma^{-1}$ is the *precision* matrix.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Conditional independence

The pair of random variables x_i, x_j are *conditionally independent* if, for x_k fixed ($k \neq i, j$), the density can be factored:

$$p(x) = p_i(x_i)p_j(x_j)$$

where p_i, p_j depend also on the other variables.

Interpretation: if all the other variables are fixed then x_i, x_j are independent. *Example:* Gray hair and shoe size are independent, conditioned on age.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Conditional independence
C.I. and the precision matrix

Theorem (C.I. for Gaussian RVs)
The variables x_i, x_j are conditionally independent if and only if the i, j element of the precision matrix is zero:
 $(\Sigma^{-1})_{ij} = 0$.

Proof.
The coefficient of $x_i x_j$ in $\log p(x)$ is $(\Sigma^{-1})_{ij}$. ■

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

Sparse precision matrix estimation

Let us encourage sparsity of the precision matrix in the maximum-likelihood problem:

$$\max_X \log \det X - \text{Tr } \hat{C}X - \lambda \|X\|_1,$$

with $\|X\|_1 := \sum_{i,j} |X_{ij}|$, and $\lambda > 0$ a parameter.

- ▶ The above provides an invertible result, even if \hat{C} is not positive-definite.
- ▶ The problem is convex.
- ▶ The result allows to discover a sparse graph revealing conditional independencies: look pairs (i, j) for which $X_{ij} = 0$.
- ▶ Motivations for the use of the l_1 -norm: encourages sparsity.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer
2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood


Examples

References


Notes

Example

Data: Interest rates



Using covariance matrix ($\lambda = 0$).



Using $\lambda = 0.1$.

The original precision matrix is dense, but the sparse version reveals the maturity structure (an information that was not given to the algorithm).


Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Example

Data: US Senate voting, 2002-2004



Again the sparse version reveals information, here political blocks within each party.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Code

- Python: <http://scikit-learn.org/stable/modules/covariance.html>
Implements a few methods for covariance estimation, including the sparse inverse covariance estimator.
- R: <http://strimmerlab.org/software/corpcor/>
Focuses on a special type of shrinkage estimator (James-Stein)

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence


Penalized maximum likelihood


Examples


References


Notes


References I


**N. El Karoui.**
Spectrum estimation for large dimensional covariance matrices using random matrix theory.
The Annals of Statistics, 36(6):2767–2790, 2008.


**N. El Karoui.**
High-dimensionality effects in the markowitz problem and other quadratic programs with linear constraints: Risk underestimation.
The Annals of Statistics, 38(6):3487–3566, 2010.


**Jiangqiang Fan, Yuan Liao, and Han Liu.**
An overview of the estimation of large covariance and precision matrices.
The Econometrics Journal, 18(1):C1–C32, 2015.
Online version at <https://arxiv.org/pdf/1504.02995.pdf>.

**Jürgen Franke, Wolfgang K. Härdle, and Christian M. Hafner.**
Statistics of Financial Markets: An Introduction.
2008.

**T. L. Lai and H. Xing.**
Statistical models and methods for financial markets.
Springer, 2008.

**Olivier Ledoit and Michael Wolf.**
Honey, i shrunk the sample covariance matrix.
The Journal of Portfolio Management, 30(4):110–119, 2004.

**Olivier Ledoit and Michael Wolf.**
A well-conditioned estimator for large-dimensional covariance matrices.
Journal of Multivariate Analysis, 88:365–411, February 2004.

**O.Banerjee, L. El Ghaoui, and A. d’Aspremont.**
Model selection through sparse maximum likelihood estimation for multivariate gaussian or binary data.
Journal of Machine Learning Research, 9:465–516, March 2008.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017

Introduction

Motivations

Recap: Eigenvalues

Covariance Matrices

Empirical covariance

Directional variance

Total variance

Estimation problem

Gaussian Models

Maximum likelihood

Issues

Regularization

Factor models

Shrinkage

Graphical models

Conditional independence

Penalized maximum likelihood

Examples

References

Notes

References II



Adam J Rothman, Elizaveta Levina, and Ji Zhu.
A new approach to cholesky-based covariance regularization in high dimensions.
Biometrika, pages 539–550, 2010.

Finance Data Science
3. Covariance Matrix
MFE 230P, Summer 2017
Introduction
Motivations
Recap: Eigenvalues
Covariance Matrices
Empirical covariance
Directional variance
Total variance
Estimation problem
Gaussian Models
Maximum likelihood
Issues
Regularization
Factor models
Shrinkage
Graphical models
Conditional independence
Penalized maximum likelihood
Examples
References

Notes

Notes
