

T3 group factors: EFT_2_after_S1_S2

Active theory: SMEFT with the Weinberg operator C_5 . No T3 heavy field remains dynamical.

$$j = \frac{d-1}{2}, \qquad C_2(d) = \frac{d^2-1}{4}, \qquad T(d) = \frac{d(d^2-1)}{12}.$$

In every comparison table, $\mathcal{S}$ denotes the RGE structure multiplying the displayed group factor in $16\pi^2\beta$.

Notation key

symbol	interaction / definition
g_1	$D_\mu \supset ig_1 Y B_\mu$
g_2	$D_\mu \supset ig_2 T^A W_\mu^A$
g_3	$D_\mu \supset ig_3 t^A G_\mu^A$
Y_u	$-\bar{Q} Y_u \widetilde{H} u + \text{h.c.}$
Y_d	$-\bar{Q} Y_d H d + \text{h.c.}$
Y_e	$-\bar{L} Y_e H e + \text{h.c.}$
T	$\text{Tr}\left(Y_e^\dagger Y_e + 3Y_u^\dagger Y_u + 3Y_d^\dagger Y_d\right)$
λ_1	$\frac{1}{2}\lambda_1(H^\dagger H)^2$
C_5^{ij}	$(L_i^T C L_j) H H$

Stage field content

run	S_1	S_2	F	integrated at this stage	active after matching
T3-A-m4	$(1,-2)$	$(3,-1)$	$(2,-\frac{3}{2})$	S1, S2	none
T3-A-m2	$(1,-1)$	$(3,0)$	$(2,-\frac{1}{2})$	S1, S2	none
T3-A-p0	$(1,0)$	$(3,1)$	$(2,\frac{1}{2})$	S1, S2	none
T3-B-m3	$(2,-\frac{3}{2})$	$(2,-\frac{1}{2})$	$(1,-1)$	S1, S2	none
T3-B-m1	$(2,-\frac{1}{2})$	$(2,\frac{1}{2})$	$(1,0)$	S1, S2	none
T3-B-p1	$(2,\frac{1}{2})$	$(2,\frac{3}{2})$	$(1,1)$	S1, S2	none
T3-C-m3	$(2,-\frac{3}{2})$	$(2,-\frac{1}{2})$	$(3,-1)$	S1, S2	none
T3-C-m1	$(2,-\frac{1}{2})$	$(2,\frac{1}{2})$	$(3,0)$	S1, S2	none
T3-C-p1	$(2,\frac{1}{2})$	$(2,\frac{3}{2})$	$(3,1)$	S1, S2	none
T3-D-m2	$(3,-1)$	$(1,0)$	$(2,-\frac{1}{2})$	S1, S2	none
T3-D-p0	$(3,0)$	$(1,1)$	$(2,\frac{1}{2})$	S1, S2	none
T3-D-p2	$(3,1)$	$(1,2)$	$(2,\frac{3}{2})$	S1, S2	none

T3-E-m4	$(3, -2)$	$(3, -1)$	$(2, -\frac{3}{2})$	S1, S2	none
T3-E-m2	$(3, -1)$	$(3, 0)$	$(2, -\frac{1}{2})$	S1, S2	none
T3-E-p0	$(3, 0)$	$(3, 1)$	$(2, \frac{1}{2})$	S1, S2	none
T3-E-p2	$(3, 1)$	$(3, 2)$	$(2, \frac{3}{2})$	S1, S2	none

Decoupling statement

All dependence on the original S_1 , S_2 and F representations is contained in the matched boundary value $C_5(\mu_{\text{th}})$. The one-loop running below the final threshold is common to all T3 classes.

β_{C_5}

Run block 1: T3-A-m4–T3-B-m1

$\mathcal{S}$	T3 – A – m4 ($\alpha = -4$)	T3 – A – m2 ($\alpha = -2$)	T3 – A – p0 ($\alpha = 0$)	T3 – B – m3 ($\alpha = -3$)	T3 – B – m1 ($\alpha = -1$)
$\lambda_1 C_5$	2	2	2	2	2
$g_2^2 C_5$	−3	−3	−3	−3	−3
$T C_5$	2	2	2	2	2
$(Y_e^\dagger Y_e)^T C_5$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$
$C_5(Y_e^\dagger Y_e)$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$

Run block 2: T3-B-p1–T3-D-m2

$\mathcal{S}$	T3 – B – p1 ($\alpha = 1$)	T3 – C – m3 ($\alpha = -3$)	T3 – C – m1 ($\alpha = -1$)	T3 – C – p1 ($\alpha = 1$)	T3 – D – m2 ($\alpha = -2$)
$\lambda_1 C_5$	2	2	2	2	2
$g_2^2 C_5$	−3	−3	−3	−3	−3
$T C_5$	2	2	2	2	2
$(Y_e^\dagger Y_e)^T C_5$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$
$C_5(Y_e^\dagger Y_e)$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$

Run block 3: T3-D-p0–T3-E-p0

$\mathcal{S}$	T3 – D – p0 ($\alpha = 0$)	T3 – D – p2 ($\alpha = 2$)	T3 – E – m4 ($\alpha = -4$)	T3 – E – m2 ($\alpha = -2$)	T3 – E – p0 ($\alpha = 0$)
$\lambda_1 C_5$	2	2	2	2	2
$g_2^2 C_5$	−3	−3	−3	−3	−3
$T C_5$	2	2	2	2	2
$(Y_e^\dagger Y_e)^T C_5$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$

$C_5(Y_e^\dagger Y_e)$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$	$-\frac{3}{2}$
------------------------	----------------	----------------	----------------	----------------	----------------

Run block 4: T3-E-p2-T3-E-p2

$\mathcal{S}$	T3 – E – p2 ($\alpha = 2$)
$\lambda_1\,C_5$	2
$g_2^2\,C_5$	−3
$T\,C_5$	2
$(Y_e^\dagger Y_e)^T C_5$	$-\frac{3}{2}$
$C_5(Y_e^\dagger Y_e)$	$-\frac{3}{2}$

Flavor form

$$16\pi^2\frac{dC_5}{d\ln\mu}=(2\lambda_1-3g_2^2+2T)C_5\tag{1}$$

$$-\frac{3}{2}\left[(Y_e^\dagger Y_e)^TC_5+C_5(Y_e^\dagger Y_e)\right].\tag{2}$$