

Weinberg-operator coefficient

$$T3 - A, \quad \alpha = 0: \quad S_1 = (1, 0), \quad S_2 = (3, 1), \quad F = (2, \frac{1}{2})$$

$$C_5 = \frac{1}{2} \left(-\frac{\hbar M_F^3 M_{S_2}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S_2}^2}\right) \overline{y_{1,i_1,r_1} y_{2,i_2,r_1}}}{(M_F^2 - M_{S_1}^2)(M_F^2 - M_{S_2}^2)(M_{S_1}^2 - M_{S_2}^2)} + \frac{\hbar M_F^3 M_{S_1}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S_1}^2}\right) \overline{y_{1,i_1,r_1} y_{2,i_2,r_1}}}{(M_F^2 - M_{S_1}^2)(M_F^2 - M_{S_2}^2)(M_{S_1}^2 - M_{S_2}^2)} - \frac{\hbar M_F M_{S_1}^2 M_{S_2}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S_1}^2}\right) \overline{y_{1,i_1,r_1} y_{2,i_2,r_1}}}{(M_F^2 - M_{S_1}^2)(M_F^2 - M_{S_2}^2)(M_{S_1}^2 - M_{S_2}^2)} + \frac{\hbar M_F M_{S_1}^2 M_{S_2}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S_2}^2}\right) \overline{y_{1,i_1,r_1} y_{2,i_2,r_1}}}{(M_F^2 - M_{S_1}^2)(M_F^2 - M_{S_2}^2)(M_{S_1}^2 - M_{S_2}^2)} \right)$$