

Weinberg-operator coefficient

$$T3 - d4 - d6 - F5, \alpha = 0: \quad S_1 = (4, 0), \quad S_2 = (6, 1), \quad F = (5, \frac{1}{2})$$

$$C_5 = \frac{4}{3} \left(-\frac{\sqrt{6}hM_F^3M_{S2}^2\lambda_{T3}\log\left(\frac{M_F^2}{M_{S2}^2}\right)\overline{y_{1,i_1,r_1}y_{2,i_2,r_1}}}{(M_F^2-M_{S1}^2)(M_F^2-M_{S2}^2)(M_{S1}^2-M_{S2}^2)} + \frac{\sqrt{6}hM_F^3M_{S1}^2\lambda_{T3}\log\left(\frac{M_F^2}{M_{S1}^2}\right)\overline{y_{1,i_1,r_1}y_{2,i_2,r_1}}}{(M_F^2-M_{S1}^2)(M_F^2-M_{S2}^2)(M_{S1}^2-M_{S2}^2)} - \frac{\sqrt{6}hM_F M_{S1}^2 M_{S2}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S1}^2}\right)\overline{y_{1,i_1,r_1}y_{2,i_2,r_1}}}{(M_F^2-M_{S1}^2)(M_F^2-M_{S2}^2)(M_{S1}^2-M_{S2}^2)} + \frac{\sqrt{6}hM_F M_{S1}^2 M_{S2}^2 \lambda_{T3} \log\left(\frac{M_F^2}{M_{S2}^2}\right)\overline{y_{1,i_1,r_1}y_{2,i_2,r_1}}}{(M_F^2-M_{S1}^2)(M_F^2-M_{S2}^2)(M_{S1}^2-M_{S2}^2)} \right)$$