T3 group factors: EFT_1_after_F_S1_S2

Active theory: SMEFT with the Weinberg operator C_5. No T3 heavy field remains dynamical.

$$j = \frac{d-1}{2}, \qquad C_2(d) = \frac{d^2-1}{4}, \qquad T(d) = \frac{d(d^2-1)}{12}.$$

In every comparison table, $\mathcal{S}$ denotes the RGE structure multiplying the displayed group factor in $16\pi^2\beta$.

Notation key

symbol	interaction / definition
g_1	$D_\mu \supset i g_1 Y B_\mu$
g_2	$D_\mu \supset i g_2 T^A W_\mu^A$
g_3	$D_\mu \supset i g_3 t^A G_\mu^A$
Y_u	$-\bar{Q}\, Y_u\, \widetilde{H}\, u + \text{h.c.}$
Y_d	$-\bar{Q}\, Y_d\, H\, d + \text{h.c.}$
Y_e	$-\bar{L}\, Y_e\, H\, e + \text{h.c.}$
T	$\mathrm{Tr}\left(Y_e^\dagger Y_e + 3Y_u^\dagger Y_u + 3Y_d^\dagger Y_d\right)$
λ_1	$\frac{1}{2}\lambda_1 (H^\dagger H)^2$
C_5^{ij}	$(L_i^T C L_j)\, HH$

Stage field content

run	S_1	S_2	F	integrated at this stage	active after matching
T3-E	$(3,1)$	$(3,2)$	$(2,\frac{3}{2})$	F, S1, S2	none

Decoupling statement

All dependence on the original S_1, S_2 and F representations is contained in the matched boundary value $C_5(\mu_{\rm th})$. The one-loop running below the final threshold is common to all T3 classes.

β_{C_5}

$\mathcal{S}$	T3 − E ($\alpha = 2$)
$\lambda_1\, C_5$	2
$g_2^2\, C_5$	-3
$T\, C_5$	2
$(Y_e^\dagger Y_e)^T C_5$	$-\frac{3}{2}$
$C_5 (Y_e^\dagger Y_e)$	$-\frac{3}{2}$

Flavor form

$$16\pi^2 \frac{dC_5}{d\ln\mu} = (2\lambda_1 - 3g_2^2 + 2T) C_5 \tag{1}$$

$$- \frac{3}{2}\left[(Y_e^\dagger Y_e)^T C_5 + C_5 (Y_e^\dagger Y_e)\right]. \tag{2}$$