

T3 Weinberg-operator coefficient comparison

Model	Ordered coefficient extracted from Matchete	Symmetric physical Wilson coefficient
T3-B	$A_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} \overline{y_{1,i,r} y_{2,j,r}}$	$C_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} (\overline{y_{1,i,r} y_{2,j,r}} + \overline{y_{1,j,r} y_{2,i,r}})$
T3-C	$A_5^{ij} = \frac{2}{3} \hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2) \overline{y_{1,i,r} y_{2,j,r}}$	$C_5^{ij} = \frac{2}{3} \hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2) (\overline{y_{1,i,r} y_{2,j,r}} + \overline{y_{1,j,r} y_{2,i,r}})$
T3-A	$A_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} \overline{y_{1,i,r} y_{2,j,r}}$	$C_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} (\overline{y_{1,i,r} y_{2,j,r}} + \overline{y_{1,j,r} y_{2,i,r}})$
T3-D	$A_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} \overline{y_{1,i,r} y_{2,j,r}}$	$C_5^{ij} = -\frac{2\hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2)}{\sqrt{3}} (\overline{y_{1,i,r} y_{2,j,r}} + \overline{y_{1,j,r} y_{2,i,r}})$
T3-E	$A_5^{ij} = 2\sqrt{\frac{2}{3}} \hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2) \overline{y_{1,i,r} y_{2,j,r}}$	$C_5^{ij} = 2\sqrt{\frac{2}{3}} \hbar M_F \lambda_{T3} I_3 (M_F^2, M_{S1}^2, M_{S2}^2) (\overline{y_{1,i,r} y_{2,j,r}} + \overline{y_{1,j,r} y_{2,i,r}})$