

Towards Braided ∞ -Operads

by

Franziska Ahrens

born

03.07.1997

A thesis submitted to the Department of Mathematics

Universität Hamburg

In Partial Fulfillment of the Requirements
for the Degree of Master of Science

July 14, 2025

Supervisor: Prof. Dr. Tobias Dyckerhoff

Examiner: Prof. Dr. Christoph Schweigert

Contents

1	Introduction	3
2	Operads	3
	2.1 Planar and Symmetric Operads	4
	2.2 Colored Operads	12
3	Braided Operads	23
	3.1 Action Operad	24
	3.2 Universal Covering of E_2	26
4	Colored Braided Operads	31
	4.1 Braided Monoidal Structure	35
	4.2 Colored Braided Operads as Op-fibrations	42
5	Outlook	42
6	Appendix	42

1. Introduction

1 Introduction

Coming soon

2 Operads

Operads are a concept that originally occurred when studying loop spaces. Let ΩX be the space of based loops in a topological space X . Its structure reflects an interesting form of algebraic composition. The composition of loops is only associative up to homotopy equivalence meaning that loop spaces behave like a monoid in a homotopy-theoretic sense, but not in a strict algebraic sense. An operad can be thought of as a structure that encodes a family of operations, each of which has a certain number of inputs, and a way to compose these operations together. The composition of operations in an operad satisfies associativity and unit laws.

Let $\alpha: [0, 1] \rightarrow X$ and $\beta: [0, 1] \rightarrow X$ be loops in X having the same base point x_0 . Their composition $\beta \circ \alpha$ is usually defined by splitting the unit interval in half, i.e.

$$\beta \circ \alpha(s) := \begin{cases} \alpha(2s) & \text{for } s \in [0, \frac{1}{2}] \\ \beta(2s - 1) & \text{for } s \in (\frac{1}{2}, 1]. \end{cases}$$

This concatenation in itself is not associative; consider $\alpha, \beta, \gamma \in \Omega X$. Then by the given definition $\gamma \circ (\beta \circ \alpha)$ and $(\gamma \circ \beta) \circ \alpha$ are different maps:

$$\gamma \circ (\beta \circ \alpha)(s) = \begin{cases} \alpha(4s) & \text{for } s \in [0, \frac{1}{4}] \\ \beta(4s - 1) & \text{for } s \in (\frac{1}{4}, \frac{1}{2}] \\ \gamma(2s - 1) & \text{for } s \in (\frac{1}{2}, 1] \end{cases}$$

$$(\gamma \circ \beta) \circ \alpha(s) = \begin{cases} \alpha(2s) & \text{for } s \in [0, \frac{1}{2}] \\ \beta(4s - 2) & \text{for } s \in (\frac{1}{2}, \frac{3}{4}] \\ \gamma(4s - 3) & \text{for } s \in (\frac{3}{4}, 1] \end{cases}$$

The choice of splitting the unit interval in half is merely a convention. Hence, composition of loops can be thought of as splitting intervals in an arbitrary manner; or in other words embedding unit intervals into a unit interval such that the image of the inner intervals are disjoint.

$$[[I_1] \dots [I_i] \dots [I_n]]$$

So, for an ordered family of loops $\{\alpha_i: I_i \rightarrow X\}_{1 \leq i \leq n}$ there is for each order preserving concatenation β a unique map $f \in \text{Emb}(I_1, \dots, I_n, I)$ such that $\beta \circ f(I_i) = \alpha_i$ for all $1 \leq i \leq n$. We denote the set of embeddings by $E(n) = \text{Emb}(I_1, \dots, I_n, I)$ for each $n \in \mathbb{N}$. For every $(f_k, (f_{j_i})_{1 \leq i \leq k}) \in E(k) \times E(j_1) \times \dots \times E(j_k)$ we obtain an embedding $f \in E(j_1 + \dots + j_k)$ by defining $f = f_k \circ (f_{j_1} \sqcup \dots \sqcup f_{j_k})$. This defines a composition map

$$\gamma: E(k) \times E(j_1) \times \dots \times E(j_k) \rightarrow E(j_1 + \dots + j_k)$$

Since the composition of maps is associative the following diagram commutes:

$$\begin{array}{ccc} E(k) \otimes (\prod_{s=1}^k E(j_s)) \times (\prod_{r=1}^j E(i_r)) & \xrightarrow{\gamma \otimes \text{Id}} & E(j) \times (\prod_{r=1}^j E(i_r)) \\ \downarrow s & & \downarrow \gamma \\ & & E(i) \\ & & \uparrow \gamma \\ E(k) \otimes (\prod_{s=1}^k (E(j_s) \times \prod_{q=1}^{j_s} E(i_{g_{s-1}+q}))) & \xrightarrow{\text{Id} \times \gamma^s} & E(k) \times (\prod_{s=1}^k E(h_s)) \end{array}$$

2. Operads

where s shuffles the cartesian product and

$$n, j_1, \dots, j_k, i_1, \dots, i_j \in \mathbb{N}$$

with

$$\sum_{1 \leq s \leq k} j_s = j, \sum_{1 \leq r \leq j} i_r = i, g_s = \sum_{0 \leq l \leq s} j_l \text{ and } h_s = \sum_{1 \leq q \leq j_s} i_{g_{s-1}+q}.$$

Further composing with the identity embedding in $E(1)$ does nothing. Thus, we have seen our first example of an operad, providing a framework in which the composition of loop spaces satisfies associativity.

In this way, operads provide a more general and flexible formalism for capturing the algebraic structures found in loop spaces and other topological constructs where composition occurs up to homotopy. Similarly to monoids that can be considered as single object categories, operads can be considered as single object multicategories. In this section, we will introduce the notion of a planar and a symmetric operad as they were introduced by May in [IM]. Then, we will move on to the more general notion of a colored operad following [JL]. We will establish a second equivalent definition of a colored operad that is purely categorical, meaning we can therefore also phrase it in an ∞ -categorical context. In the next section we will then consider braided operads and we work towards defining braided colored operads. Ultimately, our goal is a purely functorial definition of a braided colored operad such that we can lift the concept to an ∞ -categorical context as well.

2.1 Planar and Symmetric Operads

From now on, unless it is stated otherwise, let $\mathcal{C}$ be a symmetric monoidal category with tensor product $\otimes$ and tensor unit I . For an object C in $\mathcal{C}$, we write C^j for the j -th tensor power of C .

Definition 2.1. A *planar operad* $\mathcal{P}$ in $\mathcal{C}$ consists of a family of objects $(\mathcal{P}(j))_{j \in \mathbb{N}}$ together with a unit morphism $\eta: k \rightarrow \mathcal{P}(1)$ and a product morphism

$$\gamma: \mathcal{P}(k) \otimes \mathcal{P}(j_1) \otimes \dots \otimes \mathcal{P}(j_k) \rightarrow \mathcal{P}(j),$$

where $\sum_{1 \leq i \leq k} j_i = j$ such that the following associativity and unitality constraints hold:

- For any choice of integers

$$n, j_1, \dots, j_k, i_1, \dots, i_j$$

with

$$\sum_{1 \leq s \leq k} j_s = j, \sum_{1 \leq r \leq j} i_r = i, g_s = \sum_{0 \leq l \leq s} j_l \text{ and } h_s = \sum_{1 \leq q \leq j_s} i_{g_{s-1}+q}$$

the following diagram commutes:

$$\begin{array}{ccc} \mathcal{P}(k) \otimes (\bigotimes_{s=1}^k \mathcal{P}(j_s)) \otimes (\bigotimes_{r=1}^j \mathcal{P}(i_r)) & \xrightarrow{\gamma \otimes \text{Id}} & \mathcal{P}(j) \otimes (\bigotimes_{r=1}^j \mathcal{P}(i_r)) \\ \downarrow s & & \downarrow \gamma \\ & & \mathcal{P}(i) \\ & & \uparrow \gamma \\ \mathcal{P}(k) \otimes (\bigotimes_{s=1}^k (\mathcal{P}(j_s) \otimes \bigotimes_{q=1}^{j_s} \mathcal{P}(i_{g_{s-1}+q}))) & \xrightarrow{\text{Id} \otimes (\bigotimes_s \gamma)} & \mathcal{P}(k) \otimes ((\bigotimes_{s=1}^k \mathcal{P}(h_s))) \end{array}$$

where s shuffles the entries of the tensor product.

- For all $j \in \mathbb{N}$ the following diagrams commute:

2. Operads

$$\begin{array}{ccc}
 \mathcal{P}(j) \otimes k^j & \xrightarrow{\cong} & \mathcal{P}(j) \\
 \text{Id} \otimes (\otimes_j \eta) \downarrow & \nearrow \gamma & \\
 \mathcal{P}(j) \otimes \mathcal{P}(1)^j & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 k \otimes \mathcal{P}(j) & \xrightarrow{\cong} & \mathcal{P}(j) \\
 \eta \otimes \text{Id} \downarrow & \nearrow \gamma & \\
 \mathcal{P}(1) \otimes \mathcal{P}(j) & &
 \end{array}$$

Examples 2.2. Let us consider some examples of planar operads in a symmetric monoidal category $\mathcal{C}$:

1. Later we will define what it means to be an algebra over an operad. The *associative operad* $\text{Ass}^\otimes$ is the operad whose algebras are associative monoids. The associative planar operad in $\mathcal{C}$ is the operad $\text{Ass}^\otimes_{\mathcal{C}}(n) = \mathbf{I}$ for all $i \in \mathbb{N}$ with $\text{Id}_{\mathbf{I}}$ being the unit and the composition map being the canonical isomorphism. Hence, for $\mathcal{C} = \mathbf{Set}$, the category of sets equipped with the cartesian symmetric monoidal structure, we obtain exactly one n -ary operation for each n : $\text{Ass}^\otimes_{\mathbf{Set}}$ is defined by $\text{Ass}^\otimes_{\mathbf{Set}}(n) = *$ for all n . Similarly, for $\mathcal{C} = \mathbf{Vect}_K$ we get the free abelian vector space over $*$, i.e. K .
2. At the beginning of the section we already saw that the concatenation of loops defines an operad: Let $\mathcal{C} = \mathbf{Top}$. Then we can define the *operad of little interval* $\mathbf{E}$ as

$$\mathbf{E}(n) = \text{Emb}(I_1, \dots, I_n, I),$$

the set of orientation-preserving embeddings of n disjoint intervals into the unit interval I . We equip $\mathbf{E}(n)$ with the subspace topology of $\mathbb{R}^n$. Composition is defined by inserting the intervals into another interval as we have seen above.

3. For $\mathcal{C}$ being closed, i.e. having an internal hom functor $\underline{\text{Hom}}(-, -)$, we can define another planar operad End_X for any object $X \in \mathcal{C}$:

$$\text{End}_X(n) := \underline{\text{Hom}}(X^{\otimes n}, X)$$

The unit morphism η is given by the image of Id_X under the adjunction:

$$\text{Hom}(X, X) \cong \text{Hom}(I, \underline{\text{Hom}}(X, X))$$

Further, the composition morphism

$$\underline{\text{Hom}}(X^{\otimes k}, X) \otimes \bigotimes_s \underline{\text{Hom}}(X^{\otimes j_s}, X) \rightarrow \underline{\text{Hom}}(X^{\otimes j}, X)$$

is given by composition:

$$\begin{array}{c}
 \underline{\text{Hom}}(X^{\otimes k}, X) \otimes \bigotimes_s \underline{\text{Hom}}(X^{\otimes j_s}, X) \\
 \downarrow \\
 \underline{\text{Hom}}(X^{\otimes k}, X) \otimes \underline{\text{Hom}}(X^{\otimes j}, X^{\otimes k}) \\
 \downarrow \\
 \underline{\text{Hom}}(X^{\otimes j}, X)
 \end{array}$$

We call End_X the *endomorphism operad over X* .

For another example of an operad, which we will encounter quite a lot in the following, we need to recall the definition of the braid group first.

Definition 2.3. For each integer $n > 2$ we define the *braid group* on n strands as the group with $n - 1$ generators $\sigma_1, \dots, \sigma_{n-1}$ satisfying the following relations:

$$\begin{array}{ll}
 \sigma_i \sigma_j = \sigma_j \sigma_i & \text{for } |i - j| > 1 \\
 \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} & \text{for } 1 \leq i \leq n - 2
 \end{array}$$

For $n = 2$ we define B_2 as the free group with one generator and let $B_0 = B_1 = \{1\}$ be the trivial group.

2. Operads

Remark 2.4. The name braid group B_n makes sense when we consider the following depiction of the group for $0 \leq i, j \leq n$ with $i+1 < j$:

$$\begin{aligned}
 \sigma_i &= \begin{array}{c} \text{Diagram of a braid with } i \text{ strands crossing over } i+1 \end{array} \\
 \sigma_i \circ \sigma_j &= \begin{array}{c} \text{Diagram of two braid blocks: first } \sigma_i \text{ on strands } i, i+1, \dots, \text{ then } \sigma_j \text{ on strands } j, j+1, \dots \end{array} = \sigma_j \circ \sigma_i \\
 \sigma_i \circ \sigma_{i+1} \circ \sigma_i &= \begin{array}{c} \text{Diagram of three braid blocks: } \sigma_i \text{ on } i, i+1, \text{ then } \sigma_{i+1} \text{ on } i+1, i+2, \text{ then } \sigma_i \text{ on } i, i+1 \end{array} = \begin{array}{c} \text{Diagram of three braid blocks: } \sigma_{i+1} \text{ on } i+1, i+2, \text{ then } \sigma_i \text{ on } i, i+1, \text{ then } \sigma_{i+1} \text{ on } i+1, i+2 \end{array} = \sigma_{i+1} \circ \sigma_i \circ \sigma_{i+1}
 \end{aligned}$$

Further details on how this is defined formally can be found in [CS].

We state some facts about braid groups that are known and can for example be found in [DY].

Remark 2.5. 1. There is a projection, which we denote by $\pi: B_n \rightarrow S_n$, that adds the condition $\sigma_i^2 = \text{Id}$.

2. The direct sum homomorphism

$$\bigoplus_S: S_{k_1} \times \cdots \times S_{k_n} \rightarrow S_{k_1 + \cdots + k_n}$$

can be lifted to the braid groups:

$$\bigoplus_B: B_{k_1} \times \cdots \times B_{k_n} \rightarrow B_{k_1 + \cdots + k_n}$$

such that $\bigoplus_B \circ \pi = (\pi)^k \circ \bigoplus_S$. Geometrically, $b_1 \oplus \cdots \oplus b_n \in B_{k_1 + \cdots + k_n}$ describes the braid that places $b_1, \dots, b_n$ side by side from left to right.

3. For $n \geq 1$ and $k_1, \dots, k_n \in \mathbb{N}$, the block permutation map

$$(-)_S(k_1, \dots, k_n): S_n \rightarrow S_{k_1 + \cdots + k_n}$$

can be lifted to the braid groups:

$$(-)_B(k_1, \dots, k_n): B_n \rightarrow B_{k_1 + \cdots + k_n}$$

such that $(-)_B(k_1, \dots, k_n) \circ \pi = \pi \circ (-)_S(k_1, \dots, k_n)$. Geometrically, for $b \in B_n$, the corresponding block braid $b(k_1, \dots, k_n)$ is obtained by replacing the i -th strand by k_i many parallel strands for $1 \leq i \leq n$.

2. Operads

4. Let $\sigma, \rho \in B_n$ and $\tau_j \in B_{k_j}$ for $1 \leq j \leq n$. Then the following hold in $B_{k_1+\dots+k_n}$:

$$\begin{aligned} (\sigma\rho)(k_1, \dots, k_n) &= \sigma(k_{\bar{\rho}^{-1}(1)}, \dots, k_{\bar{\rho}^{-1}(n)}) \cdot \rho(k_1, \dots, k_n) \\ \sigma(k_1, \dots, k_n) \cdot (\tau_1 \oplus \dots \oplus \tau_n) &= (\tau_{\bar{\sigma}^{-1}(1)} \oplus \dots \oplus \tau_{\bar{\sigma}^{-1}(n)}) \cdot \sigma(k_1, \dots, k_n) \end{aligned}$$

where $\bar{\sigma} + \pi(\sigma)$ and $\bar{\rho} + \pi(\rho)$.

Example 2.6. Let $\mathcal{B}$ be the *braid operad* defined by:

$$\mathcal{B}(n) := B_n,$$

where B_n denotes the braid group on n strands. The composition morphism


$$\begin{aligned} \gamma: B_k \times B_{j_1} \times \dots \times B_{j_k} &\rightarrow B_j \\ (\tau, (\sigma_i)_{1 \leq i \leq k}) &\mapsto \tau(\sigma_1 \oplus \dots \oplus \sigma_k) \end{aligned}$$

where $\tau(\sigma_1 \oplus \dots \oplus \sigma_k)$ denotes the braid that τ -braids k many block braids $\sigma_1, \dots, \sigma_k$ as if they were strands in B_k .


So for the composition map

$$B_2 \times B_2 \times B_2 \rightarrow B_4$$

let us observe what happens with the following element in $B_2 \times B_2 \times B_2$:



After applying the composition map we obtain the following braiding:



The associativity and unitality conditions follow from the group conditions.

Definition 2.7. Let $\mathcal{P}, \mathcal{P}'$ be planar operads in a symmetric monoidal category $\mathcal{C}$. A *morphism of planar operads* $\beta: \mathcal{P} \rightarrow \mathcal{P}'$ consists of morphisms $\beta_n: \mathcal{P}(n) \rightarrow \mathcal{P}'(n)$ for all $n \in \mathbb{N}$ such that the following diagrams commute:

$$\begin{array}{ccc} \mathcal{P}(k) \otimes \mathcal{P}(j_1) \otimes \dots \otimes \mathcal{P}(j_k) & \xrightarrow{\beta_k \otimes \beta_{j_1} \otimes \dots \otimes \beta_{j_k}} & \mathcal{P}'(k) \otimes \mathcal{P}'(j_1) \otimes \dots \otimes \mathcal{P}'(j_k) \\ \gamma \downarrow & & \downarrow \gamma' \\ \mathcal{P}(j) & \xrightarrow{\beta_j} & \mathcal{P}'(j) \end{array}$$

$$\begin{array}{ccc} & k & \\ \eta \swarrow & & \searrow \eta' \\ \mathcal{P}(1) & \xrightarrow{\beta_1} & \mathcal{P}'(1) \end{array}$$

Therefore, for any closed symmetric monoidal category $\mathcal{C}$ the planar operads and their morphisms form a category $\mathbf{PlanarOp}_{\mathcal{C}}$.

When we later discuss operads, we will seldom focus on planar operads but will work with symmetric operads instead.

2. Operads

Definition 2.8. A (*symmetric*) *operad* $\mathcal{O}$ in $\mathcal{C}$ is a planar operad as defined in definition 2.1 with an action of S_n on $\mathcal{O}(n)$ together with the following equivariance condition: Let $\sigma(j_1, \dots, j_k) \in S_j$ denote the element that permutes k blocks of length j_l for $1 \leq l \leq k$. Then, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) & \xrightarrow{\sigma \otimes \sigma^{-1}} & \mathcal{O}(k) \otimes \mathcal{O}(j_{\sigma(1)}) \otimes \dots \otimes \mathcal{O}(j_{\sigma(k)}) \\ \gamma \downarrow & & \downarrow \gamma \\ \mathcal{O}(j) & \xrightarrow{\sigma(j_1 \dots j_k)} & \mathcal{O}(j). \end{array}$$

Furthermore, let $\tau_l \in S_{j_l}$ and $\tau_1 \oplus \dots \oplus \tau_k$ denotes the sum of permutations on k blocks of length j_l for $1 \leq l \leq k$. Then the following diagram commutes:

$$\begin{array}{ccc} \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) & \xrightarrow{\text{Id} \otimes \tau_1 \otimes \dots \otimes \tau_k} & \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) \\ \gamma \downarrow & & \downarrow \gamma \\ \mathcal{O}(j) & \xrightarrow{\tau_1 \oplus \dots \oplus \tau_k} & \mathcal{O}(j). \end{array}$$

As already mentioned we introduce operads as a tool to generally encode algebraic structure.

Definition 2.9. Let $\mathcal{O}$ be an operad in $\mathcal{C}$. An $\mathcal{O}$ -*algebra* is an object A in $\mathcal{C}$ with maps

$$\theta: \mathcal{O}(j) \otimes A^j \rightarrow A$$

for $j \in \mathbb{N}$ such that the following associativity, unitality and equivariant conditions hold:

- Associative:

$$\begin{array}{ccc} \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) \otimes A^j & \xrightarrow{\gamma \otimes \text{Id}} & \mathcal{O}(j) \otimes A^j \\ \downarrow s & & \downarrow \theta \\ \mathcal{O}(k) \otimes (\bigotimes (\mathcal{O}(j_l) \otimes A^{j_l})) & \xrightarrow{\text{Id} \otimes \theta^k} & \mathcal{O}(k) \otimes A^k \\ & & \uparrow \theta \\ & & A \end{array}$$

- Unital:

$$\begin{array}{ccc} \text{Id} \otimes A & \xrightarrow{\cong} & A \\ \eta \otimes \text{Id} \downarrow & \nearrow \theta & \\ \mathcal{O}(1) \otimes A & & \end{array}$$

- Equivariant: For all $j \in \mathbb{N}$ and $\sigma \in S_j$:

$$\begin{array}{ccc} \mathcal{O}(j) \otimes A^j & \xrightarrow{\sigma \otimes \sigma^{-1}} & \mathcal{O}(j) \otimes A^j \\ & \searrow \theta & \swarrow \theta \\ & A & \end{array}$$

Definition 2.10. Let $\mathcal{O}$ be an operad in $\mathcal{C}$, A an $\mathcal{O}$ -algebra. We call an object M in $\mathcal{C}$ with maps

$$\omega: \mathcal{O}(j) \otimes A^{j-1} \otimes M \rightarrow M$$

an A -*module* if it satisfies the following associativity, unitality and equivariance conditions:

2. Operads

- Associative: For all $\sum j_l = j$ in $\mathbb{N}$ the following diagrams commute:

$$\begin{array}{ccc}
 \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \cdots \otimes \mathcal{O}(j_k) \otimes A^{j-1} \otimes M & \xrightarrow{\gamma \otimes \text{Id}} & \mathcal{O}(j) \otimes A^{j-1} \otimes M \\
 \downarrow s & & \downarrow \omega \\
 \mathcal{O}(k) \otimes \left(\bigotimes_{l=1}^{k-1} \mathcal{O}(j_l) \otimes A^{j_l} \right) \otimes \mathcal{O}(j_k) \otimes A^{j_k-1} \otimes M & \xrightarrow{\text{Id} \otimes \theta^{k-1} \otimes \omega} & \mathcal{O}(k) \otimes A^{k-1} \otimes M \\
 & & \uparrow \omega \\
 & & M
 \end{array}$$

- Unital:

$$\begin{array}{ccc}
 \text{I} \otimes M & \xrightarrow{\cong} & M \\
 \eta \otimes \text{Id} \downarrow & \nearrow \omega & \\
 \mathcal{O}(1) \otimes A^0 \otimes M & &
 \end{array}$$

- Equivariant: For all $j \in \mathbb{N}$ and $\sigma \in S_{j-1} \subset S_j$, the following diagram commutes:

$$\begin{array}{ccc}
 \mathcal{O}(j) \otimes A^{j-1} \otimes M & \xrightarrow{\sigma \otimes \sigma^{-1} \otimes \text{Id}} & \mathcal{O}(j) \otimes A^{j-1} \otimes M \\
 \searrow \omega & & \swarrow \omega \\
 & M &
 \end{array}$$

Remark 2.11. We can of course also define algebras over planar operads by dropping the equivariance condition as there is no symmetric group action. So, algebras over the operad $\text{Ass}^\otimes$ which we defined in examples 2.2 are indeed monoids since the corresponding squares commute due to canonical isomorphisms:

$$\begin{array}{ccc}
 \text{Ass}^\otimes(2) \otimes \text{Ass}^\otimes(2) \otimes \text{Ass}^\otimes(1) \otimes A^3 & \xrightarrow{\cong} & \text{Ass}^\otimes(3) \otimes A^3 \\
 \downarrow \cong & & \downarrow \theta \\
 \text{Ass}^\otimes(2) \otimes (\text{Ass}^\otimes(2) \otimes A^2) \otimes (\text{Ass}^\otimes(1) \otimes A) & \xrightarrow{\text{Id} \otimes \theta^2} & \text{Ass}^\otimes(2) \otimes A^2 \\
 & & \uparrow \theta \\
 & & A \\
 & & \downarrow \theta \\
 \text{Ass}^\otimes(2) \otimes \text{Ass}^\otimes(1) \otimes \text{Ass}^\otimes(2) \otimes A^3 & \xrightarrow{\cong} & \text{Ass}^\otimes(3) \otimes A^3 \\
 \downarrow \cong & & \downarrow \theta \\
 \text{Ass}^\otimes(2) \otimes (\text{Ass}^\otimes(1) \otimes A) \otimes (\text{Ass}^\otimes(2) \otimes A^2) & \xrightarrow{\text{Id} \otimes \theta^2} & \text{Ass}^\otimes(2) \otimes A^2 \\
 & & \uparrow \theta \\
 & & A \\
 & & \downarrow \theta
 \end{array}$$

Examples 2.12. 1. A natural question is whether the associative planar operad and the associative operad in a category $\mathcal{C}$ coincide. For $\mathcal{C} = \mathbf{Set}$, we can equip the associative operad

$$\text{Ass}^{\otimes \text{planar}}_{\mathbf{Set}}(n) = *$$

with the trivial action of S_n . Yet this appears to be too restrictive: The equivariance condition implies that algebras over this operad are not just monoids but commutative monoids. To distinguish operations after acting by S_n , we can think of a more suited candidate: $\text{Ass}^\otimes(n) = S_n$ for all $n \in \mathbb{N}$ where S_n acts on $\text{Ass}^\otimes(n)$ by translation. The

2. Operads

composition map is defined by considering a sum of permutations and then permuting k blocks of length j_i for $1 \leq i \leq k$:

$$\begin{aligned} \gamma: \text{Ass}^\otimes(k) \times \text{Ass}^\otimes(j_1) \times \cdots \times \text{Ass}^\otimes(j_k) &\rightarrow \text{Ass}^\otimes(k) \\ (\sigma, \tau_1, \dots, \tau_k) &\mapsto \sigma(\tau_1 \oplus \cdots \oplus \tau_k) \end{aligned}$$

We might also refer to this operad as the symmetry operad $\mathcal{S}$ later on.

2. The operad $\text{Comm}^\otimes_{\mathbf{Set}}(n) = *$ with the trivial action of S_n is an interesting operad nevertheless as it is the *commutative operad* in the category $\mathbf{Set}$. The commutative operad in a category $\mathcal{C}$ is the operad such that its algebras are commutative monoids in $\mathcal{C}$.
3. The *little n -disk operad* D_n is an important example of an operad in the category of topological spaces $\mathbf{Top}$. For all $n \in \mathbb{N}$ we will define $D_n(k)$ as a topological space representing the space of k disjoint little disks inside the unit disk in the plane. Let $n \in \mathbb{N}$ and

$$D^n = \{(r_1, \dots, r_n) \in \mathbb{R}^n : r_1^2 + \cdots + r_n^2 \leq 1\} \subset \mathbb{R}^n$$

is the closed unit n -disk. A little n -disk is defined as the embedding $f: D^n \rightarrow D^n$ of the following form:

$$f(r_1, \dots, r_n) := (c_1, \dots, c_n) + \lambda(r_1, \dots, r_n)$$

for $(c_1, \dots, c_n) \in D^n$ and $\lambda \in \mathbb{R}$ such that

$$0 < \lambda \leq 1 - \sum_{i=1}^n c_i^2.$$

We denote the interior of a little disk by $\text{int}(f)$ and define for $k > 0$:

$$D_n(k) := \left\{ f_1 \sqcup \cdots \sqcup f_n \in C \left(\bigsqcup_{i=1}^k D^n, D^n \right) : \text{int}(f_i) \cap \text{int}(f_j) = \emptyset \text{ for all } i \neq j \right\}$$

We equip $C \left(\bigsqcup_{i=1}^k D^n, D^n \right)$ with the compact-open topology and $D_n(k)$ with the subspace topology. $D_n(0)$ is the one point space. There is an operad structure to be defined on

$$D_n = \{D_n(k)\}_{k \geq 0}.$$


The construction of the composition map

$$\gamma: D_n(k) \times D_n(l_1) \times \cdots \times D_n(l_k) \rightarrow D_n(l_1 + \cdots + l_k)$$

is geometrically interpreted as:

- Start with a configuration in $D_n(k)$, which gives k disks inside the unit disk.
- Each of the k disks is treated as a new "unit disk" in which a configuration from $D_n(l_1), D_n(l_2), \dots, D_n(l_k)$ can be inserted.
- The configurations from $D_n(l_1), D_n(l_2), \dots, D_n(l_k)$ are scaled down to fit inside the respective disks from the configuration in $D_n(k)$, and they are inserted.
- After placing all configurations inside the k disks, the boundaries of the k disks are erased, yielding a new configuration in $D_n(n_1 + \cdots + n_k)$.

2. Operads



Formally, this means that for $(f, (g_i)_{1 \leq i \leq k}) \in D_n(l_1) \times D_n(l_2) \times \cdots \times D_n(l_k)$ we define $\gamma((f, (g_i)_{1 \leq i \leq k}))$ via the following commuting diagram:

$$\begin{array}{ccc}
 \bigsqcup_{j=1}^l D^n & \xrightarrow{\gamma(f, (g_i)_{1 \leq i \leq k})} & D^n \\
 \downarrow \cong & & \uparrow f \\
 \bigsqcup_{i=1}^k \left(\bigsqcup_{p=1}^{l_k} D^n \right) & \xrightarrow{\bigsqcup g_i} & \bigsqcup_{i=1}^k D^n
 \end{array}$$

The unit is the identity configuration, where the single disk remains unchanged, i.e. the identity map $\text{Id}_{D^n} \in C(D^n, D^n)$. This satisfies the unitality condition for operads by definition, where inserting a single disk into another disk does not alter any configuration: $\gamma((f, (\text{Id})_{1 \leq i \leq k})) = f$ and $\gamma((\text{Id}, g)) = g$.

The composition map satisfies the associativity condition by definition.

The disks are labeled, and there is an action of the symmetric group S_k on $D_n(k)$, which permutes the labels of the disks: For $f_1 \sqcup \cdots \sqcup f_k \in D_n(k)$ and $\sigma \in S_k$:

$$\sigma(f_1 \sqcup \cdots \sqcup f_k) := f_{\sigma(1)} \sqcup \cdots \sqcup f_{\sigma(k)}$$

The composition map γ respects the action of S_k : For any $\sigma \in S_k$, $\tau_i \in S_{l_i}$ and

2. Operads

$(f, (g_i)_{1 \leq i \leq k}) \in \mathcal{D}_n(k) \times \mathcal{D}_n(l_1) \times \cdots \times \mathcal{D}_n(l_k)$ we have by definition

$$\begin{aligned} \sigma \circ \gamma((f, (g_i)_{1 \leq i \leq k})) &= \sigma \left(\bigsqcup_{i=1}^k f_i \circ g_i \right) \\ &= \bigsqcup_{i=1}^k f_{\sigma(i)} \circ g_{\sigma(i)} \\ &= \gamma((f_{\sigma(1)} \sqcup \cdots \sqcup f_{\sigma(k)}, g_{\sigma(1)}, \dots, g_{\sigma(k)}) \\ &= \gamma \circ (\sigma \times \sigma^{-1})((f, (g_i)_{1 \leq i \leq k})) \end{aligned}$$

and

$$\begin{aligned} (\tau_1 \oplus \cdots \oplus \tau_k) \circ \gamma((f, (g_i)_{1 \leq i \leq k})) &= (\tau_1 \oplus \cdots \oplus \tau_k) \left(\bigsqcup_{i=1}^k f_i \circ g_i \right) \\ &= \bigsqcup_{i=1}^k f_i \circ \tau_i(g_i) \\ &= \gamma((f, \tau_1(g_1), \dots, \tau_k(g_k))) \\ &= \gamma \circ (\tau_1 \times \cdots \times \tau_k)((f, (g_i)_{1 \leq i \leq k})). \end{aligned}$$

2.2 Colored Operads

Following [JL], the goal of this section is to introduce colored operads in such a way that we can define colored ∞ -operads later on. A symmetric monoidal category is an example of a colored operad. Thus, we will focus on this special case first. We will show that a symmetric monoidal category can be defined in a purely functorial way. After that we will then follow this approach and define the notion of a colored operad in a similar way.

Let us start by introducing some terminology from [Kerodon].

Definition 2.13. Let $P: \mathcal{C} \rightarrow \mathcal{D}$ be a functor of arbitrary categories and $f: x \rightarrow y$ a morphism in $\mathcal{C}$. We call f *cartesian* if for any morphism $g: z \rightarrow y$ in $\mathcal{C}$ and $\omega: P(z) \rightarrow P(x)$ in $\mathcal{D}$ with $P(f) \circ \omega = P(g)$, there exists a unique morphism $\omega': z \rightarrow x$ in $\mathcal{C}$ such that $g = f \circ \omega'$ and $P(\omega') = \omega$. In diagrams we have:

$$\begin{array}{ccc} \begin{array}{ccc} z & \xrightarrow{g} & y \\ \exists! \downarrow \vdots & \nearrow f & \\ x & & \end{array} & \xrightarrow{P} & \begin{array}{ccc} P(z) & \longrightarrow & P(y) \\ \downarrow & \nearrow & \\ P(x) & & \end{array} \end{array}$$

And vice versa, we call f *cocartesian* if we can flip arrows of the above case, i.e. for any morphism $g: y \rightarrow z$ in $\mathcal{C}$ and $\omega: P(x) \rightarrow P(z)$ in $\mathcal{D}$ with $\omega \circ P(f) = P(g)$, there is a unique morphism $\omega': x \rightarrow z$ such that $P(\omega') = \omega$ and $\omega' \circ f = g$. In a diagram this means the following:

$$\begin{array}{ccc} \begin{array}{ccc} y & \longrightarrow & z \\ & \searrow f & \uparrow \exists! \vdots \\ & & x \end{array} & \xrightarrow{P} & \begin{array}{ccc} P(y) & \longrightarrow & P(z) \\ & \searrow & \uparrow \\ & & P(x) \end{array} \end{array}$$

Note that equivalently we can define a cocartesian morphism by requiring a bijection:

$$\mathrm{Hom}_{\mathcal{C}}(y, z) \times_{\mathrm{Hom}_{\mathcal{D}}(P(y), P(z))} \mathrm{Hom}_{\mathcal{D}}(P(x), P(z)) \cong \mathrm{Hom}_{\mathcal{C}}(x, z)$$

2. Operads

where $\text{Hom}_{\mathcal{C}}(y, z) \times_{\text{Hom}_{\mathcal{D}}(P(y), P(z))} \text{Hom}_{\mathcal{D}}(P(x), P(z))$ is the pullback of applying the functor P to $\text{Hom}_{\mathcal{C}}(y, z)$ and precomposing with $P(f): P(y) \rightarrow P(x)$. So, for the bijection we send a pair (g, ω) to the unique morphism ω' and vice versa we send a morphism $h: x \rightarrow z$ to the pair $(h \circ f, P(h))$.

Definition 2.14. Let $P: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. P is called a *Grothendieck fibration* if for every object y in $\mathcal{C}$ and morphisms $f': x' \rightarrow P(y)$ there exists a cartesian morphism $f: x \rightarrow y$ such that $f' = P(f)$. We call f a cartesian lift of f' and f' a cleavage. Dually we define the notion of a *Grothendieck op-fibration*, i.e. P is a Grothendieck op-fibration if for every object x in $\mathcal{C}$ and morphism $f': P(x) \rightarrow y'$ there exists a cocartesian morphism $f: x \rightarrow y$ such that $P(f) = f'$.

If not stated otherwise, when we talk about fibrations and op-fibrations in this thesis, we will always mean Grothendieck fibrations and Grothendieck op-fibrations, respectively. Also note, that in [Kerodon], these terms are referred to as cartesian and cocartesian fibration. However, we chose to use the terminology used in [JL].

In the next part we will construct a functor having certain properties from a symmetric monoidal category. We will then see that the functor encodes all the necessary data of the symmetric monoidal category.

Definition 2.15. Let $\mathcal{C}$ be a symmetric monoidal category. We define a new category $\mathcal{C}^{\otimes}$ in the following way:

- Objects of $\mathcal{C}^{\otimes}$ are finite sequences $[C_1, \dots, C_n]$ where C_i is an object in $\mathcal{C}$ for all i . It is possible that the sequence is empty.
- A morphism $f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]$ in $\mathcal{C}^{\otimes}$ consists of a subset $S_f \subset \{1, \dots, n\}$, a map of finite sets $\alpha_f: S_f \rightarrow \{1, \dots, m\}$ and a family of morphisms

$$\left\{ f_j: \bigotimes_{\alpha_f(i)=j} C_i \rightarrow C'_j \right\}_{1 \leq j \leq m}$$

in $\mathcal{C}$.

- Composition is defined as follows: Let

$$f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m] \quad \text{and} \quad g: [C'_1, \dots, C'_m] \rightarrow [C''_1, \dots, C''_k]$$

be morphisms in $\mathcal{C}^{\otimes}$, then

$$g \circ f: [C_1, \dots, C_n] \rightarrow [C''_1, \dots, C''_k]$$

is defined via $S_{g \circ f} = \alpha_f^{-1}(S_g)$, $\alpha_{g \circ f} = \alpha_g \circ \alpha_f|_{S_{g \circ f}}$ and for all $1 \leq l \leq k$ we have:

$$\bigotimes_{\alpha_{g \circ f}(i)=l} C_i \cong \bigotimes_{\alpha_g(j)=k} \bigotimes_{\alpha_f(i)=j} C_i \rightarrow \bigotimes_{\alpha_g(j)=k} C'_j \rightarrow C''_l.$$

Let I be a finite set. We write $I_* = I \coprod *$.

Definition 2.16. We define Fin_* , the *category of finite pointed sets*, as follows:

- For each $n \in \mathbb{N}$, we write $\langle n \rangle^\circ := \{1, \dots, n\}$. We define the objects of Fin_* as $\langle n \rangle := \langle n \rangle_*^\circ$.
- A morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ is a map of finite sets where $\alpha(*) = *$

2. Operads

It is clear that this defines a category. Further note that this category is equivalent to the category of finite sets with a distinguished point since for any finite set I we have an isomorphism $I_* \cong \langle n \rangle$, where n is the cardinality of I . For all pairs of $1 \leq i \leq n$ we write $\rho^i: \langle n \rangle \rightarrow \langle 1 \rangle$ for the morphism in $\mathcal{F}in_*$ which is defined by

$$\rho^i(j) = \begin{cases} 1 & i = j \\ * & \text{else.} \end{cases}$$

For any symmetric monoidal category $\mathcal{C}$ there exists a forgetful functor

$$P: \mathcal{C}^\otimes \rightarrow \mathcal{F}in_*$$

which sends a sequence $[C_1, \dots, C_n]$ to $\langle n \rangle$ and a morphism $f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]$ to $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ with

$$\alpha(x) = \begin{cases} \alpha_f(x) & x \in S_f \\ * & \text{else.} \end{cases}$$

Remark 2.17. Let $\mathcal{C}_0$ denote the trivial category of one object and one morphism. We claim that the forgetful functor $P: \mathcal{C}_0^\otimes \rightarrow \mathcal{F}in_*$ is an isomorphism of categories. This holds, since for all morphisms in $\mathcal{C}_0^\otimes$ the family of morphisms f_j in $\mathcal{C}_0$ are trivial, in other words there is nothing to forget. Hence, $\mathcal{F}in_*$ is obtained as the simplest example of the construction of a $\mathcal{C}^\otimes$ category.

Furthermore, we note that the forgetful functor $P: \mathcal{C}^\otimes \rightarrow \mathcal{F}in_*$ can be obtained from a unique symmetric monoidal functor $F: \mathcal{C} \rightarrow \mathcal{C}_0$ using the functoriality of F .

Remark 2.18. For every object $\langle n \rangle$ we denote the fiber category by $\mathcal{C}_{\langle n \rangle}$, which is defined to be the pullback of the following diagram

$$\begin{array}{ccc} \mathcal{C}_{\langle n \rangle}^\otimes & \dashrightarrow & \mathcal{C}^\otimes \\ \downarrow & & \downarrow P \\ \{\langle n \rangle\} & \hookrightarrow & \mathcal{F}in_* \end{array}$$

where $\{\langle n \rangle\}$ is interpreted as the trivial one object category. In other words, $\mathcal{C}_{\langle n \rangle}$ consists of all collections of objects of length n and all morphisms f such that $\alpha_f = \text{Id}_{\langle n \rangle}$.

We are now ready to state two very important properties of the functor $P: \mathcal{C}^\otimes \rightarrow \mathcal{F}in_*$, which as it turns out is all we need to know.

Proposition 2.19. *Let $P: \mathcal{C}^\otimes \rightarrow \mathcal{F}in_*$ be the forgetful functor as describes above. Then, the following conditions hold:*

(M1) **Op-fibration condition:** *P is a Grothendieck op-fibration.*

(M2) **Segal condition:** *We have that $\mathcal{C} \cong \mathcal{C}_{\langle 1 \rangle}$ and more precisely we obtain that $\mathcal{C}_{\langle n \rangle}^\otimes \cong \mathcal{C}^n$ where $\mathcal{C}^n$ denotes the n -fold product of copies of $\mathcal{C}$.*

Proof. (M1) Let $[C_1, \dots, C_n]$ be an object in $\mathcal{C}^\otimes$ and $f': \langle n \rangle \rightarrow \langle m \rangle$ a morphism in $\mathcal{F}in_*$. We need to construct a cocartesian morphism $f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]$ such that $P(f) = f'$. We define $\alpha_f: S_f \rightarrow \langle m \rangle^\circ$ as $\alpha_f := f'|_{S_f}$ where $S_f := f'^{-1}(\langle m \rangle^\circ)$. Further we define a selection of objects in $\mathcal{C}$ for $1 \leq j \leq m$:

$$C'_j := \begin{cases} \bigotimes_{\alpha_f(i)=j} C_i & j \in \alpha_f(S_f) \\ \text{I} & \text{else} \end{cases}$$

For $f_j := \text{Id}_{C'_j}$ we have then constructed a well defined morphism

$$f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]$$

2. Operads

such that $P(f) = f'$. We are left to show that f is indeed cocartesian. Therefore, let $g: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_l]$ be a morphism in $\mathcal{C}^\otimes$ and $\omega': \langle m \rangle \rightarrow \langle l \rangle$ a morphism in $\mathcal{F}\text{in}_*$ such that $P(g) = \omega' \circ P(f)$. We need to find a unique morphism

$$\omega: [C'_1, \dots, C'_m] \rightarrow [C''_1, \dots, C''_l]$$

such that $g = \omega \circ f$ and $P(\omega) = \omega'$. We are forced to define $\alpha_\omega := \omega'|_{S_\omega}$ where

$$S_\omega = \omega'^{-1}(\langle l \rangle^\circ).$$

By construction we have $\alpha_g = \alpha_\omega \circ \alpha_f$ and by definition of C'_j we are forced to define for all $1 \leq j \leq l$:

$$\omega_j: \bigotimes_{\alpha_\omega(k)=j} C'_k \cong \bigotimes_{\alpha_\omega(k)=j} \bigotimes_{\alpha_f(i)=k} C_i \cong \bigotimes_{\alpha_g(k)=j} C_k \xrightarrow{g_j} C''_j$$

Uniqueness of ω follows from $\mathcal{C}$ being a symmetric monoidal category. Hence, f is cocartesian.

(M2) Let $F: \mathcal{C}_{\langle 1 \rangle}^\otimes \rightarrow \mathcal{C}$ be a functor that sends $[C]$ to C . Let $f: [C] \rightarrow [C']$ be a morphism in $\mathcal{C}_{\langle 1 \rangle}^\otimes$, then $\alpha_f = \text{Id}_{\{1\}}$ and there is just one morphism $f_1: C \rightarrow C'$ corresponding to f . So, we define $F(f) = f_1$. It follows by definition that F is fully faithful and essentially surjective, i.e. an equivalence of categories.

First, we will show that for any morphism $\langle n \rangle \rightarrow \langle m \rangle$ the op-fibration condition (M1) induces a functor of fiber categories $A: \mathcal{C}_{\langle n \rangle}^\otimes \rightarrow \mathcal{C}_{\langle m \rangle}^\otimes$: Let $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ be a morphism in $\mathcal{F}\text{in}_*$ and $[C_1, \dots, C_n]$ an object in $\mathcal{C}_{\langle n \rangle}^\otimes$. By the op-fibration condition (M1), we can find a cocartesian morphism

$$f_\alpha: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m].$$

Therefore, we can define

$$A([C_1, \dots, C_n]) := [C'_1, \dots, C'_m].$$

To see that the choice of $[C'_1, \dots, C'_m]$ is unique up to isomorphism, let

$$f'_\alpha: [C_1, \dots, C_n] \rightarrow [\tilde{C}'_1, \dots, \tilde{C}'_m]$$

be a different choice of cocartesian morphism for $[C_1, \dots, C_n]$ and α . Then there are morphisms

$$g: [C'_1, \dots, C'_m] \rightarrow [\tilde{C}'_1, \dots, \tilde{C}'_m] \quad \text{and} \quad g': [\tilde{C}'_1, \dots, \tilde{C}'_m] \rightarrow [C'_1, \dots, C'_m]$$

corresponding to the cocartesian property of f_α and f'_α respectively. Because of uniqueness we obtain $g \circ g' = \text{Id}_{[\tilde{C}'_1, \dots, \tilde{C}'_m]}$ and $g' \circ g = \text{Id}_{[C'_1, \dots, C'_m]}$. Therefore, the choice of a cocartesian morphism is unique up to isomorphism.

Let $f: [C_1, \dots, C_n] \rightarrow [\bar{C}_1, \dots, \bar{C}_n]$ be a morphism in $\mathcal{C}_{\langle n \rangle}^\otimes$. We obtain a unique morphism $f': [\bar{C}_1, \dots, \bar{C}_n] \rightarrow [\bar{C}'_1, \dots, \bar{C}'_m]$ from the fact that f_α is cocartesian.

$$\begin{array}{ccc} [C_1, \dots, C_n] & \xrightarrow{f_\alpha} & [C'_1, \dots, C'_m] \\ \downarrow f & & \downarrow f' \\ [\bar{C}_1, \dots, \bar{C}_n] & \xrightarrow{\bar{f}_\alpha} & [\bar{C}'_1, \dots, \bar{C}'_m] \end{array} \xrightarrow{P} \begin{array}{ccc} \langle n \rangle & \xrightarrow{\alpha} & \langle m \rangle \\ \downarrow \text{Id} & & \downarrow \text{Id} \\ \langle n \rangle & \xrightarrow{\alpha} & \langle m \rangle \end{array}$$

2. Operads

Hence, we can define $A(f) := f'$. Due to uniqueness of f' , A preserves composition and maps identities to identities.

Let $P^i: \mathcal{C}_{\langle n \rangle}^{\otimes} \rightarrow \mathcal{C}_{\langle 1 \rangle}^{\otimes} \cong \mathcal{C}$ be the functor corresponding to $\rho^i: \langle n \rangle \rightarrow \langle 1 \rangle$ which we defined previously for all $1 \leq i \leq n$. Then from the universal property of the product, it follows that

$$\prod_{1 \leq i \leq n} P^i: \mathcal{C}_{\langle n \rangle}^{\otimes} \rightarrow \mathcal{C}^n$$

is an equivalence of categories. □

Note that in the following we might refer to the constructed functor $\mathcal{C}_{\langle n \rangle}^{\otimes} \rightarrow \mathcal{C}_{\langle m \rangle}^{\otimes}$ originated from $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ also by α .

We are now well prepared to see that the symmetric monoidal structure on $\mathcal{C}$ is encoded in $\mathcal{C}^{\otimes}$ and its forgetful functor $P: \mathcal{C}^{\otimes} \rightarrow \mathcal{Fin}_*$ satisfying the above conditions. In other words, for any category $\mathcal{D}$ together with a functor $P: \mathcal{D} \rightarrow \mathcal{Fin}_*$ satisfying the op-fibration condition (M1) and the Segal condition (M2), we can construct a symmetric monoidal category $\mathcal{C}$ such that $\mathcal{D} \cong \mathcal{C}^{\otimes}$ and P is the usual forgetful functor. The right candidate for $\mathcal{C}$ is clearly $\mathcal{D}_{\langle 1 \rangle}$.

Proposition 2.20. *The construction of $\pi: \mathcal{C}^{\otimes} \rightarrow \mathcal{Fin}_*$ from a symmetric monoidal category $\mathcal{C}$ given above is a bijection:*

$$\{\text{symmetric monoidal categories}\} \cong \{\text{op-fibrations } \mathcal{D} \rightarrow \mathcal{Fin}_* \text{ satisfying (M2)}\}$$

Proof. Let $\mathcal{D}$ be a category and $P: \mathcal{D} \rightarrow \mathcal{Fin}_*$ a functor satisfying the op-fibration condition (M1) and the Segal condition (M2). We will equip $\mathcal{C} := \mathcal{D}_{\langle 1 \rangle}$ with a symmetric monoidal structure and see that there is an equivalence of categories $\mathcal{C}^{\otimes} \cong \mathcal{D}$.

1. For the map $\alpha: \langle 2 \rangle \rightarrow \langle 1 \rangle$ with $\alpha^{-1}(1) = \{1, 2\}$ we obtain the following functor using (M2):

$$\bigotimes: \mathcal{C} \times \mathcal{C} \cong \mathcal{D}_{\langle 2 \rangle} \xrightarrow{\alpha} \mathcal{C}$$

2. There is a unique morphism $\eta: \langle 0 \rangle \rightarrow \langle 1 \rangle$ in $\mathcal{Fin}_*$ defining a functor $\mathcal{D}_{\langle 0 \rangle} \rightarrow \mathcal{C}$ which gives us a unique object I in $\mathcal{C}$. This object I will serve as the tensor unit which follows from the following commuting diagram:

$$\begin{array}{ccccc} \mathcal{D}_{\langle 0 \rangle} \times \mathcal{D}_{\langle 1 \rangle} & \xrightarrow{\eta \times \text{Id}} & \mathcal{D}_{\langle 1 \rangle} \times \mathcal{D}_{\langle 1 \rangle} & \xrightarrow{\cong} & \mathcal{D}_{\langle 2 \rangle} \\ \uparrow \cong & & & & \searrow \otimes \\ \mathcal{C} & & & & \mathcal{D}_{\langle 1 \rangle} \\ \downarrow \cong & & & & \nearrow \otimes \\ \mathcal{D}_{\langle 1 \rangle} \times \mathcal{D}_{\langle 0 \rangle} & \xrightarrow{\text{Id} \times \eta} & \mathcal{D}_{\langle 1 \rangle} \times \mathcal{D}_{\langle 1 \rangle} & \xrightarrow{\cong} & \mathcal{D}_{\langle 2 \rangle} \end{array}$$

3. Furthermore, we need a symmetric braiding. This we obtain from the morphism

$$\begin{aligned} \beta: \langle 2 \rangle &\rightarrow \langle 2 \rangle \\ 1 &\mapsto 2 \\ 2 &\mapsto 1. \end{aligned}$$

We note that $\alpha \circ \beta = \alpha$ and $\rho^i \beta = \rho^{\beta(i)}$. So, we obtain the following commuting diagramm:

$$\begin{array}{ccccc} \mathcal{D}_{\langle 1 \rangle} \times \mathcal{D}_{\langle 1 \rangle} & \xleftarrow{\rho^1 \times \rho^2} & \mathcal{D}_{\langle 2 \rangle} & \xrightarrow{\alpha} & \mathcal{D}_{\langle 1 \rangle} \\ & \swarrow \rho^2 \times \rho^1 & \downarrow \beta & \searrow \alpha & \\ & & \mathcal{D}_{\langle 2 \rangle} & & \end{array}$$

2. Operads

Hence, $A \otimes B \cong B \otimes A$ holds.

4. The associator corresponds to the following maps in $\mathcal{Fin}_*$:

$$\tau_i^n: \langle n \rangle \rightarrow \langle n-1 \rangle$$

$$j \mapsto \begin{cases} j & 1 \leq j \leq i \\ j-1 & i < j \leq n \\ * & j = * \end{cases}$$

So, the following commuting diagram

$$\begin{array}{ccc} \langle 3 \rangle & \xrightarrow{\tau_1^3} & \langle 2 \rangle \\ \tau_2^3 \downarrow & & \downarrow \tau_1^2 \\ \langle 2 \rangle & \xrightarrow{\tau_1^2} & \langle 1 \rangle \end{array}$$

translates to $(A \otimes B) \otimes C \cong A \otimes (B \otimes C)$. We need to see the pentagon identities:

$$\begin{array}{ccc} & (A \otimes B) \otimes (C \otimes D) & \\ \swarrow & & \searrow \\ A \otimes (B \otimes (C \otimes D)) & & ((A \otimes B) \otimes C) \otimes D \\ \uparrow & & \downarrow \\ A \otimes ((B \otimes C) \otimes D) & \xleftarrow{\quad} & (A \otimes (B \otimes C)) \otimes D \end{array}$$

The commutativity of the following diagram in $\mathcal{Fin}_*$ translates to the pentagon identities:

$$\begin{array}{ccccc} & & \langle 3 \rangle & \xrightarrow{\tau_2^3} & \langle 2 \rangle \\ & \nearrow \tau_1^4 & & \searrow \tau_1^3 & \searrow \tau_1^2 \\ \langle 4 \rangle & \xrightarrow{\tau_2^4} & \langle 3 \rangle & \xrightarrow{\tau_1^3} & \langle 2 \rangle & \xrightarrow{\tau_1^2} & \langle 1 \rangle \\ & \searrow \tau_3^4 & & \searrow \tau_2^3 & \nearrow \tau_1^2 \\ & & \langle 3 \rangle & \xrightarrow{\tau_2^3} & \langle 2 \rangle \end{array}$$

$$\begin{array}{ll} \tau_1^2 \circ \tau_2^3 \circ \tau_1^4 & \rightsquigarrow (A \otimes B) \otimes (C \otimes D) \\ \tau_1^2 \circ \tau_1^3 \circ \tau_1^4 & \rightsquigarrow ((A \otimes B) \otimes C) \otimes D \\ \tau_1^2 \circ \tau_1^3 \circ \tau_2^4 & \rightsquigarrow (A \otimes (B \otimes C)) \otimes D \\ \tau_1^2 \circ \tau_2^3 \circ \tau_2^4 & \rightsquigarrow A \otimes ((B \otimes C) \otimes D) \\ \tau_1^2 \circ \tau_2^3 \circ \tau_3^4 & \rightsquigarrow A \otimes (B \otimes (C \otimes D)) \end{array}$$

This construction is in bijection to the previous given construction. $\square$

Remark 2.21. With some adjustments this construction can also be done for a monoidal category $\mathcal{C}$. The category $\mathcal{C}^\otimes$ is defined analogously to the symmetric case. But we choose ordered indexing sets $[n]^\circ := \{0 < \dots < n\}$ and for a morphism $f: [C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]$ we require that α_f preserves the order. Further, we can define the *pointed simplex category* Δ_* as follows:

- For each $n \in \mathbb{N}$ we define an object in Δ_* as $[n] := [n]^\circ \sqcup \{*\}$.

2. Operads

- The single pointed set $\{*\}$ is an object in Δ_* . There are morphisms $\{*\} \hookrightarrow [n]$ for all n .
- A morphism $\alpha: [n] \rightarrow [m]$ is a order preserving map on $\alpha^{-1}([m])$ and $\alpha(*) = *$.

Note that for all pairs $0 \leq i \leq n$, we have $\rho^i: [n] \rightarrow [0]$ in Δ_* with

$$\rho^i(j) = \begin{cases} 0 & i = j \\ * & \text{else.} \end{cases}$$

Similarly to the symmetric case, we obtain a forgetful functor:

$$P: \mathcal{C}^\otimes \rightarrow \Delta_*$$

This functor is a op-fibration. The proof in proposition 2.19 applies. Note that for the proof the fact that morphisms in Δ_* preserve order is essential, otherwise we could not define

$$\omega_j: \bigotimes_{\alpha_\omega(k)=j} C'_k \cong \bigotimes_{\alpha_\omega(k)=j} \bigotimes_{\alpha_f(i)=k} C_i \cong \bigotimes_{\alpha_g(k)=j} C_k \xrightarrow{g_j} C''_j$$

in a monoidal category $\mathcal{C}$. Further, the Segal condition (M2) holds as it is a result of the op-fibration condition (M1) and the fact that ρ^i is a morphism in Δ_* . It then also follows that this similar construction is again a bijection

$$\{\text{monoidal categories}\} \cong \{\text{op-fibration } \mathcal{D} \rightarrow \Delta_* \text{ satisfying (M2)}\}.$$

The proof for this is analogously to proposition 2.20 as all morphisms in Fin_* that induce a monoidal structure on $\mathcal{C}$ are order preserving and can be considered in Δ_* .

We will now introduce the notion of a colored operad and will move on to see that, similarly to describing a symmetric monoidal structure via a functor satisfying the op-fibration condition (M1) and Segal condition (M2), there is also a functor satisfying similar properties that encode the datum of a colored operad.

Definition 2.22. A *colored operad* $\mathcal{O}$ consists of the following data:

- A collection $\{X, Y, Z; \dots\}$ which we call objects or colors of the operad.
- For every finite collection of objects $\{X_i\}_{i \in I}$ and object Y in $\mathcal{O}$, there is a set $\text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in I}, Y)$ called the set of morphisms from $\{X_i\}_{i \in I}$ to Y .
- For any finite map $I \rightarrow J$ with fibers denoted by I_j for all $j \in J$, collections of objects $\{X_i\}_{i \in I}$ and $\{Y_j\}_{j \in J}$ and object Z , there is a composition map

$$\prod_{j \in J} \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in I_j}, Y_j) \times \text{Mul}_{\mathcal{O}}(\{Y_j\}_{j \in J}, Z) \rightarrow \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in I}, Z)$$

- There is a collection of morphisms $\{\text{Id}_X \in \text{Mul}_{\mathcal{O}}(\{X\}, X)\}_{X \in \mathcal{O}}$ being a left and right unit in the composition, i.e. the following diagrams commute:

$$\begin{array}{ccc} \text{Mul}_{\mathcal{O}}(X_I, Y) & \xrightarrow{\cong} & \text{Mul}_{\mathcal{O}}(X_I, Y) \times \{\text{Id}_Y\} \\ \downarrow \text{Id} & & \downarrow \\ \text{Mul}_{\mathcal{O}}(X_I, Y) & \longleftarrow & \text{Mul}_{\mathcal{O}}(X_I, Y) \times \text{Mul}_{\mathcal{O}}((Y), Y) \end{array}$$

$$\begin{array}{ccc} \text{Mul}_{\mathcal{O}}(X_I, Y) & \xrightarrow{\cong} & (\prod_{i \in I} \{\text{Id}_{X_i}\}) \times \text{Mul}_{\mathcal{O}}(X_I, Y) \\ \downarrow \text{Id} & & \downarrow \\ \text{Mul}_{\mathcal{O}}(X_I, Y) & \longleftarrow & (\prod_{i \in I} \text{Mul}_{\mathcal{O}}((X_i), X_i)) \times \text{Mul}_{\mathcal{O}}(X_I, Y) \end{array}$$

2. Operads

- (v) For every sequence of finite maps $I \rightarrow J \rightarrow K$, collections of objects $X_I = \{X_i\}_{i \in I}$, $Y_J = \{Y_j\}_{j \in J}$, $W_K = \{W_k\}_{k \in K}$ and object $Z \in \mathcal{O}$ the following diagram commutes:

make diagram prettier

$$\begin{array}{ccc}
 \prod_{j \in J} \text{Mul}_{\mathcal{O}}(X_{I_j}, Y_j) \times \prod_{k \in K} \text{Mul}_{\mathcal{O}}(Y_j, W_k) \times \text{Mul}_{\mathcal{O}}(W_K, Z) & \xrightarrow{\text{Id} \times \gamma} & \prod_{j \in J} \text{Mul}_{\mathcal{O}}(X_{I_j}, Y_j) \times \text{Mul}_{\mathcal{O}}(Y_J, Z) \\
 \downarrow (\gamma \circ s) \times \text{Id} & & \downarrow \\
 \prod_{k \in K} \text{Mul}_{\mathcal{O}}(X_{I_k}, W_k) \times \text{Mul}_{\mathcal{O}}(W_K, Z) & \xrightarrow{\quad} & \text{Mul}_{\mathcal{O}}(X_I, Z)
 \end{array}$$

Remark 2.23. When we drop the symmetric structure we obtain a definition of a colored operad what we call colored planar operad. In other word we consider the indexing sets I in definition 2.22 to be finite ordered sets and maps between indexing sets have to be order preserving.

Remark 2.24. We note that there is a forgetful functor from the categories of colored operads to the category of small categories. Hence, for every colored operad $\mathcal{O}$ there is an underlying category which we also denote by $\mathcal{O}$. Its objects are the colors of $\mathcal{O}$ and for $X, Y \in \mathcal{O}$ we define the morphisms

$$\text{Hom}(X, Y) := \text{Mul}_{\mathcal{O}}(\{X\}, Y).$$

Composition and identity hold since they hold in $\mathcal{O}$. Thus, a colored operad can be viewed as a category with additional structure. Since we have some sort of collection of multilinear maps, some also refer to colored operads as multicategories.

Examples 2.25. 1. A colored operad with one color is a symmetric operad in **Set** as defined in definition 2.8. Let $\mathcal{O}$ be a colored operad with one color X . We will construct a symmetric operad $\mathcal{P}$ from $\mathcal{O}$. Since $\mathcal{O}$ is a colored operad with only one color, we define the family of objects $\mathcal{P}(n)$ for $n \in \mathbb{N}$ as:

$$\mathcal{P}(n) := \text{Mul}_{\mathcal{O}}(\{X\}_{1 \leq i \leq n}, X).$$

The composition maps from $\mathcal{O}$ for natural number indexing then defines the composition maps of $\mathcal{P}$

$$\gamma : \mathcal{P}(k) \times \mathcal{P}(j_1) \times \cdots \times \mathcal{P}(j_k) \rightarrow \mathcal{P}(j)$$

where $j = \sum_{i=1}^k j_i$ and $\eta : \{*\} \rightarrow \mathcal{P}(1)$ is given by Id_X . Unitality and associativity follow from the corresponding conditions in $\mathcal{O}$. However, in a colored operad (even with one color), the set of morphisms depends not only on n , the cardinality of the indexing set, but also on the specific choice of finite indexing set. Thus, for two sets N_1 and N_2 with the same number of elements, you could potentially have different morphism sets:

$$\text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in N_1}, Y) \quad \text{and} \quad \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in N_2}, Y).$$

This raises the question of how the structures of a colored operad and an operad reconcile. The reason these two different setups are still equivalent lies in the symmetric group action on the set of morphisms. The permutation of the inputs ensures that any specific labeling of the inputs (as represented by the different finite sets N_1 and N_2) can be transformed into each other via the action of S_n . Hence, although the morphisms in the colored operad are indexed by specific sets, they behave the same way up to this permutation action. Therefore, the group action effectively collapses the dependence on the specific finite sets. So, while the morphism sets in a colored operad are technically different for different finite sets, the permutation symmetry makes these differences irrelevant, aligning the structure with that of an operad.

2. Operads

2. A colored planar operad with one color is a planar operad in **Set** as defined in definition 2.1. Let $\mathcal{P}$ be a colored planar operad $\mathcal{P}$. We set for $n \in \mathbb{N}$:

$$\mathcal{P}(n) := \text{Mul}_{\mathcal{P}}((X)_{1 \leq i \leq n}, X)$$

The composition map and operadic unit we obtain similarly as in the example for a one colored operad. However, as the indexing maps need to be order preserving we will not obtain a permutation of labels as there is only one bijective order preserving map between finite ordered sets.

Examples 2.26. 1. Let $\mathcal{C}$ be a symmetric monoidal category. By setting for a finite family of objects $\{X_i\}_{i \in I}$ and object Y in $\mathcal{C}$

$$\text{Mul}_{\mathcal{C}}(\{X_i\}_{i \in I}, Y) := \text{Hom}(\otimes_{i \in I} X_i, Y)$$

we can define a colored operad $\mathcal{C}$. The composition is given by

$$\prod_{j \in J} \text{Hom}(\otimes_{i \in I_j} X_i, Y_j) \times \text{Hom}(\otimes_{j \in J} Y_j, Z) \rightarrow \text{Hom}(\otimes_{i \in I} X_i, Z)$$

$$((f_j)_{j \in J}, g) \mapsto g \circ (\otimes_{j \in J} f_j)$$

with $\{\text{Id}_X\}_{X \in \mathcal{C}}$ being the unit and associativity holds since for $(f_j: \otimes_{i \in I_j} X_i \rightarrow Y_j)_{j \in J}$, $(g_k: \otimes_{j \in J_k} Y_j \rightarrow W_k)_{k \in K}$, $h: \otimes_{k \in K} W_k \rightarrow Z$ we have

$$h \circ \bigotimes_{k \in K} \left(g_k \circ \bigotimes_{j \in J_k} f_j \right) = h \circ \bigotimes_{k \in K} g_k \circ \bigotimes_{j \in J} f_j.$$

Further, we can reconstruct the symmetric monoidal structure of $\mathcal{C}$ from this colored operad: Let $F_{X,Y}: \mathcal{C} \rightarrow \mathbf{Set}$ be the functor sending an object Z to $\text{Mul}_{\mathcal{C}}(\{X, Y\}, Z)$ then by definition and from Yoneda's lemma follows that F is uniquely (up to isomorphism) represented by $X \otimes Y$. Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be morphisms in $\mathcal{C}$ and

$$\text{Mul}_{\mathcal{C}}(\{X\}, X') \times \text{Mul}_{\mathcal{C}}(\{Y\}, Y') \times \text{Mul}_{\mathcal{C}}(\{X', Y'\}, X' \otimes Y') \xrightarrow{\sim} \text{Mul}_{\mathcal{C}}(\{X, Y\}, X' \otimes Y')$$

the composition map. Then, we define $f \otimes g := \gamma((f, g, \text{Id}))$. The tensor unit I is the representing object of the functor sending an object Z in $\mathcal{C}$ to $\text{Mul}_{\mathcal{C}}(\emptyset, Z)$. This example is even clearer when we have the functorial definition of a colored operad later on. As it will be a generalization of proposition 2.19 and proposition 2.20 illustrates how to reconstruct the symmetric monoidal category.

2. Analogously, a monoidal category $\mathcal{C}$ is an example of a colored planar operad.

Definition 2.27. Let $\mathcal{O}$ be a colored operad. We define a category $\mathcal{O}^{\otimes}$ in the following way:

- (i) The objects of $\mathcal{O}^{\otimes}$ are finite sequences of colors of $\mathcal{O}$.
- (ii) Let $\{X_i\}_{1 \leq i \leq m}$ and $\{Y_j\}_{1 \leq j \leq n}$ be objects in $\mathcal{O}^{\otimes}$. A morphism

$$\phi: \{X_i\}_{1 \leq i \leq m} \rightarrow \{Y_j\}_{1 \leq j \leq n}$$

is given by a map $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ in Fin_* together with a family of morphisms

$$\{\phi_j \in \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in \alpha^{-1}(j)}, Y_j)\}_{1 \leq j \leq n}$$

in $\mathcal{O}$.

2. Operads

- (iii) Composition of morphisms is determined by the composition in $\mathcal{Fin}_*$ and $\mathcal{O}$, i.e. for morphisms $\phi: X \rightarrow Y$ and $\phi': Y \rightarrow Z$ in $\mathcal{O}^\otimes$ with $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ and $\alpha': \langle m \rangle \rightarrow \langle k \rangle$, we have $\bar{\alpha} = \alpha' \circ \alpha$ and

$$\{(\phi' \circ \phi)_l \in \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in \bar{\alpha}^{-1}(l)}, Z_l)\}_{1 \leq l \leq k}$$

where $(\phi' \circ \phi)_l$ is applying the composition map

$$\gamma: \prod_{j \in \alpha'^{-1}(l)} \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in \alpha^{-1}(j)}, Y_j) \times \text{Mul}_{\mathcal{O}}(\{Y_i\}_{i \in \alpha'^{-1}(l)}, Z_l) \rightarrow \text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in \bar{\alpha}^{-1}(l)}, Z_l)$$

Remark 2.28. For a planar colored operad $\mathcal{P}$, we can similarly define a category $\mathcal{P}^\otimes$ by requiring $\alpha: [m] \rightarrow [n]$ to be morphism in the simplex category with a distinguished point Δ_* .

Let $\mathcal{O}$ be a colored operad. Then similarly as for the definition 2.15 we have a forgetful functor

$$\pi: \mathcal{O}^\otimes \rightarrow \mathcal{Fin}_*$$

from which we can reconstruct the underlying colored operad $\mathcal{O}$ (up to canonical equivalence): First, note that in general π does not have to be a op-fibration. For an arbitrary morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ and a collection $\{X_i\}_{i \in \langle n \rangle}$ there is no canonical choice for a collection $\{X'_j\}_{j \in \langle m \rangle}$ with morphism $f: \{X_i\}_{i \in \langle n \rangle} \rightarrow \{X'_j\}_{j \in \langle m \rangle}$ such that $\pi(f) = \alpha$ as in the proof of proposition 2.19. However, there is a canonical choice if α is a bijection on $\alpha^{-1}(\langle m \rangle^\circ)$. Then for all $j \in \langle m \rangle$ there is exactly one $i_j \in \langle n \rangle$ with $\alpha(i_j) = j$. Hence, we define $X'_j := X_{i_j}$ and for all $j \in J$ we have $f_j = \text{Id}_{X_{i_j}}$. This defines a morphism $f: \{X_i\}_{i \in I} \rightarrow \{X'_j\}_{j \in J}$.

It is left to show that f is in fact cocartesian. Let $g: \{X_i\}_{i \in I} \rightarrow \{X''_k\}_{k \in \langle l \rangle}$ be a morphism in $\mathcal{O}^\otimes$ and $\omega: \langle m \rangle \rightarrow \langle l \rangle$ a morphism in $\mathcal{Fin}_*$ such that $\pi(g) = \omega \circ \pi(f)$. Due to the fact of $\pi(f)$ being a bijection on the restriction, we observe that ω is determined by $\pi(g)$. It follows that

$$\text{Mul}(\{X'_j\}_{j \in \omega^{-1}(k)}, X''_k) = \text{Mul}(\{X_{a_j}\}_{j \in \omega^{-1}(k)}, X''_k) = \text{Mul}(\{X_i\}_{i \in \pi(g)^{-1}(k)}, X''_k)$$

and we can define for all $k \in \langle l \rangle$ $h_k = g_k$. So, we have constructed a morphism

$$h: \{X'_j\}_{j \in \langle m \rangle} \rightarrow \{X''_k\}_{k \in \langle l \rangle}$$

such that $h \circ f = g$ and $\pi(h) = \omega$. The underlying category of $\mathcal{O}$ is the fiber category $\mathcal{O}_{\langle 1 \rangle}$ as the objects are 1-collections of colors and a morphism from X to Y is defined by $\text{Id}_{\langle 1 \rangle}$ with $\phi \in \text{Mul}_{\mathcal{O}}(\{X\}, Y)$. Furthermore, the equivalence $\mathcal{O}_{\langle n \rangle} \cong \mathcal{O}^n$ holds as the weaker op-fibration condition (M1) property still suffices to admit the proof given in proposition 2.19.

Further, let $\{X_i\}_{i \in \langle n \rangle}$ be a collection of colors, $\vec{X}$ denoting the corresponding object in $\mathcal{O}_{\langle n \rangle}^\otimes$ and $Y \in \mathcal{O}$. The set $\text{Mul}_{\mathcal{O}}(\{X_i\}_{i \in \langle n \rangle}, Y)$ can be identified with the set of morphisms $f: \vec{X} \rightarrow Y$ in $\mathcal{O}^\otimes$ such that $\pi(f)$ being the morphism with $\pi(f)^{-1}(*) = \{*\}$. In addition, the composition law for morphisms in the colored operad $\mathcal{O}$ can be reconstructed from the composition law in $\mathcal{O}^\otimes$.

Remark 2.29. Let $\mathcal{P}$ be a colored planar operad. Then analogously there is also a forgetful functor

$$\pi: \mathcal{P}^\otimes \rightarrow \Delta_*.$$

which satisfies the weaker op fibration condition (M1) and Segal condition (M2) property.

This gives us an alternative definition of a colored operad. Instead of seeing a colored operad as a categorical structure $\mathcal{O}$ equipped with an extensive notion of morphisms, we can now view colored operads as a category $\mathcal{O}^\otimes$ equipped with a forgetful functor $\pi: \mathcal{O}^\otimes \rightarrow \mathcal{Fin}_*$ satisfying conditions such as $\mathcal{O}_{\langle n \rangle}^\otimes \cong (\mathcal{O}_{\langle 1 \rangle}^\otimes)^n$ and $\mathcal{O} = \mathcal{O}_{\langle 1 \rangle}^\otimes$ inherit the structure of a colored operad. Even though the conditions π has to satisfy seem tedious, the second definition is phrased entirely in the language of category theory and can therefore be generalized to an ∞ -categorical notion of a colored operad.

2. Operads

Definition 2.30. A morphism $f: \langle n \rangle \rightarrow \langle m \rangle$ is called *inert* if for every $i \in \langle m \rangle^\circ$ the inverse image $f^{-1}(i)$ has exactly one element.

Definition 2.31. An ∞ -operad is a functor $p: \mathcal{O}^\otimes \rightarrow \mathcal{N}(\mathcal{F}\text{in}_*)$ between ∞ -categories satisfying the following conditions:

- (i) For every inert morphism $f: \langle n \rangle \rightarrow \langle m \rangle$ in $\mathcal{N}(\mathcal{F}\text{in}_*)$ and object $C \in \mathcal{O}_{\langle m \rangle}^\otimes$ there exists a p -cocartesian morphism $f: C \rightarrow C'$ in $\mathcal{O}^\otimes$ lifting f . In particular f induces a functor $f_!: \mathcal{O}_{\langle n \rangle}^\otimes \rightarrow \mathcal{O}_{\langle m \rangle}^\otimes$.
- (ii) Let $C \in \mathcal{O}_{\langle m \rangle}^\otimes$ and $C' \in \mathcal{O}_{\langle n \rangle}^\otimes$ be objects, $f: \langle m \rangle \rightarrow \langle n \rangle$ be a morphism in $\mathcal{F}\text{in}_*$ and let $\text{Map}_{\mathcal{O}^\otimes}^f(C, C')$ be the union of connected components of $\text{Map}_{\mathcal{O}^\otimes}(C, C')$ which lie over f . Choose p -cocartesian morphisms $C' \rightarrow C'_i$ lying over the inert morphism $\rho^i: \langle n \rangle \rightarrow \langle 1 \rangle$ for $1 \leq i \leq n$. Then the induced map

$$\text{Map}_{\mathcal{O}^\otimes}^f(C, C') \rightarrow \prod_{1 \leq i \leq n} \text{Map}_{\mathcal{O}^\otimes}^{\rho^i \circ f}(C, C'_i)$$

is a homotopy equivalence

- (iii) For every finite collection of objects $C_1, \dots, C_n \in \mathcal{O}_{\langle 1 \rangle}^\otimes$, there exists an object $C \in \mathcal{O}_{\langle n \rangle}^\otimes$ and a collection of p -cocartesian morphism $C \rightarrow C_i$ covering $\rho^i: \langle n \rangle \rightarrow \langle 1 \rangle$.

3 Braided Operads

Braided operads were first mentioned by Fiedorowicz in his paper [ZF] on the construction of E_n operads. Let A be an operad in the category of compactly generated Hausdorff spaces. The first theorem states:

Theorem 3.1 (Recognition principles for E_n operads). *A is an E_2 operad iff $A(k)$ is connected and the collection of universal covering spaces $\{\tilde{A}(k)\}_{k \in \mathbb{N}}$ forms a B_∞ operad, i.e.:*

- (1) *For all $k \in \mathbb{N}$, the space $\tilde{A}(k)$ is contractible.*
- (2) *The braid group B_k acts freely on $\tilde{A}(k)$ for each $k \in \mathbb{N}$.*

In the sketch of the proof, it is stated that "the definition of operads can be reformulated using actions by braid groups in place of actions by symmetric groups" [ZF]. As braided operads are a useful tool, e.g. recognizing E_n operads, we want to gain a deeper understanding of braided operads. Further, we have already seen that colored operads correspond to op-fibrations over finite pointed sets Fin_* as well as colored planar operads correspond to op-fibrations over Δ_* . We are hoping to find a suitable category such that colored braided operads correspond to op-fibrations over this category.

It might be interesting to try to define braided operads in braided monoidal categories rather than symmetric monoidal categories potentially yielding interesting examples. However, it is not straight forward how to do this as shuffling objects would not be a canonical isomorphism. So for now, we will stay in a symmetric monoidal category $\mathcal{C}$.

Definition 3.2. A *braided operad* $\mathcal{O}$ in $\mathcal{C}$ is a planar operad as defined in definition 2.1 with an action of B_n on $\mathcal{O}(n)$ for all $n \in \mathbb{N}$ together with the following equivariance condition: For $\sigma \in B_k$, let $\sigma(j_1, \dots, j_k) \in B_j$ denote the element that σ -braids k blocks with each block having j_l strands for $1 \leq l \leq k$. Then the following diagram commutes:

$$\begin{array}{ccc} \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) & \xrightarrow{\sigma \otimes \bar{\sigma}^{-1}} & \mathcal{O}(k) \otimes \mathcal{O}(j_{\sigma(1)}) \otimes \dots \otimes \mathcal{O}(j_{\sigma(k)}) \\ \gamma \downarrow & & \downarrow \gamma \\ \mathcal{O}(j) & \xrightarrow{\sigma(j_1 \dots j_k)} & \mathcal{O}(j), \end{array}$$

where $\bar{\sigma}$ denotes the underlying permutation of σ . Furthermore, let $\tau_l \in B_{j_l}$ for $1 \leq l \leq k$ and $\tau_1 \oplus \dots \oplus \tau_k$ denotes the sum of braidings on k blocks with each block having j_l strands. Then the following diagram commutes

$$\begin{array}{ccc} \mathcal{O}(k) \otimes \mathcal{O}(j_1) \otimes \dots \otimes \mathcal{O}(j_k) & \xrightarrow{\text{Id} \otimes \tau_1 \otimes \dots \otimes \tau_k} & \mathcal{O}(k) \otimes \mathcal{O}(j_{\sigma(1)}) \otimes \dots \otimes \mathcal{O}(j_{\sigma(k)}) \\ \gamma \downarrow & & \downarrow \gamma \\ \mathcal{O}(j) & \xrightarrow{\tau_1 \oplus \dots \oplus \tau_k} & \mathcal{O}(j) \end{array}$$

Remark 3.3. Similarly to symmetric operads, we can define algebras and modules over braided operads by changing the equivariance condition:

Let $\mathcal{O}$ be a braided operad and A an algebra over the underlying planar operad of $\mathcal{O}$. For all $j \in \mathbb{N}$, $\sigma \in B_j$ and $\bar{\sigma} \in S_j$ being the underlying permutation of σ :

$$\begin{array}{ccc} \mathcal{O}(j) \otimes A^j & \xrightarrow{\sigma \otimes \bar{\sigma}^{-1}} & \mathcal{O}(j) \otimes A^j \\ & \searrow \theta & \swarrow \theta \\ & A & \end{array}$$

Then, we call A an algebra over the braided operad $\mathcal{O}$.

Further, let A be an algebra over the braided operad $\mathcal{O}$ and M an object in $\mathcal{C}$ with

$$\omega: \mathcal{O}(j) \otimes A^{j-1} \otimes M \rightarrow M$$

3. Braided Operads

satisfying associativity and unitality conditions from definition 2.10. Then we call M an A -module if the following equivariance condition holds for all $j \in \mathbb{N}$, $\sigma \in B_{j-1} \subset B_j$ and $\bar{\sigma} \in S_{j-1}$ being the underlying permutation of σ :

$$\begin{array}{ccc} O(j) \otimes A^{j-1} \otimes M & \xrightarrow{\sigma \otimes \bar{\sigma}^{-1} \otimes \text{Id}} & O(j) \otimes A^{j-1} \otimes M \\ & \searrow \omega & \swarrow \omega \\ & M & \end{array}$$

- Example 3.4.** 1. Again, let us consider the operad in **Set** defined by the singleton set in each degree $n \in \mathbb{N}$. The braid group acts trivially. When we consider algebras over this operad we will again obtain commutative algebras.
2. We have already encountered an example of a planar operad in example 2.6 which can be lifted to a braided operad—the braid operad $\mathcal{B}$. The braid group B_n acts by composition. This group action is equivariant.

3.1 Action Operad

For this section, we remain in the symmetric monoidal category of sets equipped with the cartesian symmetric monoidal structure. We will introduce an action operad that acts on planar operads and this relates to braided operads following [DY]. An idea will be that the theory on braided operads can be extended in a sense to group action operads.

move to-
wards the
outlook?

Definition 3.5. An *action operad* is a tuple

$$(\{G(n)\}_{n \in \mathbb{N}}, \gamma^G, 1^G, \omega)$$

satisfying the following conditions:

- (i) For all $n \in \mathbb{N}$, $G(n)$ is a group with unit Id_n .
- (ii) There is a group homomorphism $\omega: G(n) \rightarrow S_n$ for all n which we call augmentation. For each $g \in G(n)$ we call $\omega(g)$ the underlying permutation of g .
- (iii) The tuple $(\{G(n)\}_{n \in \mathbb{N}}, \gamma^G, 1^G)$ is a one-colored planar operad in **Set** which we denote by G . It fulfills the following conditions:

- (a) The augmentation

$$\omega: G \rightarrow \mathcal{S}$$

is a morphism of planar operads in **Set** where $\mathcal{S}$ denotes the symmetry operad from examples 2.12.

- (b) For $n \geq 1$, let $\sigma, \rho \in G(n)$ and for $k_j \in \mathbb{N}$, $1 \leq j \leq n$ let $\tau_j, \tau'_j \in G(k_j)$. Then the following equation holds in $G(k)$ for $\sum_{1 \leq j \leq n} k_j = k$:

$$\gamma^G(\sigma \rho, \tau_1 \tau'_1, \dots, \tau_n \tau'_n) = \gamma^G(\sigma, \tau_{\bar{\rho}^{-1}(1)}, \dots, \tau_{\bar{\rho}^{-1}(n)}) \cdot \gamma^G(\rho, \tau'_1, \dots, \tau'_n)$$

where $\bar{\rho} = \omega(\rho) \in S_n$.

We call the condition (iii)(b) the action operad equation.

- Examples 3.6.** 1. The *planar group operad* P is the action operad where $P(n) = \{*\}$ is the trivial group for all $n \in \mathbb{N}$.
2. The symmetry operad $\mathcal{S}$ from examples 2.12 equipped with the identity augmentation forms an action operad.
3. Another example which we have already seen is the braid operad which we equip with the projection to the symmetric group as the augmentation.

3. Braided Operads

Proposition 3.7. *The braid operad $\mathcal{B}$ from example 2.6 equipped with the projection*

$$\pi: \mathcal{B} \rightarrow \mathcal{S}$$

forms an action operad

$$B = (\{B_n\}_{n \in \mathbb{N}}, \gamma, \text{Id}_1 \in \mathcal{B}_1, \pi)$$

Proof. We have already seen that $\mathcal{B}$ forms a one-colored planar operad in **Set**. Hence, it remains to show that π is a morphism of planar operads. We note that $\pi(\text{Id}_1)$ is the trivial permutation and π preserves the operadic composition as it preserves the group multiplication. The action operad equation follows from remark 2.5. $\square$

3. Braided Operads

3.2 Universal Covering of E_2

In this section we want to establish an operadic structure on the universal covering of E_2 . First, we need to define what it means to be a universal covering of a topological operad.

Definition 3.8. Let $\mathcal{P}$ be a topological operad. A covering operad $\tilde{\mathcal{P}}$ is a topological operad such that there is a morphism of operads $c: \tilde{\mathcal{P}} \rightarrow \mathcal{P}$ where $c_n: \tilde{\mathcal{P}}(n) \rightarrow \mathcal{P}(n)$ is a covering of $\mathcal{P}(n)$ for all $n \geq 1$.

Definition 3.9. Let $\mathcal{P}$ be a topological operad. A universal covering operad $\tilde{\mathcal{P}}$ is a covering operad that is universal in every degree $n \geq 1$.

While the operadic structure requires labels in the sense that they are necessary to define the composition maps in E_2 , they do not matter from a topological perspective. Hence, in this section we want to establish a universal covering of E_2 without labels. We will then see in the next step that this universal covering is homeomorphic to the universal covering of E_2 . So, instead of E_2 we will consider the quotient:

$$E_2 \rightarrow E_2 / S$$


We identify configurations that only differ in their labels, i.e. we consider the quotient of E_2 by the action of the symmetric group S on the labels. So, in $E_2^\otimes$ we identify $f, g: \langle n \rangle \rightarrow \langle m \rangle$ if there exists a commuting diagram in $E_2^\otimes$

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{f} & \langle m \rangle \\ \downarrow \sigma & & \downarrow \tau \\ \langle n \rangle & \xrightarrow{g} & \langle m \rangle \end{array}$$

context of
why and
how we can
consider
things in
 $E_2^\otimes$

where $\sigma \in S_n$ and $\tau \in S_m$.

Before, we can find a universal covering of E_2 / S , we need to choose basepoints in each configuration space. So, for all $n \geq 1$ we choose the basepoint $\bar{b}_n \in E_2(n)/S_n$ to be the following configuration:



Note that since $E_2(n)$ is connected and simply connected then so is $E_2(n)/S_n$. Hence, it has a universal covering $\widetilde{E_2(n)/S_n}$ which we define in the usual sense:

$$\widetilde{E_2(n)/S_n} := \{\gamma: [0, 1] \rightarrow E_2(n)/S_n \mid \gamma(0) = \bar{b}_n\} / \text{homotopy with fixed endpoints}$$

with projection map:


$$\begin{aligned} p: \widetilde{E_2(n)/S_n} &\rightarrow E_2(n)/S_n \\ \gamma &\mapsto \gamma(1) \end{aligned}$$

It is known that this in fact defines a universal covering of $E_2(n)/S_n$. Analogously, there is a universal covering of E_2 with labels, when we choose the basepoint $b_n \in E_2(n)$:

reference

add univer-
sal covern
with labels

3. Braided Operads



and define its universal covering analogously:

$$\widetilde{E_2(n)} := \{\gamma: [0, 1] \rightarrow E_2(n) \mid \gamma(0) = b_n\} / \text{homotopy with fixed endpoints}$$

with projection map:

$$\begin{aligned} p: \widetilde{E_2(n)} / S_n &\rightarrow E_2(n) / S_n \\ \gamma &\mapsto \gamma(1) \end{aligned}$$

The labels have no purpose from a topological perspective, so it should not come as a surprise that there is a homeomorphism

$$\begin{aligned} \widetilde{E_2} &\cong \widetilde{E_2} / S \\ (\gamma: b_n \rightarrow c) &\mapsto (\bar{\gamma}: \bar{b}_n \rightarrow \bar{c}) \\ (\gamma: b_n \rightarrow c') &\mapsto (\bar{\gamma}: \bar{b}_n \rightarrow \bar{c}) \end{aligned}$$

where we forget labels in one directions and in the reverse direction we label the basepoint $\bar{b}_n$ from left to right. The labeling of c' is then induced by the path γ . clearly this is continuous in both directions and inverse to one another. This shows that topologically the path γ does not need the labels which is summarized in the following representation. We observe that for every $\gamma \in \widetilde{E_2(n)} / S_n$ there is a commuting diagram in $E_2^\otimes$ for some $\sigma \in S_n$:

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{c_n} & \langle 1 \rangle \\ \sigma \downarrow & & \downarrow \gamma \\ \langle n \rangle & \xrightarrow{b_n} & \langle 1 \rangle \end{array}$$

Essentially, $\sigma \in S_n$ fixes the labels such that we can consider γ as a path in $\widetilde{E_2}$. We will use this representation to define an operadic structure on $\widetilde{E_2} / S$ such that the covering is a morphism of operads and then also use it to define a braid action on the universal covering.

Proposition 3.10. *There is an operadic structure on $\widetilde{E_2}$ such that the covering p is a morphism of operads.*

Proof. Let

$$\gamma: E_2(k) \times E_2(j_1) \times \cdots \times E_2(j_k) \rightarrow E_2(j)$$

be a composition map in E_2 . Then, we define a composition map in $\widetilde{E_2}$

$$\tilde{\gamma}: \widetilde{E_2}(k) \times \widetilde{E_2}(j_1) \times \cdots \times \widetilde{E_2}(j_k) \rightarrow \widetilde{E_2}(j),$$

which we construct as follows: Let $(\bar{\alpha}, \alpha_1, \dots, \alpha_k) \in \widetilde{E_2}(k) \times \widetilde{E_2}(j_1) \times \cdots \times \widetilde{E_2}(j_k)$ with

$$\alpha_l: b_{j_l} \rightarrow c_l$$

3. Braided Operads

for $1 \leq l \leq k$ and

$$\bar{\alpha}: b_k \rightarrow c_k.$$

Then

$$\bar{\alpha} \circ (\alpha_1, \dots, \alpha_k)$$

denotes the path from $\gamma((b_k, b_{j_1}, \dots, b_{j_k}))$ to $\gamma(c_k, c_{j_1}, \dots, c_{j_k})$ where we first apply

$$\alpha_l: b_{j_l} \rightarrow c_l$$

on the configuration in the l -th disk of b_k for all $1 \leq l \leq k$ and then apply

$$\bar{\alpha}: b_k \rightarrow c_k$$

on the configuration b_k .




Figure 1: example of $\bar{\alpha} \circ (\alpha_1, \dots, \alpha_3)$ for $\bar{\alpha} \in \widetilde{E}_2(3)$ and $\alpha_1, \alpha_2, \alpha_3 \in \widetilde{E}_2(2)$

Further, let $\tau \in \widetilde{E}_1(j) \subset \widetilde{E}_2(j)$ be a path from b_j to $\gamma((b_k, b_{j_1}, \dots, b_{j_k}))$ that resizes and translates the disks along the horizontal axis. This path τ is unique up to homotopy. This generates the following commuting diagram in $E_2^\otimes$:

$$\begin{array}{ccc} \langle j \rangle & \xrightarrow{\gamma(c_k, c_{j_1}, \dots, c_{j_k})} & \langle 1 \rangle \\ S_j \Downarrow & & \uparrow \bar{\alpha} \circ (\alpha_1, \dots, \alpha_k) \\ \langle j \rangle & \xrightarrow{\gamma(b_k, b_{j_1}, \dots, b_{j_k})} & \langle 1 \rangle \\ S_j \Downarrow & & \uparrow \tau \\ \langle j \rangle & \xrightarrow{b_j} & \langle 1 \rangle \end{array}$$


Therefore, we can set

$$\tilde{\gamma}((\bar{\alpha}, \alpha_1, \dots, \alpha_k)) := \bar{\alpha} \circ (\alpha_1, \dots, \alpha_k) \circ \tau$$

We observe, that this composition map is well-defined, since the path τ is unique up to homotopy. To see that $\tilde{\gamma}$ is associative we need to have a look at the definition. Note, that in the definition $\tau \in E_1(j)$ is a path that resizes and translates the disks, so it has no effect on the paths from a homotopy perspective. Further, we know that composition of paths is associative upto homotopy. So, for

$$(\alpha, (\alpha_s)_{1 \leq s \leq k}, (\alpha_r)_{1 \leq r \leq j}) \in E_2(k) \times \prod_{s=1}^k E_2(j_s) \times \prod_{r=1}^j E_2(i_r)$$

3. Braided Operads



Then, we obtain a braid action on $\widetilde{\mathbb{E}}_2$. Let $\alpha: b_n \rightarrow c$ be in $\widetilde{\mathbb{E}}_2/S$ which is represented by the following diagram in $\mathbb{E}_2^\otimes$ for some $\tau \in S_n$:

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{c} & \langle 1 \rangle \\ \tau \downarrow & & \uparrow \alpha \\ \langle n \rangle & \xrightarrow{b_n} & \langle 1 \rangle \end{array}$$

and let $\sigma \in B_n$, then we set:

$$\sigma.\alpha := \alpha \circ \sigma^{-1}$$

Concatinating these paths results in the following commuting diagram, where $\bar{\sigma} \in S_n$ refers to the underlying permutation of σ .

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{c} & \langle 1 \rangle \\ \tau \downarrow & & \uparrow \alpha \\ \langle n \rangle & \xrightarrow{b_n} & \langle 1 \rangle \\ \bar{\sigma} \downarrow & & \uparrow \sigma^{-1} \\ \langle n \rangle & \xrightarrow{b_n} & \langle 1 \rangle \end{array}$$

What is left to show is that this braid action is equivariant with respect to the operadic structure on $\widetilde{\mathbb{E}}_2$. Note that by definition of $\widetilde{\gamma}$ we move disks in a large disk D from the basepoint configuration to another configuration to another configuration of D again afterwards but only move D as a part of another configuration. Hence, in that sense the order in which we apply paths to a configuration does not matter from a homotopic perspective. This fact is analogous to the fact that we can pull inside braids through outside braids, i.e. the second relation of braids in remark 2.4. This means that for all $\sigma \in B_k$ we have the following commutative diagram:

$$\begin{array}{ccc} \widetilde{\mathbb{E}}_2(k) \times \widetilde{\mathbb{E}}_2(j_1) \times \cdots \times \widetilde{\mathbb{E}}_2(j_k) & \xrightarrow{\sigma \times \bar{\sigma}} & \widetilde{\mathbb{E}}_2(k) \times \widetilde{\mathbb{E}}_2(j_{\bar{\sigma}(1)}) \times \cdots \times \widetilde{\mathbb{E}}_2(j_{\bar{\sigma}(k)}) \\ \downarrow \widetilde{\gamma} & & \downarrow \widetilde{\gamma} \\ \widetilde{\mathbb{E}}_2(j) & \xrightarrow{\sigma(j_1, \dots, j_k)} & \widetilde{\mathbb{E}}_2(j) \end{array}$$

picture of paths of disks that correspond to the braid remark

and for $\tau_i \in B_{j_i}$ with $1 \leq i \leq k$, we have the following commutative diagram:

$$\begin{array}{ccc} \widetilde{\mathbb{E}}_2(k) \times \widetilde{\mathbb{E}}_2(j_1) \times \cdots \times \widetilde{\mathbb{E}}_2(j_k) & \xrightarrow{(\text{Id}, \tau_1, \dots, \tau_k)} & \widetilde{\mathbb{E}}_2(k) \times \widetilde{\mathbb{E}}_2(j_1) \times \cdots \times \widetilde{\mathbb{E}}_2(j_k) \\ \downarrow \widetilde{\gamma} & & \downarrow \widetilde{\gamma} \\ \widetilde{\mathbb{E}}_2(j) & \xrightarrow{\tau_1 \oplus \cdots \oplus \tau_k} & \widetilde{\mathbb{E}}_2(j) \end{array}$$

Hence, the braid group action on $\widetilde{\mathbb{E}}_2$ respects to the operadic structure.

4 Colored Braided Operads

In this section, we will define colored braided operads and will aim to find a op-fibration that encodes all the necessary data to give an alternative definition. First, let us recall the definition of $E_2^\otimes$ and build a related category $\mathcal{Br}_*$ that will serve as the target category for our op-fibration later on.


Remark 4.1. For each $n \in \mathbb{N}$ we have an object $\langle n \rangle$. A morphism $\langle n \rangle \rightarrow \langle m \rangle$ in $E_2^\otimes$ consists of a morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in $\mathcal{Fin}_*$ and a collection of configurations

$$\{c_j \in E_2(|\alpha^{-1}(j)|)\}_{1 \leq j \leq m}$$

where each configuration is labeled by $\alpha^{-1}(j)$. So for example,

$$\begin{aligned} \alpha: \langle 3 \rangle &\rightarrow \langle 2 \rangle \\ 1 &\mapsto 1 \\ 2 &\mapsto 2 \\ 3 &\mapsto 1 \end{aligned}$$

together with the configurations:



We denote the trivial configuration with label x by $\text{Id}^x \in E_2(1)$. Further, recall the composition in $E_2^\otimes$. Let

$$\alpha: \langle n \rangle \rightarrow \langle m \rangle \text{ with configurations } \{c_j\}_{1 \leq j \leq m}$$

and

$$\alpha': \langle m \rangle \rightarrow \langle l \rangle \text{ with configurations } \{c'_k\}_{1 \leq k \leq l}.$$

Then, the configuration c'_k is labeled by $\alpha'^{-1} = \{j_1^{(k)}, \dots, j_p^{(k)}\}$. So, we obtain a configuration

$$\gamma(c'_k, c_{j_1}, \dots, c_{j_p})$$

labeled by $(\alpha' \circ \alpha)^{-1}(k)$ for all $1 \leq k \leq l$ giving us the well-defined composition $\alpha' \circ \alpha$

Definition 4.2. A morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in $E_2^\otimes$ is called active if it is active in $\mathcal{Fin}_*$, i.e. $\alpha^{-1}(*) = \{*\}$.

Next, we partially enrich $E_2^\otimes$ with a groupoid structure.

Definition 4.3. Let $\mathcal{Br}_*$ be the following partially **Grpd**-enriched category:

- (i) The objects of $\mathcal{Br}_*$ are finite sets $\langle n \rangle$ for $n \in \mathbb{N}$.

define active morphism definition here?

define BR category. How to define objects?

4. Colored Braided Operads

- (ii) A morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in $\mathcal{Br}_*$ consists of a morphism $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in $\mathcal{Fin}_*$ and a collection of configurations

$$\{\tilde{\alpha}_j \in E_2(|\alpha^{-1}(j)|)\}_{1 \leq j \leq m}$$

where each configuration $\tilde{\alpha}_j$ is labeled by $\alpha^{-1}(j)$. The composition is defined analogously to the composition map in $E_2^\otimes$ as defined above. Further, the identity morphism $\text{Id}_{\langle n \rangle}$ is given by the collection of trivial configurations $\{\text{Id}^j \in E_2(1)\}_{1 \leq j \leq n}$.

- (iii) The active morphisms into $\langle 1 \rangle$

$$\text{Hom}^{\text{active}}(\langle n \rangle, \langle 1 \rangle)$$

form the groupoid of homotopy classes of paths in the configuration space $E_2(n)$. The remaining hom-categories are discrete groupoids.

It is clear that this defines a category. We can extend the definition of $\mathcal{Br}_*$ to include any finite set I . Note, that that category would be equivalent to $\mathcal{Br}_*$ in the analogous way as $\mathcal{Fin}_*$ is equivalent to the category of finite sets with a distinguished point. We are now set up to understand the following definition:

Definition 4.4. A *colored braided operad* $\mathcal{B}$ consists of the following data:

- (i) A collection $\{X, Y, Z; \dots\}$ of colors.
(ii) Let I be a finite set and $\tilde{I}$ an I -labeled configuration in E_2 . For a collection of colors $\{X_i\}_{i \in I}$ and a color Y in $\mathcal{B}$, there is a set

$$\text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Y)$$

called the set of morphisms from $\{X_i\}_{i \in I}$ to Y .

- (iii) Let $\alpha: I \rightarrow J$ be finite maps together with a collection of configurations labeled by the preimages of α , i.e.

$$\tilde{\alpha} = (\tilde{I}_j)_{j \in J}$$

where I_j denotes the fiber over j for all $j \in J$ and a path of configurations η :

$$\begin{array}{ccc} I & \xrightarrow{\tilde{\alpha}} & J \\ \tilde{I} \searrow & \Downarrow \eta & \swarrow \tilde{J} \\ & \langle 1 \rangle & \end{array}$$

Further, let there be collections of colors $\{X_i\}_{i \in I}$ and $\{Y_j\}_{j \in J}$ and color Z . Then, there is a composition map:

$$\gamma: \prod_{j \in J} \text{Mul}_{\mathcal{B}}^{\tilde{I}_j}(\{X_i\}_{i \in I_j}, Y_j) \times \text{Mul}_{\mathcal{B}}^{\tilde{J}}(\{Y_j\}_{j \in J}, Z) \rightarrow \text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Z)$$

- (iv) There is a collection of morphisms $\{\text{Id}_X \in \text{Mul}_{\mathcal{B}}^{\text{Id}^1}(\{X\}, X)\}_{X \in \mathcal{B}}$ being a left and right unit in the composition, i.e. the following diagrams commute:

$$\begin{array}{ccc} \text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Y) & \xrightarrow{\cong} & \text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Y) \times \{\text{Id}_Y\} \\ \downarrow \text{Id} & & \downarrow \\ \text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Y) & \xleftarrow{\gamma} & \text{Mul}_{\mathcal{B}}^{\tilde{I}}(\{X_i\}_{i \in I}, Y) \times \text{Mul}_{\mathcal{B}}^{\text{Id}^1}((Y), Y) \end{array}$$

4. Colored Braided Operads

$$\begin{array}{ccc}
 \text{Mul}_{\mathcal{B}}^{\tilde{I}}(X_I, Y) & \xrightarrow{\cong} & (\prod_{i \in I} \{\text{Id}_{X_i}\}) \times \text{Mul}_{\mathcal{B}}^{\tilde{I}}(X_I, Y) \\
 \downarrow \text{Id} & & \downarrow \\
 \text{Mul}_{\mathcal{B}}^{\tilde{I}}(X_I, Y) & \xleftarrow{\gamma} & (\prod_{i \in I} \text{Mul}_{\mathcal{B}}^{\text{Id}^i}((X_i), X_i)) \times \text{Mul}_{\mathcal{B}}^{\tilde{I}}(X_I, Y)
 \end{array}$$

(v) For every sequence of finite maps $I \xrightarrow{\alpha} J \xrightarrow{\beta} K$, with paths

$$\begin{array}{ccccc}
 I & \xrightarrow{\tilde{\alpha}} & J & \xrightarrow{\tilde{\beta}} & K \\
 & \searrow \tilde{I} & \downarrow \tilde{J} & \swarrow \tilde{K} & \\
 & & \langle 1 \rangle & &
 \end{array}$$

and collections of colors $X_I = \{X_i\}_{i \in I}$, $Y_J = \{Y_j\}_{j \in J}$, $W_K = \{W_k\}_{k \in K}$ and a color $Z \in \mathcal{B}$ the following diagram commutes:

$$\begin{array}{ccc}
 \prod_{j \in J} \text{Mul}_{\mathcal{B}}^{\tilde{I}_j}(X_{I_j}, Y_j) \times \prod_{k \in K} \text{Mul}_{\mathcal{B}}^{\tilde{J}}(Y_J, W_k) \times \text{Mul}_{\mathcal{B}}^{\tilde{K}}(W_K, Z) & \xrightarrow{\text{Id} \times \gamma} & \prod_{j \in J} \text{Mul}_{\mathcal{B}}^{\tilde{I}_j}(X_{I_j}, Y_j) \times \text{Mul}_{\mathcal{B}}^{\tilde{J}}(Y_J, Z) \\
 \downarrow (\gamma \circ s) \times \text{Id} & & \downarrow \\
 \prod_{k \in K} \text{Mul}_{\mathcal{B}}^{\tilde{I}_k}(X_{I_k}, W_k) \times \text{Mul}_{\mathcal{B}}^{\tilde{K}}(W_K, Z) & \xrightarrow{\quad} & \text{Mul}_{\mathcal{B}}(X_I, Z)
 \end{array}$$

make diagram prettier

Example 4.5. Let $\mathcal{B}_{col}$ be a colored braided operad with one color X . Similar to the symmetric case, $\mathcal{B}_{col}$ induces a braided operad $\mathcal{B}$ as in definition 3.2. For $n \in \mathbb{N}$, we set

$$\mathcal{B}(n) = \text{Mul}_{\mathcal{B}_{col}}^{b_n}(\{X\}_{1 \leq i \leq n}, X)$$

where b_n is the basepoint configuration in $E_2(n)$. Further, we define the composition map

$$\theta: \mathcal{B}(k) \times \mathcal{B}(j_1) \times \cdots \times \mathcal{B}(j_k) \rightarrow \mathcal{B}(j)$$

where $j = j_1 + \cdots + j_k$, from the composition map γ in $\mathcal{B}_{col}$:

$$\theta: \prod_{1 \leq i \leq k} \text{Mul}_{\mathcal{B}_{col}}^{b_{j_i}}(\{X_i\}^{j_i}, X) \times \text{Mul}_{\mathcal{B}_{col}}^{b_k}(\{X\}^k, X) \xrightarrow{\gamma} \text{Mul}_{\mathcal{B}_{col}}^{b_k(b_{j_1}, \dots, b_{j_k})}(\{X\}^j, X) \cong \text{Mul}_{\mathcal{B}_{col}}^{b_j}(\{X\}^j, X)$$

where the postcomposition is the bijection induced by morphism There is a unit

$$\eta: * \rightarrow \mathcal{B}(1)$$

which is defined by the identity morphism $\text{Id}_X \in \text{Mul}_{\mathcal{B}_{col}}^{b_1}(\{X\}, X)$. The unitality condition is satisfied as the identity morphism Id_X is the left and right unit in $\text{Mul}_{\mathcal{B}_{col}}^{b_1}(\{X\}, X)$. The associativity condition as it holds for $\mathcal{B}_{col}$: For any choice of integers

$$n, j_1, \dots, j_k, i_1, \dots, i_j$$

??? how to handle E_1 translation like we need it here?

4. Colored Braided Operads

with

$$\sum_{1 \leq s \leq k} j_s = j, \quad \sum_{1 \leq r \leq j} i_r = i, \quad g_s = \sum_{0 \leq l \leq s} j_l \text{ and } h_s = \sum_{1 \leq q \leq j_s} i_{g_{s-1}+q}$$

the following diagram

$$\begin{array}{ccc} \mathcal{B}(k) \times (\prod_{s=1}^k \mathcal{B}(j_s)) \times (\prod_{r=1}^j \mathcal{B}(i_r)) & \xrightarrow{\gamma \times \text{Id}} & \mathcal{B}(j) \times (\prod_{r=1}^j \mathcal{B}(i_r)) \\ \downarrow s & & \downarrow \gamma \\ \mathcal{B}(k) \times \prod_{s=1}^k (\mathcal{B}(j_s) \times \prod_{q=1}^{j_s} \mathcal{B}(i_{g_{s-1}+q})) & \xrightarrow{\text{Id} \times (\times_s \gamma)} & \mathcal{B}(k) \times (\prod_{s=1}^k \mathcal{B}(h_s)) \\ & & \uparrow \gamma \\ & & \mathcal{B}(i) \end{array}$$

where s shuffles the entries of the tensor product, commutes as the composition map γ in $\mathcal{B}_{col}$ is associative for the following maps for $1 \leq s \leq k$, $1 \leq q \leq j_s$ and $h'_q = \sum_{t=1}^{g_{s-1}+1} i_t$.

$$\begin{aligned} \alpha: \langle i \rangle &\rightarrow \langle j \rangle \\ h'_{q-1} + 1, \dots, h'_q &\mapsto g_{s-1} + q \end{aligned}$$

with $\widetilde{\alpha_{g_{s-1}+q}} = b_{i_{g_{s-1}+q}}$ and

$$\begin{aligned} \alpha': \langle j \rangle &\rightarrow \langle k \rangle \\ g_{s-1} + 1, \dots, g_s &\mapsto s \end{aligned}$$

with $\widetilde{\alpha'_s} = b_{j_s}$. So we have defined a planar operad $\mathcal{B}$. It remains to define a well-defined action of the braid group B_n on $\mathcal{B}(n)$ and verify that it respects θ .

Definition 4.6. Let $\mathcal{B}$ be a colored operad. We define a category $\mathcal{B}^\otimes$ in the following way:

- (i) The objects of $\mathcal{B}^\otimes$ are finite sequences of colors of $\mathcal{B}$.
- (ii) Let $\{X_i\}_{1 \leq i \leq m}$ and $\{Y_j\}_{1 \leq j \leq n}$ be objects in $\mathcal{B}^\otimes$. A morphism

$$\phi: \{X_i\}_{1 \leq i \leq m} \rightarrow \{Y_j\}_{1 \leq j \leq n}$$

is given by a map $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ in $\mathcal{B}_*$ together with a family of morphisms

$$\{\phi_j \in \text{Mul}_{\mathcal{B}}^{\widetilde{\alpha_j}}(\{X_i\}_{i \in \alpha^{-1}(j)}, Y_j)\}_{1 \leq j \leq n}$$

in $\mathcal{B}$.

- (iii) Composition of morphisms is determined by the composition in $\mathcal{B}_*$ and $\mathcal{B}$, i.e. for morphisms $\phi: X \rightarrow Y$ and $\phi': Y \rightarrow Z$ in $\mathcal{B}^\otimes$ with $\alpha': \langle n \rangle \rightarrow \langle m \rangle$ and $\alpha'': \langle m \rangle \rightarrow \langle k \rangle$, we have $\alpha = \alpha'' \circ \alpha'$ and

$$\{(\phi' \circ \phi)_l \in \text{Mul}_{\mathcal{B}}^{\widetilde{\alpha_l}}(\{X_i\}_{i \in \alpha^{-1}(l)}, Z_l)\}_{1 \leq l \leq k}$$

where $(\phi' \circ \phi)_l$ is applying the composition map

$$\gamma: \prod_{j \in \alpha''^{-1}(l)} \text{Mul}_{\mathcal{B}}^{\widetilde{\alpha'_j}}(\{X_i\}_{i \in \alpha'^{-1}(j)}, Y_j) \times \text{Mul}_{\mathcal{B}}^{\widetilde{\alpha'_l}}(\{Y_i\}_{i \in \alpha''^{-1}(l)}, Z_l) \rightarrow \text{Mul}_{\mathcal{B}}^{\widetilde{\alpha_l}}(\{X_i\}_{i \in \alpha^{-1}(l)}, Z_l).$$

add braid group action: how to fix labeling and Id_X problem?

4. Colored Braided Operads

4.1 Braided Monoidal Structure

In the following we will construct a braided monoidal structure obtained from an op-fibration over $\mathcal{B}r_*$. To do so, we will need to choose paths of configurations in $\mathcal{B}r_*$ on various occasions. We will refrain from specifying the paths in detail but rather describe the paths enough to understand from which homotopy class we choose a path so we then see why certain compositions of paths are homotopy equivalent. Also, we do not care much about the size of the disks as we can expand and shrink them continuously as needed.

We will now follow the same approach as in the symmetric monoidal case in section 2, i.e. we will define a category $\mathcal{C}^\otimes$ and a functor $\pi: \mathcal{C}^\otimes \rightarrow \mathcal{B}r_*$ such that π is an op-fibration and the fibers of π are given by the categories $\mathcal{C}^n := \mathcal{C}_{\langle n \rangle}^\otimes$ for $n \geq 1$.

If not stated otherwise, a morphism $\langle n \rangle \rightarrow \langle 1 \rangle$ always refers to an active morphism in $\mathcal{B}r_*$ from now on for all $n \in \mathbb{N}$.

Definition 4.7. An $\mathcal{B}r_*$ -operad is a category $\mathcal{C}^\otimes$ with a functor

$$\mathcal{C}^\otimes \rightarrow \mathcal{B}r_*$$

satisfying

(M1) **Op-fibration condition:** The functor π is an op-fibration.

(M2) **Segal condition:** For $\mathcal{C} := \mathcal{C}_{\langle 1 \rangle}^\otimes$, we have $\mathcal{C}_{\langle n \rangle}^\otimes \cong \mathcal{C}^n$ for all $n \geq 1$.

So for the next part, let $\pi: \mathcal{C}^\otimes \rightarrow \mathcal{B}r_*$ be a $\mathcal{B}r_*$ -operad and we denote $\mathcal{C} := \mathcal{C}_{\langle 1 \rangle}^\otimes$. In the following section we aim to define a braided monoidal structure on $\mathcal{C}$. First, we introduce the following abuse of notation: For a 2-morphism η , i.e. a path between two configurations $c_1, c_2 \in E_2(n)$ in $\mathcal{B}r_*$, we would write the following diagram:

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{\sigma} & \langle n \rangle \\ & \searrow c_1 \quad \swarrow c_2 & \\ & \langle 1 \rangle & \end{array}$$

However, we will use a slightly different notation for the sake of readability. We will write the following diagram instead:

$$\begin{array}{ccc} \langle n \rangle & \xrightarrow{\sigma} & \langle n \rangle \\ \downarrow c_1 & & \downarrow c_2 \\ \langle 1 \rangle & \dashrightarrow \eta & \langle 1 \rangle \end{array}$$

Let β denote the following morphism in $\mathcal{B}r_*$


$$\begin{array}{l} \beta: \langle 2 \rangle \rightarrow \langle 2 \rangle \\ 1 \mapsto 2 \\ 2 \mapsto 1 \end{array}$$

with the trivial collection $(\text{Id}^2, \text{Id}^1)$. Further, let

$$\begin{array}{l} \mu: \langle 2 \rangle \rightarrow \langle 1 \rangle \\ 1 \mapsto 1 \\ 2 \mapsto 1 \end{array}$$

be the morphism with the basepoint configuration $b_2 \in E_2(2)$:

4. Colored Braided Operads




Since π satisfies the op-fibration condition (M1) and the Segal condition (M2), we have that for any $(A, B) \in \mathcal{C}^2 \cong \mathcal{C}_{\langle 2 \rangle}^{\otimes}$ there is a cocartesian lift $(A, B) \rightarrow A \otimes B$. The cocartesian property ensures that the choice is unique up to unique isomorphism. So, we obtain a well-defined functor

$$\otimes: \mathcal{C}^2 \rightarrow \mathcal{C}.$$

Further, we consider the following diagramm in $\mathcal{B}r_*$:

$$\begin{array}{ccc} \langle 2 \rangle & \xrightarrow{\beta} & \langle 2 \rangle \\ \downarrow \mu & & \downarrow \mu \\ \langle 1 \rangle & \dashrightarrow & \langle 1 \rangle \end{array}$$

Filling the diagram, i.e. providing commutativity data means choosing a path from the basepoint configuration, corresponding to μ , to the twisted basepoint configuration, corresponding to $\mu \circ \beta$. We choose this to be the half twist which we denote by hT.



Since π is an op-fibration, we obtain for a pair (A, B) a cocartesian morphism $\beta': (A, B) \rightarrow (B, A)$ in $\mathcal{C}^2$ with $\pi(\beta') = \beta$.

Since $\otimes$ and β' are cocartesian and composites of cocartesian morphism are again cocartesian, we can lift the diagram in both directions:

$$\begin{array}{ccc} \langle 2 \rangle & \xrightarrow{\beta} & \langle 2 \rangle \\ \downarrow \mu & & \downarrow \mu \\ \langle 1 \rangle & \dashrightarrow & \langle 1 \rangle. \end{array}$$

and obtain two commuting diagram in $\mathcal{C}^{\otimes}$:

$$\begin{array}{ccc} (A, B) & \xrightarrow{\beta} & (B, A) \\ \downarrow \otimes & & \downarrow \otimes \\ A \otimes B & \xrightarrow[b^{-1}]{b} & B \otimes A \end{array}$$

4. Colored Braided Operads

The morphism b is the induced morphism from the cocartesian morphism $\otimes$ while b^{-1} is induced by $\otimes \circ \beta'$. They are inverse to one another since lifts with chosen source are unique. The induced morphism b is a unique isomorphism due to the fact that $\otimes$ and β are cocartesian morphisms. Note, that for a homotopic different choice of path instead of the half twist, we would obtain a different braiding. So, this choice is essential for our construction. We recall the following morphisms from $\mathcal{F}in_*$ for $n \in \mathbb{N}$ and $1 \leq i \leq n$:

$$\tau_i^n: \langle n \rangle \rightarrow \langle n-1 \rangle$$

$$j \mapsto \begin{cases} j & 1 \leq j \leq i \\ j-1 & i < j \leq n \\ * & j = * \end{cases}$$


We consider these morphisms in $\mathcal{B}r_*$ with the following collection of configurations:

$$\widetilde{\tau}_{i,j}^n := \begin{cases} b_2 & i = j \\ \text{Id}^j & \text{otherwise} \end{cases}$$

For the associator we consider the following diagram in $\mathcal{B}r_*$:

$$\begin{array}{ccc} \langle 3 \rangle & \xrightarrow{\tau_2^3} & \langle 2 \rangle \\ \downarrow \tau_1^3 & & \downarrow \mu \\ \langle 2 \rangle & \xrightarrow{\mu} & \langle 1 \rangle \end{array}$$

where the associated configurations are either the trivial configuration or the basepoint configuration. To fill the diagram, we choose a path γ that only translates the middle points of the disks along the horizontal line and shrinks or expands the disks as needed.



γ lies in a sense in E_1 and we do not care too much about $E_1 \dots$

Again using the op-fibration condition (M1) property of π we obtain a commuting diagram in $\mathcal{C}^\otimes$:

$$\begin{array}{ccccc} (A \otimes B, C) & \xleftarrow{(\otimes, \text{Id})} & (A, B, C) & \xrightarrow{(\text{Id}, \otimes)} & (A, B \otimes C) \\ \downarrow \otimes & & & & \downarrow \otimes \\ (A \otimes B) \otimes C & \xrightarrow{a} & & & A \otimes (B \otimes C) \end{array}$$

Similarly to Proposition 2.20, the tensor unit is given by the unique morphism $\eta: \langle 0 \rangle \rightarrow \langle 1 \rangle$ in $\mathcal{B}r_*$ which induces a functor $\mathcal{C}_{(0)}^\otimes \rightarrow \mathcal{C}$ giving us a tensor unit $\mathbf{1}$.

Proposition 4.8. *The category $\mathcal{C}$ equipped with $(\otimes, b, a, \mathbf{1})$ defines a braided monoidal category.*

Proof. First, we observe that the associator refers to E_1 being embedded into E_2 along the horizontal line, as we translate and resize the disks along the horizontal line. So, we can arbitrarily translate and resize configurations with middle points on the horizontal axis, since it is only a associative translation in E_1 .

So, the pentagon identity holds as E_1 is the associative operad. The unitality condition holds analogously to the proof of Proposition 2.20 using the Segal condition. Hence, we focus on verifying the hexagon identities

include a proof here if spare time, at least elaborate on E_1

4. Colored Braided Operads






Figure 2: example of a configuration of E_1 being embedded in E_2 along the horizontal line


1.




2.



Each edge in the hexagon diagrams refers to a commuting diagram in $\mathcal{B}r_*$ of the form



with configurations:




4. Colored Braided Operads

The commutativity of the triangle diagrams on the side means choosing a path from one configuration to another while filling the tetrahedron in $\mathcal{Br}_*$ means having a homotopy between the different paths. As it turns out we do not make any new choices of paths. The paths are induced by the half twist hT we chose to define $\otimes$:

Let us understand where the diagram or more precisely the configurations come from. The morphisms corresponding to $c_{(A \otimes B) \otimes C}$ and $c_{C \otimes (A \otimes B)}$ respectively factor through $\langle 2 \rangle$ in the following way:


$$\begin{array}{ccc}
 & \langle 3 \rangle & \\
 c_{(A \otimes B) \otimes C} \swarrow & \downarrow (b_2, \text{Id}^3) & \searrow c_{C \otimes (A \otimes B)} \\
 & \langle 2 \rangle & \\
 \mu \swarrow & & \searrow \mu \circ \beta \\
 \langle 1 \rangle & \xrightarrow{\text{hT}} & \langle 1 \rangle
 \end{array}$$

We observe, that the outer triangle commutes precisely because the inner triangle commutes which is the case since we already chose the path to be the half twist hT.




Similar arguments for all the corresponding diagrams induced by the hexagon identity (1) give us paths that are already defined by the associativity path and the half twist up to homotopy.

4. Colored Braided Operads




The resulting paths are the following representatives of their homotopy class and they are clearly homotopy equivalent.

4. Colored Braided Operads




Analogously for the hexagon identity (2), we consider the following composition of paths in $\mathcal{Br}_*$:



5. Outlook

The resulting paths are again homotopy equivalent and therefore the second hexagon identity holds.



This concludes the proof that $\mathcal{B}r_*$ induces a braided monoidal structure on $\mathcal{C}$.

□

4.2 Colored Braided Operads as Op-fibrations

Let $\mathcal{B}$ be a colored braided operad, then there is a forgetful functor

$$\pi: \mathcal{B}^\otimes \rightarrow \mathcal{B}r_*$$

In this section we will focus on characterizing π .

5 Outlook

Notes on outlook:

- action operads: Could we define a op-fibration definition for any group operad?

6 Appendix

We wanted to define braided operads as fibrations over a suitable category. A first intuition led us to the assumption that colored braided operads might fiber over the crossed simplicial braid group as defined in [DK]. However, we discovered that we were too quick to make this assumption. We will elaborate in the following why this did not work out as we hoped originally.

Definition 6.1. The *crossed simplicial braid group* is a category ΔB equipped with an embedding $\iota: \Delta \rightarrow \Delta B$ such that:

1. the functor ι is bijective on objects,
2. any morphism $u: \iota[m] \rightarrow \iota[n]$ in ΔB can be uniquely expressed as a composition $\iota(\phi) \circ b$ where $\phi: [m] \rightarrow [n]$ is a morphism in Δ and $b \in B_m$ is acting on $\iota[m]$ in ΔB .

Note that we sometimes refer to $\iota[m]$ by writing $[m]$ instead.

Remark 6.2. To understand composition in the category ΔB let us consider two morphisms

$$f: [m] \rightarrow [n] \quad \text{and} \quad g: [n] \rightarrow [l]$$

with the factorization $f = \phi_f \circ b_f$ and $g = \phi_g \circ b_g$. We define for every $j \in [n]$ an f -pencil p_j as a collection $[j, i_1, \dots, i_s]$ with $\phi_f^{-1}(j) = \{i_1 < \dots < i_s\}$. We call s the width of the pencil. Note that we say it is an empty pencil p_j if $\phi_f^{-1}(j) = \emptyset$.

Then we obtain a braid on m strands in the following way: First, we consider $p_1, \dots, p_n$ as n strand such that we can braid by b_g . Then, we forget the empty pencil strands; in other


if time: Define colored braided operad from a braided mon cat.

in this section we finally try to define a fibration from the colored braided operad to $\mathcal{B}r_*$ - good luck!

43

6. Appendix

So, we identify the following graphically:



As it turns out, a short graphical calculation shows that the hexagon identities hold for $\Delta B'$. Unfortunately, we face the next challenge as we discovered that we obtain a braid group action where we would not expect one and moreover where we not want a braid group action: Let $\sigma \in B_n$, then for $\sigma: [n] \rightarrow [n]$ we obtain cocartesian morphisms in $\mathcal{O}_{[n]}^\otimes$ for each $X \in \mathcal{O}_{[n]}^\otimes \cong \mathcal{O}^n$, so the braid group B_n acts on $\mathcal{O}^n$. However, we would expect only to gain a symmetric group action on $\mathcal{O}^n$.

Thus, we have to conclude that the crossed simplicial braid group is not the right candidate for our op-fibration.

add
hexagon
identities

explain
braid action
on cartesian
product

Bibliography

- [AB] Adrian Boerner. “Comparing different models of the little disk operad”. June 2025.
- [DK] T. Dyckerhoff and M. Kapranov. *Crossed simplicial groups and structured surfaces*. 2015.
- [ZF] Z. Fiedorowicz. *Constructions of E_n Operads*. 1999.
- [IM] Igor Kříž and J. P. May. *Operads, algebras, modules and motives*. en. Astérisque 233. Société mathématique de France, 1995.
- [JL] Jacob Lurie. “Higher algebra”. Unpublished. Available online at <https://www.math.ias.edu/~lurie/>. Sept. 2017.
- [Kerodon] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2018.
- [CS] Christoph Schweigert. “Hopf algebras, quantum groups and topological field theory”. Unpublished. Available online at <https://www.math.uni-hamburg.de/home/schweigert/skripten/hskript.pdf>. Mar. 2023.
- [DY] Donald Yau. *Infinity Operads and Monoidal Categories with Group Equivariance*. WORLD SCIENTIFIC, 2021.