

Logical Formalism

Set Theory

Adrien Pommellet, LRE




August 31, 2023

What Is a Set?

Collecting objects

- We need to be able to **gather** and manipulate objects.
- A **set** is merely a well-defined collection of objects.

 \mathbb{N} , \mathbb{R} , the prime numbers \mathbb{P} , the points of a line, etc.

What Is a Set?

The \in predicate

- Formally, a set E is matched to a **predicate** \in such that x **belongs** to E if and only if the statement $x \in E$ is true.
- We also introduce the predicate $x \notin E \iff \neg(x \in E)$.
- We define the **empty set** \emptyset such that no element belongs to \emptyset : the proposition $x \in \emptyset$ is always false.

What Is a Set?

Representing sets

- Each object in a set can appear **only once**: there are no duplicates.
 - A finite set containing n elements can be written $\{x_1, \dots, x_n\}$. This indexing is a purely arbitrary choice and may be **unordered**.
 - The set $\{x\}$ containing only the element x is called a **singleton**. It should not be mistaken with x itself.
- E** $\{0, 1, 2\}$ and $\{1, 0, 2\}$ represent the same set, whereas $\{1, 1, 2\}$ is not a set.

What Is a Set?

Shaky foundations

Set theory is somewhat flawed because of the **catalogue paradox**.

Consider a library that follows **three rules**:

- 1 Each library section features a catalogue listing its contents.
- 2 Some catalogues may list themselves, others do not.
- 3 A special catalogue lists all the catalogues that do not list themselves.




The paradox consists in the following question: **does the special catalogue list itself?**

Operations on Sets

The union \cup

Given two sets A and B , we introduce their **union** $A \cup B$ such that $x \in A \cup B$ if and only if $(x \in A) \vee (x \in B)$.




Consider $\mathbb{Z} = \mathbb{Z}_+ \cup \mathbb{Z}_-$. Note that 0 belongs to **both** \mathbb{Z}_+ and \mathbb{Z}_- .

Operations on Sets

The Intersection \cap

Given two sets A and B , we introduce their **intersection** $A \cap B$ such that $x \in A \cap B$ if and only if $(x \in A) \wedge (x \in B)$.



The intersection of two distinct lines in \mathbb{R}^2 is either \emptyset or a singleton.

Operations on Sets

The subset relation \subseteq

Given two sets A and B , we introduce the **subset relation**: $A \subseteq B$ if and only if $(x \in A) \implies (x \in B)$.



You should already know that $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.

Operations on Sets

The simple inclusion pattern

The logical implication \implies yields the following **proof pattern**:

Goal. Prove that $A \subseteq B$.

Let $x \in A \dots$
 \dots then $x \in B$.

Exercise 1. Let $A = \{a, b, c\}$ and $B = \{a, d\}$. Explicit the sets:

$$A \cup A$$

$$A \cap A$$

$$A \cup B$$

$$A \cap B$$

$$A \cap B \cup \{e\}$$

Answer

Operations on Sets

Equality of sets

- Given two sets A and B , we introduce the **equality relation**: $A = B$ if and only if $(x \in A) \iff (x \in B)$.
- Remember that $(P \iff Q) \iff (P \implies Q) \wedge (Q \implies P)$.
- As a consequence, $(A = B) \iff (A \subseteq B) \wedge (B \subseteq A)$.

Operations on Sets

The double inclusion pattern

Goal. Prove that $A = B$.

Subgoal 1. Prove that $A \subseteq B$.

Let $x \in A \dots$
... then $x \in B$.

Subgoal 2. Prove that $B \subseteq A$.

Let $x \in B \dots$
... then $x \in A$.

Set Theory

Power Sets

- Given a set E , we introduce $\mathcal{P}(E)$, the **power set** of E such that $X \in \mathcal{P}(E) \iff X \subseteq E$.
- $\mathcal{P}(E)$ is a **set of sets**.
- Note that $\emptyset \in \mathcal{P}(E)$ and $E \in \mathcal{P}(E)$ for any set E .


Exercise 2. Explicit the set $\mathcal{P}(\{a, b, c\})$.

Answer

Set Theory

The set subtraction \setminus

Given two sets A and B , we introduce the **subtraction** $A \setminus B$ of A by B such that $x \in A \setminus B$ if and only if $(x \in A) \wedge (x \notin B)$.




B isn't necessarily a subset of A : $\{0, 1, 2, 3\} \setminus \{1, 3, 4\} = \{0, 2\}$.

Set Theory

The complement \complement

Given an **implicit** set E and a subset $A \subseteq E$, the **complement** of A with regard to E is the set $A^\complement = E \setminus A$.



E

Given $A = \{0\}$, $A^\complement = \mathbb{N}^*$ with regard to \mathbb{N} and $A^\complement = \mathbb{R}^*$ with regard to \mathbb{R} .

Properties of Sets

Sets and propositional logic

Sets

$$A \cup B$$

$$A \cap B$$

$$A \subseteq B$$

$$A = B$$

$$A^c$$

Logical Operators

$$P \vee Q$$

$$P \wedge Q$$

$$P \implies Q$$

$$P \iff Q$$

$$\neg P$$

Properties of Sets

Common equalities

The following properties on sets hold, assuming $A, B, C \in \mathcal{P}(E)$:

$$A^{\complement\complement} = A$$

$$A \cup A^{\complement} = E$$

$$A \setminus B = A \cap B^{\complement}$$

$$(A \cup B)^{\complement} = A^{\complement} \cap B^{\complement}$$

$$(A \cap B)^{\complement} = A^{\complement} \cup B^{\complement}$$

$$\emptyset^{\complement} = E$$

$$A \subseteq B \iff A \cap B^{\complement} = \emptyset$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Practical Application

Exercise 3. Are the following properties true in the general case, assuming $A, B, C \in \mathcal{P}(E)$? If they're not, use **counter-examples** to falsify them.

$$A = B \iff A \cup C = B \cup C$$

$$A = B \iff A \cap B = A$$

$$A \cup B = C \iff A \cup C = B$$

$$A \subseteq B \cup C \iff (A \subseteq B) \vee (A \subseteq C)$$

$$A \subseteq B \cap C \iff (A \subseteq B) \wedge (A \subseteq C)$$

$$A = B \iff A \setminus C = B \setminus C$$

$$(A \subseteq C) \wedge (B \subseteq C) \implies A \subseteq B$$

$$A \cup B = C \iff A = C \setminus B$$

$$A \cup B + A \cap B = A + B \iff A \cup B = A + B - A \cap B$$

Answer I

Answer II

Practical Application

Exercise 4. Are the following equalities true in the general case, assuming $A, B, C \in \mathcal{P}(E)$? If they're not, use **counter-examples** to falsify them.

$$(A \setminus B) \cup B = A$$

$$(A \cup B) \setminus B = A$$

$$(A \cup B \cup B) \setminus B = A \cup B$$

$$(A \setminus B) \cup (A \cap B) = A$$

$$\emptyset \cup A = E \cap A$$

$$\emptyset \cap A = E \cup A$$

$$A + B - B = A$$

$$A \cup B \cup B = A + 2B$$

$$A \cup B = (A \cup B) \setminus (A \cap B)$$

Answer 1

Answer II

A Last Few Definitions

The Cartesian product \times

- We consider **ordered pairs** of objects. We use the notation (x, y) .
- Do not mistake the pair (x, y) for the set (also called an unordered pair) $\{x, y\}$.
- Given two sets U and V , we introduce their **Cartesian product** $U \times V$ such that $(x, y) \in U \times V$ if and only if $(x \in U) \wedge (y \in V)$.



Consider $\mathbb{R}^2 = (\mathbb{R}_+ \times \mathbb{R}_+) \cup (\mathbb{R}_+ \times \mathbb{R}_-) \cup (\mathbb{R}_- \times \mathbb{R}_-) \cup (\mathbb{R}_- \times \mathbb{R}_+)$.

A Last Few Definitions

Qualifying sets

- A common way to define a set is to select all the elements of a set **verifying a given proposition**.
- Formally, given a proposition $P(x)$ and a set E , we can introduce a subset S of E such that $x \in S \iff (x \in E) \wedge P(x)$.
- We often write $S = \{x \in E \mid P(x)\}$ instead.

E

For $a, b \in \mathbb{R}$, the set $S = \{(x, y) \in \mathbb{R}^2 \mid y = a \cdot x + b\}$ is a line of the real plane \mathbb{R}^2 .