

Finite Dimensional Approximations of Spectral Triples on Quantum tori

Frédéric Latrémolière



North Atlantic Noncommutative Geometry Seminar
March, 24th 2021.



- 1 *Introduction*
- 2 *Quantum Compact Metric Spaces*
- 3 *Convergence of Metric Spectral Triples*
- 4 *Examples*

Why a geometry for a space of quantum spaces

Our goal is to construct *geometries* on *spaces of quantum spaces*, typically using metrics.

Why a geometry for a space of quantum spaces

Our goal is to construct *geometries* on *spaces of quantum spaces*, typically using metrics.

Quantum Gravity

Is a theory of quantum gravity a form of dynamics over a space of (quantum) space-times?

Why a geometry for a space of quantum spaces

Our goal is to construct *geometries* on *spaces of quantum spaces*, typically using metrics.

Quantum Gravity

Is a theory of quantum gravity a form of dynamics over a space of (quantum) space-times?

Finite Dimensional Approximations for quantum spaces

Can a physical model be a statistical limit of finite models over quantum spaces?

Why a geometry for a space of quantum spaces

Our goal is to construct *geometries* on *spaces of quantum spaces*, typically using metrics.

Quantum Gravity

Is a theory of quantum gravity a form of dynamics over a space of (quantum) space-times?

Finite Dimensional Approximations for quantum spaces

Can a physical model be a statistical limit of finite models over quantum spaces?

New C^ -algebraic constructions*

What are the properties of classes of quantum metric spaces, and in particular, what does metric geometry tells us about noncommutative spaces?

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

Founding Allegory of Noncommutative Geometry

Noncommutative geometry is the study of noncommutative generalizations of algebras of smooth functions over geometric spaces.

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

*Founding Allegory of Noncommutative **Metric** Geometry*

Noncommutative *metric* geometry is the study of noncommutative generalizations of algebras of *Lipschitz* functions over *metric* spaces.

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

*Founding Allegory of Noncommutative **Metric** Geometry*

Noncommutative *metric* geometry is the study of noncommutative generalizations of algebras of *Lipschitz* functions over *metric* spaces.

- Pioneered by *Rieffel (1998–)*, inspired by *Connes (1989)*.

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

*Founding Allegory of Noncommutative **Metric** Geometry*

Noncommutative *metric* geometry is the study of noncommutative generalizations of algebras of *Lipschitz* functions over *metric* spaces.

- Pioneered by *Rieffel (1998–)*, inspired by *Connes (1989)*.
- Motivated by mathematical physics.

A geometry for the space of quantum spaces

The Project

My project brings together ideas from *metric geometry* and *noncommutative geometry* to construct a geometry on *hyperspaces of quantum spaces*.

*Founding Allegory of Noncommutative **Metric** Geometry*

Noncommutative *metric* geometry is the study of noncommutative generalizations of algebras of *Lipschitz* functions over *metric* spaces.

- Pioneered by *Rieffel (1998–)*, inspired by *Connes (1989)*.
- Motivated by mathematical physics.
- The focus is on various generalizations of the *Gromov-Hausdorff* distance.

- 1 *Introduction*
- 2 *Quantum Compact Metric Spaces*
- 3 *Convergence of Metric Spectral Triples*
- 4 *Examples*

The Monge-Kantorovich metric

Let (X, \mathfrak{m}) be a compact metric space. The *Lipschitz seminorm* L induced by \mathfrak{m} is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$ (allowing ∞).

The Monge-Kantorovich metric

Let (X, \mathfrak{m}) be a compact metric space. The *Lipschitz seminorm* L induced by \mathfrak{m} is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$ (allowing ∞).

The *Monge-Kantorovich metric* on $\mathcal{S}(C(X))$ is given for all Borel-regular probability measures μ, ν by:

$$\mathsf{mk}_{\mathsf{L}}(\mu, \nu) = \sup \left\{ \left| \int_X f d\mu - \int_X f d\nu \right| : f \in \mathfrak{sa}(C(X)), \mathsf{L}(f) \leq 1 \right\}.$$

The Monge-Kantorovich metric

Let (X, \mathfrak{m}) be a compact metric space. The *Lipschitz seminorm* L induced by \mathfrak{m} is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$ (allowing ∞).

The *Monge-Kantorovich metric* on $\mathcal{S}(C(X))$ is given for all Borel-regular probability measures μ, ν by:

$$\mathsf{mk}_{\mathsf{L}}(\mu, \nu) = \sup \left\{ \left| \int_X f d\mu - \int_X f d\nu \right| : f \in \mathfrak{sa}(C(X)), \mathsf{L}(f) \leq 1 \right\}.$$

The Gelfand map $x \in (X, \mathfrak{m}) \mapsto \delta_x \in (\mathcal{S}(C(X)), \mathsf{mk}_{\mathsf{L}})$ is an isometry.

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

(\mathfrak{A}, L) is a *quantum compact metric space* when:

- 1 \mathfrak{A} is a *unital C^* -algebra*,

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathsf{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② L is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathsf{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathsf{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② L is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathsf{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,
- ④ The *weak* topology on $\mathcal{S}(\mathfrak{A})$* is metrized by the *Monge-Kantorovich metric* mk_{L} , defined $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$ by:

$$\text{mk}_{\mathsf{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathsf{L}(a) \leq 1 \}$$

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathbb{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② \mathbb{L} is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathbb{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathbb{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,
- ④ The *weak* topology on $\mathcal{S}(\mathfrak{A})$* is metrized by the *Monge-Kantorovich metric* $\text{mk}_{\mathbb{L}}$, defined $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$ by:

$$\text{mk}_{\mathbb{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathbb{L}(a) \leq 1 \}$$

- ⑤ $\max \{ \mathbb{L}(a \circ b), \mathbb{L}(\{a, b\}) \} \leq \|a\|_{\mathfrak{A}} \mathbb{L}(b) + \|b\|_{\mathfrak{A}} \mathbb{L}(a),$

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathsf{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② L is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathsf{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,
- ④ The *weak* topology on $\mathcal{S}(\mathfrak{A})$* is metrized by the *Monge-Kantorovich metric* mk_{L} , defined $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$ by:

$$\text{mk}_{\mathsf{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathsf{L}(a) \leq 1 \}$$

- ⑤ $\max \{ \mathsf{L}(a \circ b), \mathsf{L}(\{a, b\}) \} \leq F(\|a\|_{\mathfrak{A}}, \|b\|_{\mathfrak{A}}, \mathsf{L}(a), \mathsf{L}(b)),$

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathbb{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② \mathbb{L} is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathbb{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathbb{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,
- ④ The *weak* topology on $\mathcal{S}(\mathfrak{A})$* is metrized by the *Monge-Kantorovich metric* $\text{mk}_{\mathbb{L}}$, defined $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$ by:

$$\text{mk}_{\mathbb{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathbb{L}(a) \leq 1 \}$$

- ⑤ $\max \{ \mathbb{L}(a \circ b), \mathbb{L}(\{a, b\}) \} \leq F(\|a\|_{\mathfrak{A}}, \|b\|_{\mathfrak{A}}, \mathbb{L}(a), \mathbb{L}(b))$,
- ⑥ \mathbb{L} is lower semi-continuous wrt $\|\cdot\|_{\mathfrak{A}}$.

Compact Quantum Metric Spaces

Definition (Connes, 89; Rieffel, 98; L., 13)

$(\mathfrak{A}, \mathsf{L})$ is a *quantum compact metric space* when:

- ① \mathfrak{A} is a *unital C^* -algebra*,
- ② L is a *seminorm* defined on a (dense) Jordan-Lie subalgebra $\text{dom}(\mathsf{L})$ of $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$,
- ③ $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$,
- ④ The *weak* topology on $\mathcal{S}(\mathfrak{A})$* is metrized by the *Monge-Kantorovich metric* mk_{L} , defined $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$ by:

$$\text{mk}_{\mathsf{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathsf{L}(a) \leq 1 \}$$

- ⑤ $\max \{ \mathsf{L}(a \circ b), \mathsf{L}(\{a, b\}) \} \leq F(\|a\|_{\mathfrak{A}}, \|b\|_{\mathfrak{A}}, \mathsf{L}(a), \mathsf{L}(b))$,
- ⑥ L is lower semi-continuous wrt $\|\cdot\|_{\mathfrak{A}}$.

We call L an *L -seminorm*.

Some examples

- *Rieffel, 98*: unital C^* -algebras with ergodic metric compact group actions,
- *Ozawa, Rieffel, 02*: word-Hyperbolic discrete group C^* -algebras with Connes' length function spectral triple,
- *Li, 03*: Connes-Landi-du Bois-Voilette's spheres wrt spectral triple,
- *Hawkins, Skalski, White, Zacharias, 13*: C^* -crossed products for equicontinuous actions wrt spectral triple,
- *L., Aguilar*: AF algebras with faithful tracial state,
- *L., 16*: Curved quantum tori, with metric from spectral triple,
- *L., 16*: Perturbations (conformal and otherwise) of metric spectral triples,
- *Christ, Rieffel, 16*: Nilpotent group C^* -algebras with Connes' length function spectral triple,
- *Aguilar, Kaad, 17*: Podles spheres,
- *Kaad, Kyed, 19*: C^* -crossed products for actions of \mathbb{Z} by "Lipschitz" morphisms (twisted spectral triple),

Lipschitz morphisms

Theorem-Definition (Lipschitz Morphisms)

Let $(\mathfrak{A}, L_{\mathfrak{A}})$ and $(\mathfrak{B}, L_{\mathfrak{B}})$ be two quasi-Leibniz quantum compact metric spaces. A *Lipschitz morphism* $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$ is a unital $*$ -morphism from \mathfrak{A} to \mathfrak{B} such that any of the following equivalent statement holds:

- ❶ $(\exists k) \varphi \in \mathcal{S}(\mathfrak{B}) \mapsto \varphi \circ \pi$ is a *k-Lipschitz map from $(\mathcal{S}(\mathfrak{B}), \text{mk}_{L_{\mathfrak{B}}})$ to $(\mathcal{S}(\mathfrak{A}), \text{mk}_{L_{\mathfrak{A}}})$,*
- ❷ (Rieffel, 00) $(\exists k) L_{\mathfrak{B}} \circ \pi \leq k L_{\mathfrak{A}}$,
- ❸ (L., 16) $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$.

Lipschitz morphisms

Theorem-Definition (Lipschitz Morphisms)

Let $(\mathfrak{A}, L_{\mathfrak{A}})$ and $(\mathfrak{B}, L_{\mathfrak{B}})$ be two quasi-Leibniz quantum compact metric spaces. A *Lipschitz morphism* $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$ is a unital $*$ -morphism from \mathfrak{A} to \mathfrak{B} such that any of the following equivalent statements holds:

- ① $(\exists k) \varphi \in \mathcal{S}(\mathfrak{B}) \mapsto \varphi \circ \pi$ is a *k-Lipschitz map from $(\mathcal{S}(\mathfrak{B}), \text{mk}_{L_{\mathfrak{B}}})$ to $(\mathcal{S}(\mathfrak{A}), \text{mk}_{L_{\mathfrak{A}}})$,*
- ② (Rieffel, 00) $(\exists k) L_{\mathfrak{B}} \circ \pi \leq k L_{\mathfrak{A}}$,
- ③ (L., 16) $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$.

Lipschitz morphisms are arrows for a category over quasi-Leibniz quantum compact metric spaces.

Quantum Isometries

If $X \subset (Z, \mathfrak{m})$ and $f \in \mathfrak{sa}(C(X))$:

- if $g \in C(Z)$ and $g|_X = f$, the $\text{Lip}(f) \leq \text{Lip}(g)$,
- there exists $g \in \mathfrak{sa}(C(Z))$ with $g|_X = f$ and $\text{Lip}(f) = \text{Lip}(g)$ (McShane, 34).

Quantum Isometries

If $X \subset (Z, \mathfrak{m})$ and $f \in \mathfrak{sa}(C(X))$:

- if $g \in C(Z)$ and $g|_X = f$, the $\text{Lip}(f) \leq \text{Lip}(g)$,
- there exists $g \in \mathfrak{sa}(C(Z))$ with $g|_X = f$ and $\text{Lip}(f) = \text{Lip}(g)$ (McShane, 34).

Definition (Rieffel (98), L. (13))

A *quantum isometry* $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$ is a $*$ -epimorphism such that:

$$\forall b \in \text{dom}(\mathbb{L}_{\mathfrak{B}})$$

$$\mathbb{L}_{\mathfrak{B}}(b) = \inf \{ \mathbb{L}_{\mathfrak{A}}(a) : \pi(a) = b \}.$$

A *full quantum isometry* π is a $*$ -isomorphism such that $\mathbb{L}_{\mathfrak{B}} \circ \pi = \mathbb{L}_{\mathfrak{A}}$.

Quantum Isometries

If $X \subset (Z, \mathfrak{m})$ and $f \in \mathfrak{sa}(C(X))$:

- if $g \in C(Z)$ and $g|_X = f$, the $\text{Lip}(f) \leq \text{Lip}(g)$,
- there exists $g \in \mathfrak{sa}(C(Z))$ with $g|_X = f$ and $\text{Lip}(f) = \text{Lip}(g)$ (McShane, 34).

Definition (Rieffel (98), L. (13))

A *quantum isometry* $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$ is a $*$ -epimorphism such that:

$$\forall b \in \text{dom}(\mathbb{L}_{\mathfrak{B}}) \quad \mathbb{L}_{\mathfrak{B}}(b) = \inf \{ \mathbb{L}_{\mathfrak{A}}(a) : \pi(a) = b \}.$$

A *full quantum isometry* π is a $*$ -isomorphism such that $\mathbb{L}_{\mathfrak{B}} \circ \pi = \mathbb{L}_{\mathfrak{A}}$.

The dual maps of quantum isometries are indeed isometries for the Monge-Kantorovich metric.

The Gromov-Hausdorff Distance

Definition

For any two compact metric spaces (X, \mathbf{m}_X) and (Y, \mathbf{m}_Y) , we define $\text{Adm}(\mathbf{m}_X, \mathbf{m}_Y)$ as:

$$\left\{ (Z, \mathbf{m}_Z, \iota_X, \iota_Y) \left| \begin{array}{l} (Z, \mathbf{m}_Z) \text{ compact metric space,} \\ \iota_X : X \hookrightarrow Z \text{ isometry,} \\ \iota_Y : Y \hookrightarrow Z \text{ isometry.} \end{array} \right. \right\}$$

The Gromov-Hausdorff Distance

Definition

For any two compact metric spaces (X, m_X) and (Y, m_Y) , we define $\text{Adm}(m_X, m_Y)$ as:

$$\left\{ (Z, m_Z, \iota_X, \iota_Y) \left| \begin{array}{l} (Z, m_Z) \text{ compact metric space,} \\ \iota_X : X \hookrightarrow Z \text{ isometry,} \\ \iota_Y : Y \hookrightarrow Z \text{ isometry.} \end{array} \right. \right\}$$

Notation

The *Hausdorff distance* on the compact subsets of a metric space (X, m) is denoted by Haus_m .

Definition (Edwards, 75; Gromov, 81)

The *Gromov-Hausdorff distance* between two compact metric spaces (X, m_X) and (Y, m_Y) is:

$$\inf \{ \text{Haus}_{m_Z}(\iota_X(X), \iota_Y(Y)) : (Z, m_Z, \iota_X, \iota_Y) \in \text{Adm}(m_X, m_Y) \}.$$

The Dual Gromov-Hausdorff Propinquity




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

The Dual Gromov-Hausdorff Propinquity




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

Definition (The extent of a tunnel, L. 13,14)

The *extent* $\chi(\tau)$ of a tunnel $\tau = (\mathfrak{D}, L_{\mathfrak{D}}, \pi_{\mathfrak{A}}, \pi_{\mathfrak{B}})$ is:

$$\max \left\{ \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left(\mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{A}}^* (\mathcal{S}(\mathfrak{A})) \right), \right. \\ \left. \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left(\mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{B}}^* (\mathcal{S}(\mathfrak{B})) \right) \right\}.$$

The Dual Gromov-Hausdorff Propinquity




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

Definition (The Dual Propinquity, L. 13, 14)

The *extent* $\chi(\tau)$ of a tunnel $\tau = (\mathfrak{D}, L_{\mathfrak{D}}, \pi_{\mathfrak{A}}, \pi_{\mathfrak{B}})$ is:

$$\max \left\{ \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left(\mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{A}}^* (\mathcal{S}(\mathfrak{A})) \right), \right. \\ \left. \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left(\mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{B}}^* (\mathcal{S}(\mathfrak{B})) \right) \right\}.$$

The *dual propinquity* $\Lambda_F^*((\mathfrak{A}, L_{\mathfrak{A}}), (\mathfrak{B}, L_{\mathfrak{B}}))$ is given by:

$$\inf \{ \chi(\tau) : \tau \text{ any } F\text{-tunnel from } (\mathfrak{A}, L_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, L_{\mathfrak{B}}) \}.$$

The Dual Gromov-Hausdorff Propinquity




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

Theorem (L., 13)

The *dual propinquity* Λ_F^* , defined for any two quantum compact metric spaces $(\mathfrak{A}, L_{\mathfrak{A}})$ by $(\mathfrak{B}, L_{\mathfrak{B}})$ by:

$$\inf \{ \chi(\tau) : \tau \text{ any } F\text{-tunnel from } (\mathfrak{A}, L_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, L_{\mathfrak{B}}) \}$$

is a *complete metric* up to *full quantum isometry*.
 $\Lambda((\mathfrak{A}, L_{\mathfrak{A}}), (\mathfrak{B}, L_{\mathfrak{B}})) = 0$ iff there exists a $*$ -isomorphism $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ such that $L_{\mathfrak{B}} \circ \pi = L_{\mathfrak{A}}$.

The Dual Gromov-Hausdorff Propinquity




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

Theorem (L., 13)

The *dual propinquity* Λ_F^* , defined for any two quantum compact metric spaces $(\mathfrak{A}, \mathcal{L}_{\mathfrak{A}})$ by $(\mathfrak{B}, \mathcal{L}_{\mathfrak{B}})$ by:

$$\inf \{ \chi(\tau) : \tau \text{ any } F\text{-tunnel from } (\mathfrak{A}, \mathcal{L}_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, \mathcal{L}_{\mathfrak{B}}) \}$$

is a *complete metric* up to *full quantum isometry*. Moreover Λ^* induces the topology of the *Gromov-Hausdorff distance* on compact metric spaces.

Examples of Convergence

Example

- *L., 13* proved that fuzzy tori converge to quantum tori and quantum tori form a continuous family, for a natural metric from the dual actions,
- *Rieffel, 15* has constructed full matrix approximations of $C(S^2)$ using metrics arising from actions of $SU(2)$.
- *L. and Packer, 16* proved continuity for the family of noncommutative solenoids,
- *Aguilar, L., 15* proved certain continuity results for AF algebras,
- *L., 15* proved continuity for curved quantum tori,
- *L., 15* proved continuity for conformal deformations,
- *L., 15* proved that nuclear quasi-diagonal quantum compact metric spaces are limits of finite dimensional quantum compact metric spaces,
- and more ...

- 1 *Introduction*
- 2 *Quantum Compact Metric Spaces*
- 3 *Convergence of Metric Spectral Triples*
- 4 *Examples*

Standard description of a noncommutative geometry

Definition (Connes, 89)

A **spectral triple** $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is given by:

- ① a unital C^* -algebra \mathfrak{A} ,
- ② a Hilbert space \mathcal{H} on which \mathfrak{A} acts,
- ③ a self-adjoint operator \mathcal{D} on $\text{dom}(\mathcal{D}) \subseteq \mathcal{H}$, such that:
 - $\mathcal{D} \pm i$ has a compact inverse,
 - $\mathfrak{A}^1 = \{a \in \mathfrak{A} : a \cdot \text{dom}(\mathcal{D}) \subseteq \text{dom}(\mathcal{D}), [\mathcal{D}, a] \text{ has bounded closure}\}$ is dense in \mathfrak{A} .

Standard description of a noncommutative geometry

Definition (Connes, 89)

A **spectral triple** $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is given by:

- ① a unital C^* -algebra \mathfrak{A} ,
- ② a Hilbert space \mathcal{H} on which \mathfrak{A} acts,
- ③ a self-adjoint operator \mathcal{D} on $\text{dom}(\mathcal{D}) \subseteq \mathcal{H}$, such that:
 - $\mathcal{D} \pm i$ has a compact inverse,
 - $\mathfrak{A}^1 = \{a \in \mathfrak{A} : a \cdot \text{dom}(\mathcal{D}) \subseteq \text{dom}(\mathcal{D}), [\mathcal{D}, a] \text{ has bounded closure}\}$ is dense in \mathfrak{A} .

If $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is a spectral triple and if for any $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$, we set:

$$\text{mk}_{\mathcal{D}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), ||| [\mathcal{D}, a] |||_{\mathcal{H}} \leq 1 \}$$

then $\text{mk}_{\mathcal{D}}$ is an extended pseudo-distance on $\mathcal{S}(\mathfrak{A})$.

Standard description of a noncommutative geometry

Definition (Connes, 89)

A **spectral triple** $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is given by:

- ① a unital C^* -algebra \mathfrak{A} ,
- ② a Hilbert space \mathcal{H} on which \mathfrak{A} acts,
- ③ a self-adjoint operator \mathcal{D} on $\text{dom}(\mathcal{D}) \subseteq \mathcal{H}$, such that:
 - $\mathcal{D} \pm i$ has a compact inverse,
 - $\mathfrak{A}^1 = \{a \in \mathfrak{A} : a \cdot \text{dom}(\mathcal{D}) \subseteq \text{dom}(\mathcal{D}), [\mathcal{D}, a] \text{ has bounded closure}\}$ is dense in \mathfrak{A} .

If $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is a spectral triple and if for any $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$, we set:

$$\text{mk}_{\mathcal{D}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), ||| [D, a] |||_{\mathcal{H}} \leq 1 \}$$

then $\text{mk}_{\mathcal{D}}$ is an extended pseudo-distance on $\mathcal{S}(\mathfrak{A})$.

A metric condition

We want that $\text{mk}_{\mathcal{D}}$ induces the weak* topology on $\mathcal{S}(\mathfrak{A})$.

Standard description of a noncommutative geometry

Definition (Connes, 89)

A **spectral triple** $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is given by:

- ① a unital C^* -algebra \mathfrak{A} ,
- ② a Hilbert space \mathcal{H} on which \mathfrak{A} acts,
- ③ a self-adjoint operator \mathcal{D} on $\text{dom}(\mathcal{D}) \subseteq \mathcal{H}$, such that:
 - $\mathcal{D} \pm i$ has a compact inverse,
 - $\mathfrak{A}^1 = \{a \in \mathfrak{A} : a \cdot \text{dom}(\mathcal{D}) \subseteq \text{dom}(\mathcal{D}), [\mathcal{D}, a] \text{ has bounded closure}\}$ is dense in \mathfrak{A} .

Definition

A **metric spectral triple** $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is a spectral triple such that, if

$$\forall a \in \mathfrak{sa}(\mathfrak{A}^1)$$

$$\mathcal{L}_{\mathcal{D}}(a) = ||| [\mathcal{D}, a] |||_{\mathcal{H}}$$

then $(\mathfrak{A}, \mathcal{L}_{\mathcal{D}})$ is a **quantum compact metric space**.

Ingredients for Convergence of Spectral Triples

Let $(\mathfrak{A}, \mathcal{H}, D)$ be a spectral triple.

A Metric

Connes proposed a noncommutative version of the *Monge-Kantorovich metric* on the state space $\mathcal{S}(\mathfrak{A})$:

$$\mathrm{mk}_D : \varphi, \psi \in \mathcal{S}(\mathfrak{A}) \mapsto \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), ||| [D, a] |||_{\mathcal{H}} \leq 1 \}.$$

Ingredients for Convergence of Spectral Triples

Let $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ be a spectral triple.

A Metric

Connes proposed a noncommutative version of the *Monge-Kantorovich metric* on the state space $\mathcal{S}(\mathfrak{A})$:

$$\mathrm{mk}_{\mathcal{D}} : \varphi, \psi \in \mathcal{S}(\mathfrak{A}) \mapsto \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), ||| [\mathcal{D}, a] |||_{\mathcal{H}} \leq 1 \}.$$

A Sobolev Space

The space $\mathrm{dom}(\mathcal{D})$ is a Banach space for the *graph norm*

$$\xi \in \mathrm{dom}(\mathcal{D}) \mapsto \mathsf{D}(\xi) = \|\xi\|_{\mathcal{H}} + \|\mathcal{D}\xi\|_{\mathcal{H}}.$$

Ingredients for Convergence of Spectral Triples

Let $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ be a spectral triple.

A Metric

Connes proposed a noncommutative version of the *Monge-Kantorovich metric* on the state space $\mathcal{S}(\mathfrak{A})$:

$$\mathrm{mk}_{\mathcal{D}} : \varphi, \psi \in \mathcal{S}(\mathfrak{A}) \mapsto \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), ||| [\mathcal{D}, a] |||_{\mathcal{H}} \leq 1 \}.$$

A Sobolev Space

The space $\mathrm{dom}(\mathcal{D})$ is a Banach space for the *graph norm*

$$\xi \in \mathrm{dom}(\mathcal{D}) \mapsto \mathsf{D}(\xi) = \|\xi\|_{\mathcal{H}} + \|\mathcal{D}\xi\|_{\mathcal{H}}.$$

A Quantum Dynamics

Since \mathcal{D} is self-adjoint, it gives rise to an *action of \mathbb{R} by unitaries* of \mathcal{H} :

$$t \in \mathbb{R} \longmapsto \exp(2i\pi t \mathcal{D}).$$

D-norms

The Hilbert space \mathcal{H} of a spectral triple $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ is a \mathfrak{A} -module and a \mathbb{C} -Hilbert module.

Next question: modules

How far are modules over quantum compact metric spaces?

D-norms

The Hilbert space \mathcal{H} of a spectral triple $(\mathfrak{A}, \mathcal{H}, D)$ is a \mathfrak{A} -module and a \mathbb{C} -Hilbert module.

Next question: modules

How far are modules over quantum compact metric spaces?

To this end we need a key ingredient: the graph norm of D

$$\forall \xi \in \text{dom}(D) \quad D_D(\xi) = \|\xi\|_{\mathcal{H}} + \|D\xi\|_{\mathcal{H}}$$

Theorem (L., 2018,9)

If $(\mathfrak{A}, \mathcal{H}, D)$ is a metric spectral triple and D_D is the graph norm of D then:

$$\{\xi \in \mathcal{H} : D_D(\xi) \leq 1\} \text{ is compact in } \mathcal{H},$$

while $D_D(a\xi) \leq (L_D(a) + \|a\|_{\mathfrak{A}})D_D(\xi)$ (with $L_D(a) = \| [D, a] \|_{\mathcal{H}}$).

Metrical C^* -correspondences

Definition (L. (16,18,19))

A *metrical C^* -correspondence* $\Omega = (\mathcal{M}, \mathsf{D}, \mathfrak{B}, \mathsf{L}_{\mathfrak{B}}, \mathfrak{A}, \mathsf{L}_{\mathfrak{A}})$ is given by:

- ❶ two *quantum compact metric spaces* $(\mathfrak{A}, \mathsf{L}_{\mathfrak{A}})$ and $(\mathfrak{B}, \mathsf{L}_{\mathfrak{B}})$,
- ❷ a *right C^* -Hilbert module* $(\mathcal{M}, \langle \cdot, \cdot \rangle_{\mathcal{M}})$ over \mathfrak{B} , which is also a *left module* over \mathfrak{A} , with the actions of \mathfrak{A} and \mathfrak{B} commuting,
- ❸ D is a *norm* on a dense subspace of \mathcal{M} such that:
 - ❶ $\mathsf{D} \geq \| \cdot \|_{\mathcal{M}} := \sqrt{\langle \cdot, \cdot \rangle_{\mathcal{M}}}$
 - ❷ $\{\omega \in \mathcal{M} : \mathsf{D}(\omega) \leq 1\}$ is compact in $(\mathcal{M}, \| \cdot \|_{\mathcal{M}})$,
 - ❸ $\forall \eta, \omega \in \mathcal{M} \quad \max\{\mathsf{L}_{\mathfrak{B}}(\Re \langle \omega, \eta \rangle_{\mathcal{M}}), \mathsf{L}_{\mathfrak{B}}(\Im \langle \omega, \eta \rangle_{\mathcal{M}})\} \leq H(\mathsf{D}(\omega), \mathsf{D}(\eta)),$
 - ❹ $\forall \eta \in \mathcal{M} \quad \forall a \in \mathfrak{sa}(\mathfrak{A}) \quad \mathsf{D}(a\eta) \leq G(\|a\|_{\mathfrak{A}}, \mathsf{L}_{\mathfrak{A}}(b), \mathsf{D}(\eta)).$

Metrical C^* -correspondences

Definition (L. (16,18,19))

A *metrical C^* -correspondence* $\Omega = (\mathcal{M}, \mathbb{D}, \mathfrak{B}, \mathbb{L}_{\mathfrak{B}}, \mathfrak{A}, \mathbb{L}_{\mathfrak{A}})$ is given by:

- ① two *quantum compact metric spaces* $(\mathfrak{A}, \mathbb{L}_{\mathfrak{A}})$ and $(\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$,
- ② a *right C^* -Hilbert module* $(\mathcal{M}, \langle \cdot, \cdot \rangle_{\mathcal{M}})$ over \mathfrak{B} , which is also a *left module* over \mathfrak{A} , with the actions of \mathfrak{A} and \mathfrak{B} commuting,
- ③ \mathbb{D} is a *norm* on a dense subspace of \mathcal{M} such that:
 - ① $\mathbb{D} \geq \| \cdot \|_{\mathcal{M}} := \sqrt{\langle \cdot, \cdot \rangle_{\mathcal{M}}}$
 - ② $\{\omega \in \mathcal{M} : \mathbb{D}(\omega) \leq 1\}$ is compact in $(\mathcal{M}, \| \cdot \|_{\mathcal{M}})$,
 - ③ $\forall \eta, \omega \in \mathcal{M} \quad \max\{\mathbb{L}_{\mathfrak{B}}(\Re \langle \omega, \eta \rangle_{\mathcal{M}}), \mathbb{L}_{\mathfrak{B}}(\Im \langle \omega, \eta \rangle_{\mathcal{M}})\} \leq H(\mathbb{D}(\omega), \mathbb{D}(\eta)),$
 - ④ $\forall \eta \in \mathcal{M} \quad \forall a \in \mathfrak{sa}(\mathfrak{A}) \quad \mathbb{D}(a\eta) \leq G(\|a\|_{\mathfrak{A}}, \mathbb{L}_{\mathfrak{A}}(b), \mathbb{D}(\eta)).$

For our purpose, if $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$ is a *metric spectral triple*, then

$$\text{qvb}(\mathfrak{A}, \mathcal{H}, \mathbb{D}) = (\mathcal{H}, \mathbb{D}_{\mathbb{D}}, \mathbb{C}, 0, \mathfrak{A}, \mathbb{L}_{\mathbb{D}})$$

is a *metrical C^* -correspondence*.

Metrical Tunnels

Let $(\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1)$ and $(\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)$ be two metric spectral triples.




Figure: π_1, π_2 are quantum isometries

Metrical Tunnels

Let $(\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1)$ and $(\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)$ be two metric spectral triples.




Figure: π_1, π_2 are quantum isometries




Figure: $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$ are quantum isometries

We want (π_j, θ_j) to be a *module morphism* and $\mathbb{D}_j = \inf \mathbb{T} \circ \theta_j^{-1}$ for $j \in \{1, 2\}$.

Approximating metrical quantum vector bundles




Figure: A metrical modular tunnel between $(\mathfrak{A}_1, \mathcal{H}_1, \mathbf{D}_1)$ and $(\mathfrak{A}_2, \mathcal{H}_2, \mathbf{D}_2)$.

The modular propinquity

Definition (L.,16,18)

A *tunnel* τ from $\mathbb{A}_1 = (\mathcal{M}_1, \mathbf{D}_1, \mathfrak{A}_1, \mathbf{L}_{\mathfrak{A}}^1, \mathfrak{B}_1, \mathbf{L}_{\mathfrak{B}}^1)$ to $\mathbb{A}_2 = (\mathcal{M}_2, \mathbf{D}_2, \mathfrak{A}_2, \mathbf{L}_{\mathfrak{A}}^2, \mathfrak{B}_2, \mathbf{L}_{\mathfrak{B}}^2)$ is given by:

- a metrical C*-correspondence $(\mathcal{P}, \mathbf{D}, \mathfrak{D}, \mathbf{L}_{\mathfrak{D}}, \mathfrak{E}, \mathbf{L}_{\mathfrak{E}})$,
- a module morphism (θ_j, Θ_j) from \mathcal{P} to \mathcal{M}_j which preserves the inner product and such that the quotient of \mathbf{D} via Θ_j is \mathbf{D}_j , while θ_j is a quantum isometry,
- a quantum isometry $\pi_j : (\mathfrak{E}, \mathbf{L}_{\mathfrak{E}}) \rightarrow (\mathfrak{B}_j, \mathbf{L}_{\mathfrak{B}}^j)$ such that (π_j, Θ_j) is also a module morphism.

The modular propinquity

Definition (L.,16,18)

A *tunnel* τ from $\mathbb{A}_1 = (\mathcal{M}_1, \mathbb{D}_1, \mathfrak{A}_1, \mathbb{L}_{\mathfrak{A}}^1, \mathfrak{B}_1, \mathbb{L}_{\mathfrak{B}}^1)$ to $\mathbb{A}_2 = (\mathcal{M}_2, \mathbb{D}_2, \mathfrak{A}_2, \mathbb{L}_{\mathfrak{A}}^2, \mathfrak{B}_2, \mathbb{L}_{\mathfrak{B}}^2)$ is given by:

- a metrical C^* -correspondence $(\mathcal{P}, \mathbb{D}, \mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \mathfrak{E}, \mathbb{L}_{\mathfrak{E}})$,
- a module morphism (θ_j, Θ_j) from \mathcal{P} to \mathcal{M}_j which preserves the inner product and such that the quotient of \mathbb{D} via Θ_j is \mathbb{D}_j , while θ_j is a quantum isometry,
- a quantum isometry $\pi_j : (\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}) \rightarrow (\mathfrak{B}_j, \mathbb{L}_{\mathfrak{B}}^j)$ such that (π_j, Θ_j) is also a module morphism.

The extent of τ is the maximum of the extents of $(\mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \theta_1, \theta_2)$ and $(\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}, \pi_1, \pi_2)$.

The modular propinquity

Definition (L.,16,18)

A *tunnel* τ from $\mathbb{A}_1 = (\mathcal{M}_1, \mathbb{D}_1, \mathfrak{A}_1, \mathbb{L}_{\mathfrak{A}}^1, \mathfrak{B}_1, \mathbb{L}_{\mathfrak{B}}^1)$ to $\mathbb{A}_2 = (\mathcal{M}_2, \mathbb{D}_2, \mathfrak{A}_2, \mathbb{L}_{\mathfrak{A}}^2, \mathfrak{B}_2, \mathbb{L}_{\mathfrak{B}}^2)$ is given by:

- a metrical C^* -correspondence $(\mathcal{P}, \mathbb{D}, \mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \mathfrak{E}, \mathbb{L}_{\mathfrak{E}})$,
- a module morphism (θ_j, Θ_j) from \mathcal{P} to \mathcal{M}_j which preserves the inner product and such that the quotient of \mathbb{D} via Θ_j is \mathbb{D}_j , while θ_j is a quantum isometry,
- a quantum isometry $\pi_j : (\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}) \rightarrow (\mathfrak{B}_j, \mathbb{L}_{\mathfrak{B}}^j)$ such that (π_j, Θ_j) is also a module morphism.

The extent of τ is the maximum of the extents of $(\mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \theta_1, \theta_2)$ and $(\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}, \pi_1, \pi_2)$. The *propinquity* between \mathbb{A}_1 and \mathbb{A}_2 is the infimum of the extent of all possible tunnels between them.

The modular propinquity

Definition (L.,16,18)

A *tunnel* τ from $\mathbb{A}_1 = (\mathcal{M}_1, \mathbb{D}_1, \mathfrak{A}_1, \mathbb{L}_{\mathfrak{A}_1}^1, \mathfrak{B}_1, \mathbb{L}_{\mathfrak{B}_1}^1)$ to $\mathbb{A}_2 = (\mathcal{M}_2, \mathbb{D}_2, \mathfrak{A}_2, \mathbb{L}_{\mathfrak{A}_2}^2, \mathfrak{B}_2, \mathbb{L}_{\mathfrak{B}_2}^2)$ is given by:

- a metrical C*-correspondence $(\mathcal{P}, \mathbb{D}, \mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \mathfrak{E}, \mathbb{L}_{\mathfrak{E}})$,
- a module morphism (θ_j, Θ_j) from \mathcal{P} to \mathcal{M}_j which preserves the inner product and such that the quotient of \mathbb{D} via Θ_j is \mathbb{D}_j , while θ_j is a quantum isometry,
- a quantum isometry $\pi_j : (\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}) \rightarrow (\mathfrak{B}_j, \mathbb{L}_{\mathfrak{B}_j}^j)$ such that (π_j, Θ_j) is also a module morphism.

The extent of τ is the maximum of the extents of $(\mathfrak{D}, \mathbb{L}_{\mathfrak{D}}, \theta_1, \theta_2)$ and $(\mathfrak{E}, \mathbb{L}_{\mathfrak{E}}, \pi_1, \pi_2)$. The *propinquity* between \mathbb{A}_1 and \mathbb{A}_2 is the infimum of the extent of all possible tunnels between them.

Theorem (L., 16,18)

The modular propinquity is a complete metric on metrical C*-correspondence, up to full quantum isometry.

Application to metric spectral triples

Theorem (L., 19)

For any metric spectral triple $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$, we set:

$$\text{qvb}(\mathfrak{A}, \mathcal{H}, \mathbb{D}) = (\mathcal{H}, \mathbb{D}, \mathbb{C}, 0, \mathfrak{A}, \mathbb{L})$$

where

- $\mathbb{L}(a) = |||[\mathbb{D}, a]|||_{\mathcal{H}},$
- $\mathbb{D}(\xi) = \|\xi\|_{\mathcal{H}} + \|\mathbb{D}\xi\|_{\mathcal{H}}.$

The function $\Lambda^{\text{mod}}(\text{qvb}(\cdot), \text{qvb}(\cdot))$ is a pseudo-metric on metric spectral triples such that:

$$\Lambda^{\text{mod}}(\text{qvb}(\mathfrak{A}, \mathcal{H}_{\mathfrak{A}}, \mathbb{D}_{\mathfrak{A}}), \text{qvb}(\mathfrak{B}, \mathcal{H}_{\mathfrak{B}}, \mathbb{D}_{\mathfrak{B}})) = 0$$

if and only if there exists a full quantum isometry $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$ and a unitary $U : \mathcal{H}_{\mathfrak{A}} \rightarrow \mathcal{H}_{\mathfrak{B}}$ such that:

$$U\mathbb{D}_{\mathfrak{A}}^2 U^* = \mathbb{D}_{\mathfrak{B}}^2 \text{ and } \pi(a) = UaU^*.$$

Group actions

The phase of a Dirac operator

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

Group actions

The phase of a Dirac operator

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

If $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$ is a spectral triple then:

$$t \in \mathbb{R} \mapsto U_t = \exp(i t \mathbb{D})$$

is a strongly continuous action of \mathbb{R} on \mathcal{H} such that:

$$\forall t \in \mathbb{R}, \xi \in \mathcal{H} \quad \mathbb{D}(U_t \xi) = \|U_t \xi\|_{\mathcal{H}} + \|\mathbb{D} U_t \xi\|_{\mathcal{H}} = \mathbb{D}(\xi).$$

Group actions

The phase of a Dirac operator

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

If $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$ is a spectral triple then:

$$t \in \mathbb{R} \mapsto U_t = \exp(it\mathbb{D})$$

is a strongly continuous action of \mathbb{R} on \mathcal{H} such that:

$$\forall t \in \mathbb{R}, \xi \in \mathcal{H} \quad \mathbb{D}(U_t \xi) = \|U_t \xi\|_{\mathcal{H}} + \|\mathbb{D}U_t \xi\|_{\mathcal{H}} = \mathbb{D}(\xi).$$

Next step

How do we incorporate convergence of group actions?

A proper monoid Gromov-Hausdorff distance

Let (G_1, δ_1) and (G_2, δ_2) be two proper metric monoids. A *ε -isometric isomorphism* $(\varsigma_1, \varsigma_2)$ from (G_1, δ_1) to (G_2, δ_2) is a pair of functions $\varsigma_1 : G_1 \rightarrow G_2$ and $\varsigma_2 : G_2 \rightarrow G_1$ such that for all $\{j, k\} = \{1, 2\}$:

$$\forall g, g' \in G_j \left[\frac{1}{\varepsilon} \right], h \in G_k \left[\frac{1}{\varepsilon} \right]$$

$$|\delta_k(\varsigma_j(g)\varsigma_j(g'), h) - \delta_j(gg', \varsigma_k(h))| \leq \varepsilon,$$

and ς_1, ς_2 are unital.

A proper monoid Gromov-Hausdorff distance

Let (G_1, δ_1) and (G_2, δ_2) be two proper metric monoids. A *ε -isometric isomorphism* $(\varsigma_1, \varsigma_2)$ from (G_1, δ_1) to (G_2, δ_2) is a pair of functions $\varsigma_1 : G_1 \rightarrow G_2$ and $\varsigma_2 : G_2 \rightarrow G_1$ such that for all $\{j, k\} = \{1, 2\}$:

$$\forall g, g' \in G_j \left[\frac{1}{\varepsilon} \right], h \in G_k \left[\frac{1}{\varepsilon} \right]$$

$$|\delta_k(\varsigma_j(g)\varsigma_j(g'), h) - \delta_j(gg', \varsigma_k(h))| \leq \varepsilon,$$

and ς_1, ς_2 are unital.

We then set:

$$Y((G_1, \delta_1), (G_2, \delta_2)) = \inf \{ \varepsilon > 0 \mid \exists (\varsigma, \kappa) \text{ } \varepsilon\text{-iso-iso} \}.$$

Y is a *metric up to isometric isomorphism of proper monoid*, and it dominates the pointed Gromov-Hausdorff distance.

The Covariant Reach

Let (τ, τ') be a *metrical tunnel* from $(\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1)$ to $(\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)$.
Write $\tau = (\mathcal{P}, \text{TN}, (\Pi_1, \pi_2), (\Pi_2, \pi_2))$.

Let $\varepsilon > 0$. Let $(\varsigma_1, \varsigma_2)$ be an *iso-iso* from \mathbb{R} to \mathbb{R} .

Definition (L., 18)

The ε -covariant reach $\rho(\tau|\varepsilon)$ of $(\tau, \tau', \varsigma_1, \varsigma_2)$ is

$$\max_{\{j,k\}=\{1,2\}} \sup_{\substack{\xi \in \mathcal{H}_j \\ \mathbb{D}_j(\xi) \leq 1}} \inf_{\substack{\eta \in \mathcal{H}_k \\ \mathbb{D}_k(\eta) \leq 1}} \sup_{|t| \leq \frac{1}{\varepsilon}} k_{\text{TN}}(e^{it\mathbb{D}_j} \xi, e^{i\varsigma_j(t)\mathbb{D}_k} \eta),$$

where

$$k_{\text{TN}}(\xi, \eta) = \sup_{\substack{\omega \in \mathcal{P} \\ \text{TN}(\omega) \leq 1}} \left| \langle \xi, \Pi_1(\omega) \rangle_{\mathcal{H}_1} - \langle \eta, \Pi_2(\omega) \rangle_{\mathcal{H}_2} \right|.$$

The Spectral Propinquity

Definition (L., 18)

The *spectral propinquity* $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathcal{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathcal{D}_2))$ between two spectral triples is

$$\max \left\{ \frac{\sqrt{2}}{2}, \inf \{ \varepsilon > 0 : \exists \tau \in \Upsilon \quad \max \{ \chi(\tau), \rho(\tau|\varepsilon) \} \leq \varepsilon \} \right\}$$

where Υ is the set of all covariant tunnels between the given spectral triples (for the actions $t \in \mathbb{R} \mapsto e^{it\mathcal{D}_j}$).

The Spectral Propinquity

Definition (L., 18)

The *spectral propinquity* $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2))$ between two spectral triples is

$$\max \left\{ \frac{\sqrt{2}}{2}, \inf \{ \varepsilon > 0 : \exists \tau \in \Upsilon \quad \max \{ \chi(\tau), \rho(\tau|\varepsilon) \} \leq \varepsilon \} \right\}$$

where Υ is the set of all covariant tunnels between the given spectral triples (for the actions $t \in \mathbb{R} \mapsto e^{it\mathbb{D}_j}$).

Theorem (L., 18)

The following assertions are equivalent:

- ❶ $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)) = 0$,
- ❷ there exists a unitary $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$, and a $*$ -isomorphism $\theta : \mathfrak{A}_1 \rightarrow \mathfrak{A}_2$, such that

$$U\mathbb{D}_1U^* = \mathbb{D}_2 \text{ and } \theta = \text{Ad}(U).$$

- 1 *Introduction*
- 2 *Quantum Compact Metric Spaces*
- 3 *Convergence of Metric Spectral Triples*
- 4 *Examples*

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

The *Sierpiński gasket* is the attractor of the iterated function system $\{T_1, T_2, T_3\}$ with

$$T_j : x \in \mathbb{R}^2 \mapsto \frac{1}{2} (x + v_j)$$

with

$$v_1 = (0, 0), v_2 = (1, 0), v_3 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right).$$

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

The *Sierpiński gasket* can thus be seen as a limit, for the Hausdorff distance, of finite graphs, starting from the triangle $v_1 v_2 v_3$:



Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

Specifically, \mathcal{SG}_∞ is the limit, in the Hausdorff distance in \mathbb{R}^2 , of the sequence of graphs:

- 1 We let \mathcal{SG}_0 be the triangle $v_1 v_2 v_3$.
- 2 Let $\mathcal{SG}_{n+1} = T_1 \mathcal{SG}_n \cup T_2 \mathcal{SG}_n \cup T_3 \mathcal{SG}_n$ for all $n \in \mathbb{N}$.

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

Lapidus et al. constructed a spectral triple over the Sierpiński gasket. First, we define a spectral triple over $\mathfrak{CP} = \{f : [-1, 1] \mapsto \mathbb{C} : f(-1) = f(1)\}$ by setting $\mathcal{J} = L^2(\mathfrak{CP}, \lambda)$, on which \mathfrak{CP} acts by left multiplication, and ∂ is the closure of the linear map such that:

$$\forall k \in \mathbb{Z} \quad \partial \exp(i\pi k \cdot) = i\pi \left(k + \frac{1}{2}\right) \exp(i\pi k \cdot).$$

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

We then define a spectral triple over $C([0, 1])$ by identifying $C([0, 1])$ with a C^* -subalgebra of \mathfrak{CB} , sending $f \in C([0, 1])$ to $x \in [-1, 1] \mapsto f(|x|)$.

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

Let $n \in \mathbb{N} \cup \{\infty\}$ (and write $3^\infty = \infty$). Let $\mathcal{H}_n = \bigoplus_{k=0}^n \bigoplus_{j=1}^{3^k} \mathcal{J}$, and for $\xi = (\xi_{k,j})_{\substack{k=0,\dots,n \\ j=1,\dots,3^k}} \in \mathcal{H}_n$ with all components in $\text{dom}(\partial)$:

$$D_n \xi = \left(2^k \partial \xi_{k,j} \right)_{\substack{k=0,\dots,n \\ j=1,\dots,3^k}}.$$

$C(\mathcal{S}\mathcal{G}_\infty)$ acts on \mathcal{H}_n by multiplication by restrictions over edges in $\mathcal{S}\mathcal{G}_n$.

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

Theorem (Lapidus et al., 08)

For all $n \in \mathbb{N} \cup \{\infty\}$, the triple $(C(\mathcal{S}\mathcal{G}_\infty), \mathcal{H}_n, \mathbb{D}_n)$ is a metric spectral triple, whose metric restricts to the *geodesic* distance on $\mathcal{S}\mathcal{G}_n$.

Spectral Triple on Sierpiński gasket




Figure: The Sierpiński gasket

Theorem (Lapidus et al., 08)

For all $n \in \mathbb{N} \cup \{\infty\}$, the triple $(C(\mathcal{S}\mathcal{G}_\infty), \mathcal{H}_n, \mathbb{D}_n)$ is a metric spectral triple, whose metric restricts to the *geodesic* distance on $\mathcal{S}\mathcal{G}_n$.

Theorem (Landry, Lapidus, L., 20)

$$\lim_{n \rightarrow \infty} \Lambda^{\text{spec}}((C(\mathcal{S}\mathcal{G}_n), \mathcal{H}_n, \mathbb{D}_n), (C(\mathcal{S}\mathcal{G}_\infty), \mathcal{H}_\infty, \mathbb{D}_\infty)) = 0.$$

Spectral Triples for Fuzzy Tori (Informal)

One example of our work deals with the clock and shift matrices:

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & r_n & \ddots & \\ \vdots & \ddots & \ddots & \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & 1 & \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with $r_n = \exp\left(\frac{2i\pi}{n}\right)$.

This pair of matrices encodes a *dynamical system*.

Note that $S_n C_n S_n^* = r_n C_n$ and $C^*(C_n) = C(\mathrm{Sp}(C_n))$. Thus, $C^*(C_n, S_n)$ is the C^* -crossed product for the action of \mathbb{Z}_n on $\mathrm{Sp}(C_n)$ by rotation of angle $\exp\left(\frac{2i\pi}{n}\right)$.

Spectral Triples for Fuzzy Tori (Informal)

One example of our work deals with the clock and shift matrices:

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & r_n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 1 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with $r_n = \exp\left(\frac{2i\pi}{n}\right)$.

$C_n = \mathcal{F} S_n \mathcal{F}^*$ where \mathcal{F} is the *discrete Fourier transform*, so if C_n is a position observable, then S_n is the momentum (up to scaling).

Thus $n[C_n, \cdot]$ and $n[S_n, \cdot]$, acting on $C^*(C_n, S_n)$, are discrete analogues of a moving frame of \mathbb{T}^2 .

Spectral Triples for Fuzzy Tori (Informal)

One example of our work deals with the clock and shift matrices:

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & r_n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 1 & 0 \\ 1 & 0 & \cdots & \cdots & 0 \end{pmatrix},$$

with $r_n = \exp\left(\frac{2i\pi}{n}\right)$.

Let \mathcal{H}_n be the Hilbert space given by $M_n(\mathbb{C})$ with inner product

$$\forall \xi, \eta \in M_n(\mathbb{C}) \quad \langle \xi, \eta \rangle = \frac{1}{n} \text{Tr}(\xi^* \eta).$$

Of course, $C^*(C_n, S_n)$ acts on \mathcal{H}_n both the left, via multiplication, and on the right, via multiplication by the adjoint.

Spectral Triples for Fuzzy Tori (Informal)

One example of our work deals with the clock and shift matrices:

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & r_n & \ddots & \\ \vdots & \ddots & \ddots & \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & 1 & \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with $r_n = \exp\left(\frac{2i\pi}{n}\right)$.

We will study the spectral triple $(C^*(C_n, S_n), \mathcal{H} \otimes \mathbb{C}^4, \mathcal{D}_n)$, where

$$\mathcal{D}_n = \frac{n}{2} \left([C_n + C_n^*, \cdot] \otimes \gamma_1 + i[C_n - C_n^*, \cdot] \otimes \gamma_2 \right. \\ \left. + [S_n + S_n^*, \cdot] \otimes \gamma_3 + i[S_n^* - S_n, \cdot] \otimes \gamma_4 \right).$$

and $\gamma_j \gamma_s + \gamma_s \gamma_j = -21_{s=j}$ with $\gamma_1, \dots, \gamma_4$ are 4×4 matrices.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

$C^*(U, V)$ acts on $\ell^2(\mathbb{Z}^2)$ by

$$\forall \xi \in \ell^2(\mathbb{Z}^2) \quad U\xi : m \in \mathbb{Z}^2 \mapsto \xi(m - (1, 0)) \text{ and } V\xi : m \mapsto \xi(m - (0, 1)).$$

The natural Sobolev derivatives are given ($j \in \{1, 2\}$) by

$$\partial_j \xi : (m_1, m_2) \mapsto i m_j \partial_j,$$

wherever this gives an element of $\ell^2(\mathbb{Z}^2)$.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

The heuristics is that and element of the form $\sum_{j,k \in F} a_{j,k} C_n^j S_n^k$ in $C^*(C_n, S_n)$ is an approximation of an element of the form $\sum_{j,k \in F} a_{j,k} U^j V^k$, and then:

$$\begin{aligned} n \left[S_n, \sum a_{j,k} C_n^j S_n^k \right] &= \frac{S_n \left(\sum a_{j,k} C_n^j S_n^k \right) S_n^* - \sum a_{j,k} C_n^j S_n^k}{\frac{1}{n}} S_n \\ &\rightsquigarrow \partial_1 \left(\sum a_{j,k} U^j V^k \right) V. \end{aligned}$$

Similarly, $n[C_n, \cdot]$ is presumably analogue to $\partial_2(\cdot)U$.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

We will work with the Dirac-type operator:

$$D = \Re(U)\partial_2 \otimes \gamma_1 + \Im(U)\partial_2 \otimes \gamma_2 + \Re(V)\partial_1 \otimes \gamma_3 + \Im(V)\partial_1 \otimes \gamma_4.$$

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

We will work with the Dirac-type operator:

$$D = \Re(U)\partial_2 \otimes \gamma_1 + \Im(U)\partial_2 \otimes \gamma_2 + \Re(V)\partial_1 \otimes \gamma_3 + \Im(V)\partial_1 \otimes \gamma_4.$$

We use $\Re(a) = \frac{a+a^*}{2}$, $\Im(a) = \frac{a-a^*}{2i}$, and 4×4 matrices $\gamma_1, \dots, \gamma_4$ such that $\gamma_j \gamma_k + \gamma_k \gamma_j = 1_{j=k}$. This operator is densely defined, self-adjoint, with compact resolvent, over a dense subspace of $\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^4$.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

We will work with the Dirac-type operator:

$$D = \Re(U)\partial_2 \otimes \gamma_1 + \Im(U)\partial_2 \otimes \gamma_2 + \Re(V)\partial_1 \otimes \gamma_3 + \Im(V)\partial_1 \otimes \gamma_4.$$

For all $f \in C(\mathbb{T}^2)$ such that $f \cdot \text{dom}(D) \subseteq \text{dom}(D)$ and $[D, f \otimes 1_{\mathbb{C}^2}]$ is bounded, we set

$$L(f) = |||[D, f \otimes 1_{\mathbb{C}^4}]]|||_{\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^4}.$$

$(C(\mathbb{T}^2), \ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^4, D)$ is a spectral triple, and $(C(\mathbb{T}^2), L)$ is a quantum compact metric space.

Case Study: Spectral Triples on Quantum Tori (Informal)

Our work proposes, in particular, to prove that the *spectral triples on $C^*(C_n, S_n)$ converge to some spectral triple on $C(\mathbb{T}^2)$* . Write $C(\mathbb{T}^2) = C^*(U, V)$ with $VU = UV$, and U, V unitaries.

We will work with the Dirac-type operator:

$$D = \Re(U)\partial_2 \otimes \gamma_1 + \Im(U)\partial_2 \otimes \gamma_2 + \Re(V)\partial_1 \otimes \gamma_3 + \Im(V)\partial_1 \otimes \gamma_4.$$

Theorem (L., 21)

$$\lim_{n \rightarrow \infty} \Lambda^{\text{spec}} \left((C^*(C_n, S_n), \mathcal{H}_n \otimes \mathbb{C}^4, D_n), (C(\mathbb{T}^2), \ell^2(\mathbb{Z}^{2d}, \mathbb{C}^4), D) \right) = 0.$$

Thank you!

- *The Quantum Gromov-Hausdorff Propinquity*, F. Latrémolière, *Transactions of the AMS* **368** (2016) 1, pp. 365–411, ArXiv: 1302.4058
- *Convergence of Fuzzy Tori and Quantum Tori for the quantum Gromov-Hausdorff Propinquity: an explicit Approach*, F. Latrémolière, *Münster Journal of Mathematics* **8** (2015) 1, pp. 57–98, ArXiv: 1312.0069
- *The Dual Gromov-Hausdorff Propinquity*, F. Latrémolière, *Journal de Mathématiques Pures et Appliquées* **103** (2015) 2, pp. 303–351, ArXiv: 1311.0104
- *A compactness theorem for the dual Gromov-Hausdorff Propinquity*, F. Latrémolière, *Indiana Univ. Math. J.* **66** (2017) 5, 1707–1753, Arxiv: 1501.06121
- *The modular Gromov-Hausdorff propinquity*, F. Latrémolière, *Dissertationes Math.* 544 (2019), 70 pp. 46L89 (46L30 58B34)
- *The dual modular propinquity and completeness*, F. Latrémolière, accepted in *J. Noncomm. Geometry* (2020), arXiv:1811.04534
- *The Gromov-Hausdorff Propinquity for Metric Spectral Triples*, F. Latrémolière, submitted (2019), arXiv:1811.10843.
- *Convergence of Spectral Triples on Fuzzy Tori to Spectral Triples on Quantum Tori*, F. Latrémolière, submitted, arXiv:2102.03729.