

# *A Geometry for the space of spectral triples*

Frédéric Latrémolière



*Gravity, Noncommutative Geometry, Cosmology*

*BIRS-CMO*

*August, 21<sup>st</sup> 2023.*

# The Space of Noncommutative “Drums”

A. Connes introduced in 1985 a generalization of *spectral geometry* to the *noncommutative realm* by means of a structure called a *spectral triple*, i.e. a noncommutative analogue of the Dirac operators on Riemannian spin manifolds.



# The Space of Noncommutative Manifolds

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$  is an analogue of a first order pseudo-differential operator (of Dirac type), where

- $\mathfrak{A}$  is a *unital  $C^*$ -algebra*, i.e. a noncommutative analogue of  $C(X)$  for a compact Hausdorff space  $X$ ,
- $\mathcal{H}$  is a *Hilbert space* on which  $\mathfrak{A}$  acts, analogue to the space of sections of some spinor bundles,
- $\mathcal{D}$  is a *self-adjoint operator* densely defined on  $\mathcal{H}$  with *compact resolvent* and *bounded commutator* with a dense  $*$ -subalgebra of  $\mathfrak{A}$ ; formally

$$\mathfrak{A}_{\mathcal{D}} = \{a \in \mathfrak{A} : a \operatorname{dom}(\mathcal{D}) \subseteq \operatorname{dom}(\mathcal{D}) \text{ and } [\mathcal{D}, a] \text{ is bounded}\}$$


is a dense  $*$ -subalgebra of  $\mathfrak{A}$ .



*Figure:* The spectrum of oxygen

## The Space of Noncommutative Manifolds


Spectral triples are *much more flexible* than manifold structure, and can be constructed over *noncommutative  $C^*$ -algebras*, fractals and other singular spaces, and even finite sets. Much success has been achieved in extending Atiah-Singer index theorem to spectral triples to many such generalized setting.



*Figure:* The Sierpiński Triangle

## The Space of Noncommutative Manifolds

Spectral triples are *much more flexible* than manifold structure, and can be constructed over *noncommutative  $C^*$ -algebras*, fractals and other singular spaces, and even finite sets. Much success has been achieved in extending Atiah-Singer index theorem to spectral triples to many such generalized setting.



*Figure:* The Sierpiński Triangle

Spectral triples allow us to extend important notions from physics. The *spectral action* of a spectral triple  $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$ , in particular, generalizes the Hilbert action of relativity, and is relevant to physics over noncommutative spaces.

## Matrix Models: Clock and Shift, Fuzzy Tori

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ & r_n & \ddots & \\ & & \ddots & 0 \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \ddots & \\ 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 \end{pmatrix},$$

with  $r_n = \exp\left(\frac{2i\pi}{n}\right)$ . Note that  $C^*(C_n, S_n) = \mathfrak{B}(\mathbb{C}^n)$ .

## Matrix Models: Clock and Shift, Fuzzy Tori

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ & r_n & \ddots & \\ & & \ddots & 0 \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \ddots & \\ & & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1 & \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with  $r_n = \exp\left(\frac{2i\pi}{n}\right)$ . Note that  $C^*(C_n, S_n) = \mathfrak{B}(\mathbb{C}^n)$ . Let  $\mathbb{D}_n$  be defined on  $\mathcal{H}_n = \mathfrak{B}(\mathbb{C}^n) \otimes \mathbb{C}^4$  be defined by:

$$\begin{aligned} \mathbb{D}_n = \frac{n}{2\pi} & \left( \left[ \frac{S_n + S_n^*}{2}, \cdot \right] \otimes \gamma_1 + \left[ \frac{S_n - S_n^*}{2i}, \cdot \right] \otimes \gamma_2 \right. \\ & \left. + \left[ \frac{C_n^* + C_n}{2}, \cdot \right] \otimes \gamma_3 + \left[ \frac{C_n^* - C_n}{2i}, \cdot \right] \otimes \gamma_4 \right), \end{aligned}$$

with  $\gamma_j \gamma_k + \gamma_k \gamma_j = \delta_j^k$  ( $j, k \in \{1, 2, 3, 4\}$ ). Then  $(C^*(C_n, S_n), \mathcal{H}_n, \mathbb{D}_n)$  is a spectral triple.

# *A geometry for the space of quantum spaces*

## *The Questions*

- ① How do we formalize convergence of spectral triples?
- ② Can we construct new spectral triples as limits?
- ③ Can we work with entire spaces of spectral triples?



# *A geometry for the space of quantum spaces*

## *The Questions*

- ❶ How do we formalize convergence of spectral triples?
- ❷ Can we construct new spectral triples as limits?
- ❸ Can we work with entire spaces of spectral triples?

## *Our Project*

We develop a geometric framework to study the *space of (metric) spectral triples as a natural metric space* to discuss problems from mathematical physics and functional analysis.

# *A geometry for the space of quantum spaces*

## *The Questions*

- ❶ How do we formalize convergence of spectral triples?
- ❷ Can we construct new spectral triples as limits?
- ❸ Can we work with entire spaces of spectral triples?

## *Our Project*

We develop a geometric framework to study the *space of (metric) spectral triples as a natural metric space* to discuss problems from mathematical physics and functional analysis.

For example: is there a limit for the spectral triples associated with clock/shift matrix models? Are spectral triples over fractals limits of spectral triples over simpler curves? Are spectral triples over certain inductive limits natural limits of spectral triples on the factors?

## *Our approach*

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  induces three structures relevant to our goal:

## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$  induces three structures relevant to our goal:

- 1 *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathcal{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathcal{D}), \|[\mathcal{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathcal{D})$  induces three structures relevant to our goal:

- ① *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathcal{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathcal{D}) \subseteq \mathrm{dom}(\mathcal{D}), \|[\mathcal{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

Our focus is on *metric spectral triples*, whose Connes' metric induces the weak\* topology;

## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  induces three structures relevant to our goal:

- 1 *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathbb{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathbb{D}) \subseteq \mathrm{dom}(\mathbb{D}), \|[\mathbb{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

Our focus is on *metric spectral triples*, whose Connes' metric induces the weak\* topology;

- 2 An  $\mathfrak{A}$ - $\mathbb{C}$   $C^*$ -correspondence  $\mathcal{H}$ ;

## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  induces three structures relevant to our goal:

- 1 *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathbb{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathbb{D}) \subseteq \mathrm{dom}(\mathbb{D}), \|[\mathbb{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

Our focus is on *metric spectral triples*, whose Connes' metric induces the weak\* topology;

- 2 An  $\mathfrak{A}$ - $\mathbb{C}$   $C^*$ -correspondence  $\mathcal{H}$ ;
- 3 An action of  $[0, \infty)$  on  $\mathcal{H}$  by unitaries, given by  $t \in [0, \infty) \mapsto \exp(it\mathbb{D})$ .

## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  induces three structures relevant to our goal:

- 1 *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathbb{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathbb{D}) \subseteq \mathrm{dom}(\mathbb{D}), \|[\mathbb{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

Our focus is on *metric spectral triples*, whose Connes' metric induces the weak\* topology;

- 2 An  $\mathfrak{A}$ - $\mathbb{C}$   $C^*$ -correspondence  $\mathcal{H}$ ;
- 3 An action of  $[0, \infty)$  on  $\mathcal{H}$  by unitaries, given by  $t \in [0, \infty) \mapsto \exp(it\mathbb{D})$ .

We aim at devising a metric on a large space of spectral triples for which important properties of spectral triples — such as spectrum and spectral actions — are continuous.



## Our approach

A *spectral triple*  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  induces three structures relevant to our goal:

- 1 *Connes' metric* on the state space of  $\mathfrak{A}$ : Connes' distance between any two states  $\varphi, \psi \in \mathcal{S}(\mathfrak{A})$  of  $\mathfrak{A}$  is:

$$\mathrm{mk}_{\mathbb{L}}(\varphi, \psi) := \sup \{ |\varphi(a) - \psi(a)| : a \in \mathrm{dom}(\mathbb{D}) \subseteq \mathrm{dom}(\mathbb{D}), \|[\mathbb{D}, a]\|_{\mathcal{H}} \leq 1 \};$$

Our focus is on *metric spectral triples*, whose Connes' metric induces the weak\* topology;

- 2 An  $\mathfrak{A}$ - $\mathbb{C}$   $C^*$ -correspondence  $\mathcal{H}$ ;
- 3 An action of  $[0, \infty)$  on  $\mathcal{H}$  by unitaries, given by  $t \in [0, \infty) \mapsto \exp(it\mathbb{D})$ .

We aim at devising a metric on a large space of spectral triples for which important properties of spectral triples — such as spectrum and spectral actions — are continuous.

The core of our work is noncommutative metric geometry: *metric spectral triples are examples of compact quantum metric spaces*, for which we devised an analogue of the Gromov-Hausdorff distance.

- 1 *Compact Quantum Metric Spaces*
- 2 *The Gromov-Hausdorff Propinquity*
- 3 *The Spectral Propinquity*
- 4 *Examples and Applications*

## The Monge-Kantorovich metric

Let  $(X, \mathfrak{m})$  be a compact metric space. The *Lipschitz seminorm*  $\mathsf{L}$  induced by  $\mathfrak{m}$  is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all  $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$  (allowing  $\infty$ ).

## The Monge-Kantorovich metric

Let  $(X, \mathfrak{m})$  be a compact metric space. The *Lipschitz seminorm*  $\mathsf{L}$  induced by  $\mathfrak{m}$  is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all  $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$  (allowing  $\infty$ ).

The *Monge-Kantorovich metric* on  $\mathcal{S}(C(X))$  is given for all Borel-regular probability measures  $\mu, \nu$  by:

$$\mathsf{mk}_{\mathsf{L}}(\mu, \nu) = \sup \left\{ \left| \int_X f d\mu - \int_X f d\nu \right| : f \in \mathfrak{sa}(C(X)), \mathsf{L}(f) \leq 1 \right\}.$$

## The Monge-Kantorovich metric

Let  $(X, \mathfrak{m})$  be a compact metric space. The *Lipschitz seminorm*  $\mathsf{L}$  induced by  $\mathfrak{m}$  is:

$$\mathsf{L}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\mathfrak{m}(x, y)} : x, y \in X, x \neq y \right\}$$

for all  $f \in \mathfrak{sa}(C(X)) = C(X, \mathbb{R})$  (allowing  $\infty$ ).

The *Monge-Kantorovich metric* on  $\mathcal{S}(C(X))$  is given for all Borel-regular probability measures  $\mu, \nu$  by:

$$\mathsf{mk}_{\mathsf{L}}(\mu, \nu) = \sup \left\{ \left| \int_X f d\mu - \int_X f d\nu \right| : f \in \mathfrak{sa}(C(X)), \mathsf{L}(f) \leq 1 \right\}.$$

The Gelfand map  $x \in (X, \mathfrak{m}) \mapsto \delta_x \in (\mathcal{S}(C(X)), \mathsf{mk}_{\mathsf{L}})$  is an isometry.

# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathsf{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,

# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathbb{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,
- 2  $\mathbb{L}$  is a *seminorm* defined on a (dense) Jordan-Lie subalgebra  $\text{dom}(\mathbb{L})$  of  $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$ ,
- 3  $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathbb{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$ ,

# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathsf{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,
- 2  $\mathsf{L}$  is a *seminorm* defined on a (dense) Jordan-Lie subalgebra  $\text{dom}(\mathsf{L})$  of  $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$ ,
- 3  $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$ ,
- 4 The *weak\* topology on  $\mathcal{S}(\mathfrak{A})$*  is metrized by the *Monge-Kantorovich metric  $\text{mk}_{\mathsf{L}}$* , defined  $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$  by:

$$\text{mk}_{\mathsf{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathsf{L}(a) \leq 1 \}$$



# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathbb{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,
- 2  $\mathbb{L}$  is a *seminorm* defined on a (dense) Jordan-Lie subalgebra  $\text{dom}(\mathbb{L})$  of  $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$ ,
- 3  $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathbb{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$ ,
- 4 The *weak\* topology on  $\mathcal{S}(\mathfrak{A})$*  is metrized by the *Monge-Kantorovich metric  $\text{mk}_{\mathbb{L}}$* , defined  $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$  by:

$$\text{mk}_{\mathbb{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathbb{L}(a) \leq 1 \}$$

- 5  $\mathbb{L}\left(\frac{ab+ba}{2}\right) \vee \mathbb{L}\left(\frac{ab-ba}{2i}\right) \leq F(\|a\|_{\mathfrak{A}}\mathbb{L}(b) + \mathbb{L}(a)\|b\|_{\mathfrak{A}}) + K\mathbb{L}(a)\mathbb{L}(b);$

# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathbb{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,
- 2  $\mathbb{L}$  is a *seminorm* defined on a (dense) Jordan-Lie subalgebra  $\text{dom}(\mathbb{L})$  of  $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$ ,
- 3  $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathbb{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$ ,
- 4 The *weak\* topology on  $\mathcal{S}(\mathfrak{A})$*  is metrized by the *Monge-Kantorovich metric  $\text{mk}_{\mathbb{L}}$* , defined  $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$  by:

$$\text{mk}_{\mathbb{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathbb{L}(a) \leq 1 \}$$

- 5  $\mathbb{L}\left(\frac{ab+ba}{2}\right) \vee \mathbb{L}\left(\frac{ab-ba}{2i}\right) \leq F(\|a\|_{\mathfrak{A}}\mathbb{L}(b) + \mathbb{L}(a)\|b\|_{\mathfrak{A}}) + K\mathbb{L}(a)\mathbb{L}(b)$ ;
- 6  $\{a \in \text{dom}(\mathbb{L}) : \mathbb{L}(a) \leq 1\}$  is closed in  $\mathfrak{A}$ .

# Compact Quantum Metric Spaces

*Definition (Connes, 89; Rieffel, 98; L., 13)*

$(\mathfrak{A}, \mathsf{L})$  is a *quantum compact metric space* when:

- 1  $\mathfrak{A}$  is a *unital  $C^*$ -algebra*,
- 2  $\mathsf{L}$  is a *seminorm* defined on a (dense) Jordan-Lie subalgebra  $\text{dom}(\mathsf{L})$  of  $\mathfrak{sa}(\mathfrak{A}) = \{a \in \mathfrak{A} : a^* = a\}$ ,
- 3  $\{a \in \mathfrak{sa}(\mathfrak{A}) : \mathsf{L}(a) = 0\} = \mathbb{R}1_{\mathfrak{A}}$ ,
- 4 The *weak\* topology on  $\mathcal{S}(\mathfrak{A})$*  is metrized by the *Monge-Kantorovich metric  $\text{mk}_{\mathsf{L}}$* , defined  $\forall \varphi, \psi \in \mathcal{S}(\mathfrak{A})$  by:

$$\text{mk}_{\mathsf{L}}(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : a \in \mathfrak{sa}(\mathfrak{A}), \mathsf{L}(a) \leq 1 \}$$

- 5  $\mathsf{L}\left(\frac{ab+ba}{2}\right) \vee \mathsf{L}\left(\frac{ab-ba}{2i}\right) \leq F(\|a\|_{\mathfrak{A}}\mathsf{L}(b) + \mathsf{L}(a)\|b\|_{\mathfrak{A}}) + K\mathsf{L}(a)\mathsf{L}(b)$ ;
- 6  $\{a \in \text{dom}(\mathsf{L}) : \mathsf{L}(a) \leq 1\}$  is closed in  $\mathfrak{A}$ .

We call  $\mathsf{L}$  an  *$\mathsf{L}$ -seminorm*.

## Some facts about quantum compact metric spaces

- Quantum compact metric spaces form a *category* with arrows given by Lipschitz morphisms: a *Lipschitz morphism*  $\pi : (\mathfrak{A}, \mathsf{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathsf{L}_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(\mathsf{L}_{\mathfrak{A}})) \subseteq \text{dom}(\mathsf{L}_{\mathfrak{B}})$ ,

## Some facts about quantum compact metric spaces

- Quantum compact metric spaces form a *category* with arrows given by Lipschitz morphisms: a *Lipschitz morphism*  $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(\mathbb{L}_{\mathfrak{A}})) \subseteq \text{dom}(\mathbb{L}_{\mathfrak{B}})$ ,
- Every compact metric space  $(X, d)$  gives us a quantum compact metric space  $(C(X), \mathbb{L}_d)$ ,

## Some facts about quantum compact metric spaces

- Quantum compact metric spaces form a *category* with arrows given by Lipschitz morphisms: a *Lipschitz morphism*  $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(\mathbb{L}_{\mathfrak{A}})) \subseteq \text{dom}(\mathbb{L}_{\mathfrak{B}})$ ,
- Every compact metric space  $(X, d)$  gives us a quantum compact metric space  $(C(X), \mathbb{L}_d)$ ,
- Every finite dimensional space can be made into a quantum compact metric space,

## Some facts about quantum compact metric spaces

- Quantum compact metric spaces form a *category* with arrows given by Lipschitz morphisms: a *Lipschitz morphism*  $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(\mathbb{L}_{\mathfrak{A}})) \subseteq \text{dom}(\mathbb{L}_{\mathfrak{B}})$ ,
- Every compact metric space  $(X, d)$  gives us a quantum compact metric space  $(C(X), \mathbb{L}_d)$ ,
- Every finite dimensional space can be made into a quantum compact metric space,
- Examples include *quantum tori* (Rieffel, 98, 02; L., 21), *AF algebras* (Aguilar-L., 15) and other *inductive limits* (Aguilar-L., 22; Farsi-L-Packer, 23), *Podles spheres* and  $SU_q(2)$  (Aguilar-Kaad-Kyed, Kaad-Kyed, 19-22), some  *$C^*$ -crossed-products* (Hawkins-Skalski-White-Zacharias, 13; Klisse, 23), many *discrete group  $C^*$ -algebras* (Rieffel-Ozawa, 01; Rieffel-Christ, 15), and *more*.

# Metric Spectral Triples

## Definition (L., 18)

A spectral triple  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  is *metric* when  $\mathrm{mk}_{\mathbb{D}}$  is a metric on the state space  $\mathcal{S}(\mathfrak{A})$ , which induces the *weak\* topology*.



# Metric Spectral Triples

## Definition (L., 18)

A spectral triple  $(\mathfrak{A}, \mathcal{H}, D)$  is *metric* when  $\mathrm{mk}_D$  is a metric on the state space  $\mathcal{S}(\mathfrak{A})$ , which induces the *weak\* topology*.

In other words,  $(\mathfrak{A}, \mathcal{H}, D)$  is a metric spectral triple, if and only if, setting:

$$\mathrm{dom}(\mathbb{L}_D) = \{a \in \mathfrak{sa}(\mathfrak{A}) : a \cdot \mathrm{dom}(D) \subseteq \mathrm{dom}(D), [D, a] \text{ bounded}\}$$

and, for all  $a \in \mathrm{dom}(\mathbb{L}_D)$ ,

$$\mathbb{L}_D(a) = |||[D, a]|||_{\mathcal{H}},$$

then  $(\mathfrak{A}, \mathbb{L}_D)$  is a compact quantum metric space.

- 1 *Compact Quantum Metric Spaces*
- 2 *The Gromov-Hausdorff Propinquity*
- 3 *The Spectral Propinquity*
- 4 *Examples and Applications*

# The Gromov-Hausdorff Distance

The *Hausdorff distance*  $\text{Haus}_d$  between two closed subsets  $A_1$  and  $A_2$  of a compact metric space  $(X, d)$  is defined by

$$\text{Haus}_d(A_1, A_2) = \max_{\{j,k\}=\{1,2\}} \sup_{x \in A_j} \inf_{y \in A_k} d(x, y).$$

# The Gromov-Hausdorff Distance

The *Hausdorff distance*  $\text{Haus}_d$  between two closed subsets  $A_1$  and  $A_2$  of a compact metric space  $(X, d)$  is defined by

$$\text{Haus}_d(A_1, A_2) = \max_{\{j,k\}=\{1,2\}} \sup_{x \in A_j} \inf_{y \in A_k} d(x, y).$$

*Definition (Hausdorff, 1903; Edwards, 75; Gromov, 81)*

The *Gromov-Hausdorff distance* between two compact metric spaces  $(X, m_X)$  and  $(Y, m_Y)$  is:

$$\inf \left\{ \text{Haus}_{m_Z}(\iota_X(X), \iota_Y(Y)) \left| \begin{array}{l} (Z, m_Z) \text{ compact metric space,} \\ \iota_X : X \hookrightarrow Z \text{ isometry,} \\ \iota_Y : Y \hookrightarrow Z \text{ isometry.} \end{array} \right. \right\},$$

where  $\text{Haus}_d$  is the *Hausdorff distance* induced by a metric  $d$ .

## The Gromov-Hausdorff Distance

The *Hausdorff distance*  $\text{Haus}_d$  between two closed subsets  $A_1$  and  $A_2$  of a compact metric space  $(X, d)$  is defined by

$$\text{Haus}_d(A_1, A_2) = \max_{\{j,k\}=\{1,2\}} \sup_{x \in A_j} \inf_{y \in A_k} d(x, y).$$

*Definition (Hausdorff, 1903; Edwards, 75; Gromov, 81)*

The *Gromov-Hausdorff distance* between two compact metric spaces  $(X, m_X)$  and  $(Y, m_Y)$  is:

$$\inf \left\{ \text{Haus}_{m_Z}(\iota_X(X), \iota_Y(Y)) \left| \begin{array}{l} (Z, m_Z) \text{ compact metric space,} \\ \iota_X : X \hookrightarrow Z \text{ isometry,} \\ \iota_Y : Y \hookrightarrow Z \text{ isometry.} \end{array} \right. \right\},$$

where  $\text{Haus}_d$  is the *Hausdorff distance* induced by a metric  $d$ .

The *Gromov-Hausdorff distance* is a *complete metric*, up to isometry, on the class of compact metric spaces.

## Quantum Isometries

A *Lipschitz morphism*  $\pi : (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(\mathbb{L}_{\mathfrak{A}})) \subseteq \text{dom}(\mathbb{L}_{\mathfrak{B}})$ .

## Quantum Isometries

A *Lipschitz morphism*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$ .

*Definition (Rieffel (99), L. (13) )*

A *quantum isometry*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a  $*$ -epimorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$  and

$$\forall b \in \text{dom}(L_{\mathfrak{B}}) \quad L_{\mathfrak{B}}(b) = \inf \{ L_{\mathfrak{A}}(a) : \pi(a) = b \}.$$

A *full quantum isometry*  $\pi$  is a  $*$ -isomorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) = \text{dom}(L_{\mathfrak{B}})$  and  $L_{\mathfrak{B}} \circ \pi = L_{\mathfrak{A}}$ .

## Quantum Isometries

A *Lipschitz morphism*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$ .

*Definition (Rieffel (99), L. (13) )*

A *quantum isometry*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a  $*$ -epimorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$  and

$$\forall b \in \text{dom}(L_{\mathfrak{B}}) \quad L_{\mathfrak{B}}(b) = \inf \{ L_{\mathfrak{A}}(a) : \pi(a) = b \}.$$

A *full quantum isometry*  $\pi$  is a  $*$ -isomorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) = \text{dom}(L_{\mathfrak{B}})$  and  $L_{\mathfrak{B}} \circ \pi = L_{\mathfrak{A}}$ .

*Theorem (Rieffel, 99)*

If  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a quantum isometry, then  $\pi^* : \varphi \in \mathcal{S}(\mathfrak{B}) \mapsto \varphi \circ \pi \in \mathcal{S}(\mathfrak{A})$  is an isometry from  $(\mathcal{S}(\mathfrak{B}), \text{mk}_{L_{\mathfrak{B}}})$  into  $(\mathcal{S}(\mathfrak{A}), \text{mk}_{L_{\mathfrak{A}}})$ .



## Quantum Isometries

A *Lipschitz morphism*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a unital  $*$ -morphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$ .

*Definition (Rieffel (99), L. (13) )*

A *quantum isometry*  $\pi : (\mathfrak{A}, L_{\mathfrak{A}}) \rightarrow (\mathfrak{B}, L_{\mathfrak{B}})$  is a  $*$ -epimorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) \subseteq \text{dom}(L_{\mathfrak{B}})$  and


$$\forall b \in \text{dom}(L_{\mathfrak{B}}) \quad L_{\mathfrak{B}}(b) = \inf \{ L_{\mathfrak{A}}(a) : \pi(a) = b \}.$$

A *full quantum isometry*  $\pi$  is a  $*$ -isomorphism such that  $\pi(\text{dom}(L_{\mathfrak{A}})) = \text{dom}(L_{\mathfrak{B}})$  and  $L_{\mathfrak{B}} \circ \pi = L_{\mathfrak{A}}$ .

*Theorem (L., 18)*


If  $(\mathfrak{A}_1, \mathcal{H}_1, D_1)$  and  $(\mathfrak{A}_2, \mathcal{H}_2, D_2)$  are two *unitarily equivalent metric spectral triples*, then  $(\mathfrak{A}_1, L_{D_1})$  and  $(\mathfrak{A}_2, L_{D_2})$  are *fully quantum isometric*.

# The Dual Gromov-Hausdorff Propinquity



*Figure:*  $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$  are quantum isometries

# The Dual Gromov-Hausdorff Propinquity



*Figure:*  $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$  are quantum isometries

*Definition (The extent of a tunnel, L. 13,14)*

The *extent*  $\chi(\tau)$  of a tunnel  $\tau = (\mathfrak{D}, L_{\mathfrak{D}}, \pi_{\mathfrak{A}}, \pi_{\mathfrak{B}})$  is:

$$\max \left\{ \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left( \mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{A}}^* (\mathcal{S}(\mathfrak{A})) \right), \right. \\ \left. \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left( \mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{B}}^* (\mathcal{S}(\mathfrak{B})) \right) \right\}.$$

# The Dual Gromov-Hausdorff Propinquity




Figure:  $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$  are quantum isometries

## Definition (The Dual Propinquity, L. 13, 14)

The *extent*  $\chi(\tau)$  of a tunnel  $\tau = (\mathfrak{D}, L_{\mathfrak{D}}, \pi_{\mathfrak{A}}, \pi_{\mathfrak{B}})$  is:

$$\max \left\{ \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left( \mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{A}}^* (\mathcal{S}(\mathfrak{A})) \right), \right. \\ \left. \text{Haus}_{\text{mk}_{L_{\mathfrak{D}}}} \left( \mathcal{S}(\mathfrak{D}), \pi_{\mathfrak{B}}^* (\mathcal{S}(\mathfrak{B})) \right) \right\}.$$

The *dual propinquity*  $\Lambda^* ((\mathfrak{A}, L_{\mathfrak{A}}), (\mathfrak{B}, L_{\mathfrak{B}}))$  is given by:

$$\inf \{ \chi(\tau) : \tau \text{ any tunnel from } (\mathfrak{A}, L_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, L_{\mathfrak{B}}) \}.$$

# The Dual Gromov-Hausdorff Propinquity




Figure:  $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$  are quantum isometries

## Theorem (L., 13)

The *dual propinquity*  $\Lambda^*$ , defined for any two quantum compact metric spaces  $(\mathfrak{A}, L_{\mathfrak{A}})$  by  $(\mathfrak{B}, L_{\mathfrak{B}})$  by:

$$\inf \{ \chi(\tau) : \tau \text{ any tunnel from } (\mathfrak{A}, L_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, L_{\mathfrak{B}}) \}$$

is a *complete metric* up to *full quantum isometry*.  
 $\Lambda((\mathfrak{A}, L_{\mathfrak{A}}), (\mathfrak{B}, L_{\mathfrak{B}})) = 0$  iff there exists a  $*$ -isomorphism  $\pi : \mathfrak{A} \rightarrow \mathfrak{B}$  such that  $L_{\mathfrak{B}} \circ \pi = L_{\mathfrak{A}}$ .

# The Dual Gromov-Hausdorff Propinquity




Figure:  $\pi_{\mathfrak{A}}, \pi_{\mathfrak{B}}$  are quantum isometries

## Theorem (L., 13)

The *dual propinquity*  $\Lambda^*$ , defined for any two quantum compact metric spaces  $(\mathfrak{A}, \mathbb{L}_{\mathfrak{A}})$  by  $(\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  by:

$$\inf \{ \chi(\tau) : \tau \text{ any tunnel from } (\mathfrak{A}, \mathbb{L}_{\mathfrak{A}}) \text{ to } (\mathfrak{B}, \mathbb{L}_{\mathfrak{B}}) \}$$

is a *complete metric* up to *full quantum isometry*. Moreover  $\Lambda^*$  induces the topology of the *Gromov-Hausdorff distance* on compact metric spaces.

## *Some facts about the propinquity*

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.

## *Some facts about the propinquity*

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).



## *Some facts about the propinquity*

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).

## Some facts about the propinquity

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).
- *Quantum tori* (with various metrics) form a continuous family of quantum compact metric spaces in their deformation parameter for the propinquity and are limits of finite dimensional “fuzzy tori” (L., 2013, 2021).

## Some facts about the propinquity

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).
- *Quantum tori* (with various metrics) form a continuous family of quantum compact metric spaces in their deformation parameter for the propinquity and are limits of finite dimensional “fuzzy tori” (L., 2013, 2021).
- There are various naturally continuous families of AF algebras (*Effros-Shen algebras*, *UHF* algebras).

## Some facts about the propinquity

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).
- *Quantum tori* (with various metrics) form a continuous family of quantum compact metric spaces in their deformation parameter for the propinquity and are limits of finite dimensional “fuzzy tori” (L., 2013, 2021).
- There are various naturally continuous families of AF algebras (*Effros-Shen algebras*, *UHF* algebras).
- Spheres are limits for the propinquity of *fuzzy spheres* (Rieffel, 15).

## Some facts about the propinquity

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).
- *Quantum tori* (with various metrics) form a continuous family of quantum compact metric spaces in their deformation parameter for the propinquity and are limits of finite dimensional “fuzzy tori” (L., 2013, 2021).
- There are various naturally continuous families of AF algebras (*Effros-Shen algebras*, *UHF* algebras).
- Spheres are limits for the propinquity of *fuzzy spheres* (Rieffel, 15).
- There is a characterization of convergence for inductive limits in terms of the existence of certain morphisms (Farsi-L-Packer,

## Some facts about the propinquity

- The set of *classical* compact metric spaces is *nowhere dense* and *closed*.
- There is an analogue of *Gromov's compactness theorem* (L., 2015).
- There is a *covariant* version of the *propinquity* (L., 2017–2020), and group actions can “pass to the limit” (Bassi-Conti-Farsi-L., 2023).
- *Quantum tori* (with various metrics) form a continuous family of quantum compact metric spaces in their deformation parameter for the propinquity and are limits of finite dimensional “fuzzy tori” (L., 2013, 2021).
- There are various naturally continuous families of AF algebras (*Effros-Shen algebras*, *UHF* algebras).
- Spheres are limits for the propinquity of *fuzzy spheres* (Rieffel, 15).
- There is a characterization of convergence for inductive limits in terms of the existence of certain morphisms (Farsi-L-Packer,

- 1 *Compact Quantum Metric Spaces*
- 2 *The Gromov-Hausdorff Propinquity*
- 3 *The Spectral Propinquity*
- 4 *Examples and Applications*

## Metrical $C^*$ -correspondences

If  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  is a *metric spectral triple*, then

$$\text{qvb}(\mathfrak{A}, \mathcal{H}, \mathbb{D}) := (\mathcal{H}, \mathbb{D}, \mathfrak{A}, \mathbb{L}_{\mathbb{D}}, \mathbb{C}, 0)$$

where  $\mathbb{D} = \|\cdot\|_{\mathcal{H}} + \|\mathbb{D}\cdot\|_{\mathcal{H}}$ , is an example of the following structure.


*Definition (L. (16,18,19))*

A *metrical  $C^*$ -correspondence*  $\Omega = (\mathcal{M}, \mathbb{D}, \mathfrak{A}, \mathbb{L}_{\mathfrak{A}}, \mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$  is given by:

- ❶ two *quantum compact metric spaces*  $(\mathfrak{A}, \mathbb{L}_{\mathfrak{A}})$  and  $(\mathfrak{B}, \mathbb{L}_{\mathfrak{B}})$ ,
- ❷ an  $\mathfrak{A}$ - $\mathfrak{B}$   $C^*$ -correspondence  $\mathcal{M}$ , with  $\mathfrak{B}$ -valued inner product  $\langle \cdot, \cdot \rangle_{\mathcal{M}}$ ,
- ❸  $\mathbb{D}$  is a *norm* on a dense subspace of  $\mathcal{M}$  such that:
  - ❶  $\mathbb{D} \geq \|\cdot\|_{\mathcal{M}} := \sqrt{\langle \cdot, \cdot \rangle_{\mathcal{M}}}$
  - ❷  $\{\omega \in \mathcal{M} : \mathbb{D}(\omega) \leq 1\}$  is compact in  $(\mathcal{M}, \|\cdot\|_{\mathcal{M}})$ ,
  - ❸  $\forall \eta, \omega \in \mathcal{M} \quad \max\{\mathbb{L}_{\mathfrak{B}}(\Re \langle \omega, \eta \rangle_{\mathcal{M}}), \mathbb{L}_{\mathfrak{B}}(\Im \langle \omega, \eta \rangle_{\mathcal{M}})\} \leq H\mathbb{D}(\omega)\mathbb{D}(\eta),$
  - ❹  $\forall \eta \in \mathcal{M} \quad \forall a \in \text{sa}(\mathfrak{A}) \quad \mathbb{D}(a\eta) \leq G(\|a\|_{\mathfrak{A}} + \mathbb{L}_{\mathfrak{A}}(b))\mathbb{D}(\eta).$




# Tunnels between Metrical $C^*$ -correspondences




A tunnel:  $\mathbf{L}_j(a) = \inf \mathbf{L}_{\mathfrak{D}}(\theta_j^{-1}(\{a\}))$ .

# Tunnels between Metrical $C^*$ -correspondences




$\mathcal{J}$  is a  $\mathfrak{D}$ -module,  $\mathbf{D}_j(\omega) = \inf \mathbf{T}(\Pi_j^{-1}(\{\omega\}))$ ,  $T$   $\mathfrak{D}$ -norm

# Tunnels between Metrical $C^*$ -correspondences



$\mathcal{J}$  is a  $\mathfrak{D}$ - $\mathfrak{C}$ - $C^*$ -corr;  $(\mathfrak{C}, Q, \pi_1, \pi_2)$  tunnel.

# Extent of Metrical Tunnels



$$\chi(\tau) = \max \{ \chi((\mathfrak{D}, L_{\mathfrak{D}}, \theta_1, \theta_2)), \chi((\mathfrak{C}, Q, \pi_1, \pi_2)) \}.$$

# The metrical Propinquity

*Definition (L. 16,18)*

The *metrical propinquity*  $\Lambda^{\text{met}}(\mathbb{A}_1, \mathbb{A}_2)$  between two metrical  $C^*$ -correspondences  $\mathbb{A}_1$  and  $\mathbb{A}_2$ , is defined by

$$\inf \{ \chi(\tau) : \tau \text{ is a tunnel from } \mathbb{A}_1 \text{ to } \mathbb{A}_2 \}.$$

# The metrical Propinquity

## Definition (L. 16,18)

The *metrical propinquity*  $\Lambda^{\text{met}}(\mathbb{A}_1, \mathbb{A}_2)$  between two metrical  $C^*$ -correspondences  $\mathbb{A}_1$  and  $\mathbb{A}_2$ , is defined by

$$\inf \{ \chi(\tau) : \tau \text{ is a tunnel from } \mathbb{A}_1 \text{ to } \mathbb{A}_2 \}.$$

## Theorem (L. 16,18)

The *metrical propinquity* is a complete distance on metrical  $C^*$ -correspondences, up to full quantum isometry.

# The metrical Propinquity

## Definition (L. 16,18)

The *metrical propinquity*  $\Lambda^{\text{met}}(\mathbb{A}_1, \mathbb{A}_2)$  between two metrical  $C^*$ -correspondences  $\mathbb{A}_1$  and  $\mathbb{A}_2$ , is defined by

$$\inf \{ \chi(\tau) : \tau \text{ is a tunnel from } \mathbb{A}_1 \text{ to } \mathbb{A}_2 \}.$$

## Theorem (L. 16,18)

The *metrical propinquity* is a complete distance on metrical  $C^*$ -correspondences, up to full quantum isometry.

A nontrivial example of convergence of modules for our metric is given by *Heisenberg modules over quantum tori*, where the D-norm arises from *Connes' connection*.

# The metrical Propinquity

## Definition (L. 16,18)

The *metrical propinquity*  $\Lambda^{\text{met}}(\mathbb{A}_1, \mathbb{A}_2)$  between two metrical  $C^*$ -correspondences  $\mathbb{A}_1$  and  $\mathbb{A}_2$ , is defined by

$$\inf \{ \chi(\tau) : \tau \text{ is a tunnel from } \mathbb{A}_1 \text{ to } \mathbb{A}_2 \}.$$

## Theorem (L. 16,18)

The *metrical propinquity* is a complete distance on metrical  $C^*$ -correspondences, up to full quantum isometry.

When applied to spectral triples, the metrical propinquity provides metric information, but, maybe most importantly, it encodes the *convergence of the domains of the Dirac operators*.



# Phase

## *The phase of a Dirac operator*

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

# Phase

## *The phase of a Dirac operator*

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

If  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  is a spectral triple then:

$$t \in \mathbb{R} \mapsto U_t = \exp(it\mathbb{D})$$

is a strongly continuous action of  $\mathbb{R}$  on  $\mathcal{H}$  such that:

$$\forall t \in \mathbb{R}, \xi \in \mathcal{H} \quad \mathbb{D}(U_t \xi) = \|U_t \xi\|_{\mathcal{H}} + \|\mathbb{D}U_t \xi\|_{\mathcal{H}} = \mathbb{D}(\xi).$$

# Phase

## *The phase of a Dirac operator*

What do we need to add to our construction of our metric on spectral triples so that distance 0 means equivalence by conjugation?

If  $(\mathfrak{A}, \mathcal{H}, \mathbb{D})$  is a spectral triple then:

$$t \in \mathbb{R} \mapsto U_t = \exp(it\mathbb{D})$$

is a strongly continuous action of  $\mathbb{R}$  on  $\mathcal{H}$  such that:

$$\forall t \in \mathbb{R}, \xi \in \mathcal{H} \quad \mathbb{D}(U_t \xi) = \|U_t \xi\|_{\mathcal{H}} + \|\mathbb{D}U_t \xi\|_{\mathcal{H}} = \mathbb{D}(\xi).$$

## *Next step*

How do we incorporate convergence of group actions?

## *Separation for Family of Operators*

Let  $(\mathcal{M}, D, \mathfrak{A}, L_{\mathfrak{A}}, \mathfrak{B}, L_{\mathfrak{B}})$  be a metrical  $C^*$ -correspondence.

## Separation for Family of Operators

Let  $(\mathcal{M}, \mathbf{D}, \mathfrak{A}, \mathbf{L}_{\mathfrak{A}}, \mathfrak{B}, \mathbf{L}_{\mathfrak{B}})$  be a metrical  $C^*$ -correspondence.

If  $\varphi, \psi \in \mathcal{M}^*$ , then we set

$$\mathbf{mk}_{\mathbf{D}}(\varphi, \psi) := \sup \{ |\varphi(\omega) - \psi(\omega)| : \omega \in \text{dom}(\mathbf{D}), \mathbf{D}(\omega) \leq 1 \}.$$

If  $J$  is any index set, we then set

$$\mathbf{MK}_{\mathbf{D}}((\varphi_j)_{j \in J}, (\psi_j)_{j \in J}) = \sup_{j \in J} \mathbf{mk}_{\mathbf{D}}(\varphi_j, \psi_j).$$

## Separation for Family of Operators

Let  $(\mathcal{M}, \mathbf{D}, \mathfrak{A}, \mathbf{L}_{\mathfrak{A}}, \mathfrak{B}, \mathbf{L}_{\mathfrak{B}})$  be a metrical  $C^*$ -correspondence.

If  $\varphi, \psi \in \mathcal{M}^*$ , then we set

$$\mathbf{mk}_{\mathbf{D}}(\varphi, \psi) := \sup \{ |\varphi(\omega) - \psi(\omega)| : \omega \in \text{dom}(\mathbf{D}), \mathbf{D}(\omega) \leq 1 \}.$$

If  $J$  is any index set, we then set

$$\mathbf{MK}_{\mathbf{D}}((\varphi_j)_{j \in J}, (\psi_j)_{j \in J}) = \sup_{j \in J} \mathbf{mk}_{\mathbf{D}}(\varphi_j, \psi_j).$$

$\mathcal{S}(\mathbb{M})$  consists of the maps of the form  $\zeta \in \mathcal{M} \mapsto \varphi(\langle \zeta, \omega \rangle)$  for  $\omega \in \mathcal{M}$  with  $\mathbf{D}(\omega) \leq 1$ , and  $\varphi \in \mathcal{S}(\mathfrak{B})$ .

## Separation for Family of Operators

Let  $(\mathcal{M}, \mathbf{D}, \mathfrak{A}, \mathbf{L}_{\mathfrak{A}}, \mathfrak{B}, \mathbf{L}_{\mathfrak{B}})$  be a metrical  $C^*$ -correspondence.

If  $\varphi, \psi \in \mathcal{M}^*$ , then we set

$$\mathbf{mk}_{\mathbf{D}}(\varphi, \psi) := \sup \{ |\varphi(\omega) - \psi(\omega)| : \omega \in \text{dom}(\mathbf{D}), \mathbf{D}(\omega) \leq 1 \}.$$

If  $J$  is any index set, we then set

$$\mathbf{MK}_{\mathbf{D}}((\varphi_j)_{j \in J}, (\psi_j)_{j \in J}) = \sup_{j \in J} \mathbf{mk}_{\mathbf{D}}(\varphi_j, \psi_j).$$

$\mathcal{S}(\mathbb{M})$  consists of the maps of the form  $\zeta \in \mathcal{M} \mapsto \varphi(\langle \zeta, \omega \rangle)$  for  $\omega \in \mathcal{M}$  with  $\mathbf{D}(\omega) \leq 1$ , and  $\varphi \in \mathcal{S}(\mathfrak{B})$ .

Let now  $\tau : \mathbb{A} \xleftarrow{(\Pi_{\mathbb{A}}, \dots)} \mathbb{M} \xrightarrow{(\Pi_{\mathbb{B}}, \dots)} \mathbb{B}$  be a tunnel between metrical  $C^*$ -correspondences.

### Definition (L., 23)

Let  $J \neq \emptyset$  The *separation*  $\text{sep}(A, B|\tau)$  of a family  $A := (a_j)_{j \in J}$  in  $\mathfrak{B}(\mathbb{A})$ , and a family  $B := (b_j)_{j \in J}$  in  $\mathfrak{B}(\mathbb{B})$ , wrt  $\tau$ , is

$$\mathbf{Haus}_{\mathbf{MK}_{\mathbf{D}}}(\{\varphi \circ a_j \circ \Pi_{\mathfrak{A}} : \varphi \in \mathcal{S}(\mathbb{A})\}, \{\psi \circ b_j \circ \Pi_{\mathfrak{B}} : \psi \in \mathcal{S}(\mathbb{B})\})$$

# The Spectral Propinquity

## Definition (L., 18)

The *spectral propinquity*  $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2))$  between two metric spectral triples  $(\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1)$  and  $(\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)$  is

$$\inf \left\{ \frac{\sqrt{2}}{2}, \varepsilon > 0 : \exists \tau \text{ tunnel from } (\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1) \text{ to } (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2) \right. \\ \left. \text{such that } \max \left\{ \chi(\tau), \text{sep} \left( (e^{it\mathbb{D}_1})_{0 \leq t \leq \frac{1}{\varepsilon}}, (e^{it\mathbb{D}_2})_{0 \leq t \leq \frac{1}{\varepsilon}} | \tau \right) \right\} \leq \varepsilon \right\}.$$



# The Spectral Propinquity

## Definition (L., 18)

The *spectral propinquity*  $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2))$  between two metric spectral triples  $(\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1)$  and  $(\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)$  is

$$\inf \left\{ \frac{\sqrt{2}}{2}, \varepsilon > 0 : \exists \tau \text{ tunnel from } (\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1) \text{ to } (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2) \right. \\ \left. \text{such that } \max \left\{ \chi(\tau), \text{sep} \left( (e^{it\mathbb{D}_1})_{0 \leq t \leq \frac{1}{\varepsilon}}, (e^{it\mathbb{D}_2})_{0 \leq t \leq \frac{1}{\varepsilon}} | \tau \right) \right\} \leq \varepsilon \right\}.$$

## Theorem (L., 18)

The *spectral propinquity*  $\Lambda^{\text{spec}}$  is a *metric* on the class of spectral triples, up to unitary equivalence, i.e.  $\Lambda^{\text{spec}}((\mathfrak{A}_1, \mathcal{H}_1, \mathbb{D}_1), (\mathfrak{A}_2, \mathcal{H}_2, \mathbb{D}_2)) = 0$  if, and only if there exists a *unitary*  $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  such that  $U \text{dom}(\mathbb{D}_1) = \text{dom}(\mathbb{D}_2)$ ,

$$U \mathbb{D}_1 U^* = \mathbb{D}_2 \text{ and } \text{Ad}_U^* \text{-isomorphism from } \mathfrak{A}_1 \text{ to } \mathfrak{A}_2.$$

- 1 *Compact Quantum Metric Spaces*
- 2 *The Gromov-Hausdorff Propinquity*
- 3 *The Spectral Propinquity*
- 4 *Examples and Applications*

## Matrix Models: Clock and Shift, Fuzzy Tori

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ & r_n & \ddots & \\ & \vdots & \ddots & \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with  $r_n = \exp\left(\frac{2i\pi}{n}\right)$ . Note that  $C^*(C_n, S_n) = \mathfrak{B}(\mathbb{C}^n)$ .

## Matrix Models: Clock and Shift, Fuzzy Tori

$$C_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ & r_n & \ddots & \\ & & \ddots & 0 \\ 0 & \cdots & 0 & r_n^{n-1} \end{pmatrix} \text{ and } S_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \ddots & \\ & & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1 & \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

with  $r_n = \exp\left(\frac{2i\pi}{n}\right)$ . Note that  $C^*(C_n, S_n) = \mathfrak{B}(\mathbb{C}^n)$ . Let  $\mathbb{D}_n$  be defined on  $\mathcal{H}_n = \mathfrak{B}(\mathbb{C}^n) \otimes \mathbb{C}^4$  be defined by:

$$\begin{aligned} \mathbb{D}_n = \frac{n}{2\pi} & \left( \left[ \frac{S_n + S_n^*}{2}, \cdot \right] \otimes \gamma_1 + \left[ \frac{S_n - S_n^*}{2i}, \cdot \right] \otimes \gamma_2 \right. \\ & \left. + \left[ \frac{C_n^* + C_n}{2}, \cdot \right] \otimes \gamma_3 + \left[ \frac{C_n^* - C_n}{2i}, \cdot \right] \otimes \gamma_4 \right), \end{aligned}$$

with  $\gamma_j \gamma_k + \gamma_k \gamma_j = \delta_j^k$  ( $j, k \in \{1, 2, 3, 4\}$ ). Then  $(C^*(C_n, S_n), \mathcal{H}_n, \mathbb{D}_n)$  is a spectral triple.

## Convergence of Fuzzy tori

### Theorem (L., 21)

The sequence  $(C^*(C_n, S_n), \mathfrak{B}(\mathbb{C}^n) \otimes \mathbb{C}^4, \mathcal{D}_n)_{n \in \mathbb{N}}$ , where

$$\mathcal{D}_n = \frac{n}{2\pi} \left( \left[ \frac{S_n + S_n^*}{2}, \cdot \right] \otimes \gamma_1 + \left[ \frac{S_n - S_n^*}{2i}, \cdot \right] \otimes \gamma_2 \right. \\ \left. + \left[ \frac{C_n^* + C_n}{2}, \cdot \right] \otimes \gamma_3 + \left[ \frac{C_n^* - C_n}{2i}, \cdot \right] \otimes \gamma_4 \right),$$

converges, for the spectral propinquity, to the spectral triple  $(C(T^2), L^2(\mathbb{T}^2) \otimes \mathbb{C}^4, \mathcal{D}_{\mathbb{T}^2})$ , where  $C(\mathbb{T}^2) = \{(e^{i\theta}, e^{i\psi}) : \theta, \psi \in [0, 2\pi]\}$ , and on a dense subspace of  $L^2(\mathbb{T}^2) \otimes \mathbb{C}^4$ , we set

$$\mathcal{D}_{\mathbb{T}^2} = \cos(\psi) \partial_\theta \otimes \gamma_1 + \sin(\psi) \partial_\theta \otimes \gamma_2 + \cos(\theta) \partial_\psi \otimes \gamma_3 + \sin(\theta) \partial_\psi \otimes \gamma_4.$$

## Convergence of Fuzzy tori

### Theorem (L., 21)

The sequence  $(C^*(C_n, S_n), \mathfrak{B}(\mathbb{C}^n) \otimes \mathbb{C}^4, \mathcal{D}_n)_{n \in \mathbb{N}}$ , where


$$\mathcal{D}_n = \frac{n}{2\pi} \left( \left[ \frac{S_n + S_n^*}{2}, \cdot \right] \otimes \gamma_1 + \left[ \frac{S_n - S_n^*}{2i}, \cdot \right] \otimes \gamma_2 \right. \\ \left. + \left[ \frac{C_n^* + C_n}{2}, \cdot \right] \otimes \gamma_3 + \left[ \frac{C_n^* - C_n}{2i}, \cdot \right] \otimes \gamma_4 \right),$$

converges, for the spectral propinquity, to the spectral triple  $(C(T^2), L^2(\mathbb{T}^2) \otimes \mathbb{C}^4, \mathcal{D}_{\mathbb{T}^2})$ , where  $C(\mathbb{T}^2) = \{(e^{i\theta}, e^{i\psi}) : \theta, \psi \in [0, 2\pi]\}$ , and on a dense subspace of  $L^2(\mathbb{T}^2) \otimes \mathbb{C}^4$ , we set

$$\mathcal{D}_{\mathbb{T}^2} = \cos(\psi) \partial_\theta \otimes \gamma_1 + \sin(\psi) \partial_\theta \otimes \gamma_2 + \cos(\theta) \partial_\psi \otimes \gamma_3 + \sin(\theta) \partial_\psi \otimes \gamma_4.$$

This result is extended to more general fuzzy/quantum tori (L., 21).

# Piecewise $C^1$ Fractals Curves




The Sierpiński gasket

Let  $\mathcal{J} = \oplus_{n=1}^{\infty} \oplus_{j=1}^{3^n} L^2([-1, 1])$  and  $\partial(\xi) = \left( 2^n \xi'_{n,j} \right)_{n \in \mathbb{N}, j \in \{1, \dots, 3^n\}}$ .

For  $n \in \mathbb{N} \setminus \{0\}$ , let  $C_{n,1}, \dots, C_{n,3^n}$  be affine functions from  $[0, 1]$ , which parametrize every edge of the level  $n$  triangles in  $\mathcal{S}\mathcal{G}_n$ .

# Piecewise $C^1$ Fractals Curves



The Sierpiński gasket

Let  $\mathcal{J} = \oplus_{n=1}^{\infty} \oplus_{j=1}^{3^n} L^2([-1, 1])$  and  $\mathfrak{d}(\xi) = \left( 2^n \xi'_{n,j} \right)_{n \in \mathbb{N}, j \in \{1, \dots, 3^n\}}$ .

For  $n \in \mathbb{N} \setminus \{0\}$ , let  $C_{n,1}, \dots, C_{n,3^n}$  be affine functions from  $[0, 1]$ , which parametrize every edge of the level  $n$  triangles in  $\mathcal{S}\mathcal{G}_n$ . For

$f \in C(\mathcal{S}\mathcal{G}_{\infty})$ , we set  $f \cdot \xi_{n,j} := f \circ C_{n,j}(| \cdot |) \xi_{n,j}$ .

$(C(\mathcal{S}\mathcal{G}_{\infty}), \mathcal{J}, \mathfrak{d})$  is a spectral triple, constructed by *Lapidus et al.*



## Convergence of certain fractals



For each  $n \in \mathbb{N}$ , let  $\mathfrak{d}_n$  be the restriction of  $\mathfrak{d}$  to  $\mathcal{I}_N := \oplus_{n=1}^N \oplus_{j=1}^{3^N} L^2([-1, 1])$ , and let  $C(\mathcal{S}\mathcal{G}_N)$  acts on  $\mathcal{I}_N$ .

*Theorem (Landry, Lapidus, L., 20)*

Let  $(C(\mathcal{S}\mathcal{G}_\infty), \mathcal{I}, \mathfrak{d})$  be the spectral triple over the Sierpiński gasket  $\mathcal{S}\mathcal{G}_\infty$ , and for each  $n \in \mathbb{N}$ , let  $(\mathcal{S}\mathcal{G}_n, \mathcal{I}_n, \mathfrak{d}_n)$  be its “restriction” to the finite graph  $\mathcal{S}\mathcal{G}_n$ . Then

$$\lim_{n \rightarrow \infty} \Lambda^{\text{spec}}((C(\mathcal{S}\mathcal{G}_\infty), \mathcal{I}, \mathfrak{d}), (C(\mathcal{S}\mathcal{G}_n), \mathcal{I}_n, \mathfrak{d}_n)) = 0.$$

# Collapse!

## *Theorem (L., 23)*

If  $(\mathfrak{A}, \mathcal{H}, D)$  is a spectral triple with  $0 \in \text{Sp}(D)$ , then

$$\lim_{\lambda \rightarrow \infty} \Lambda^{\text{spec}}((\mathfrak{A}, \mathcal{H}, \lambda D), (\mathbb{C}, \ker D, 0)) = 0.$$

# Inductive Limits of Group $C^*$ -algebras

## Theorem (Farsi-L-Packer, 23)

Let  $G = \bigcup_{n \in \mathbb{N}} G_n$  be an Abelian discrete group, with  $(G_n)_{n \in \mathbb{N}}$  a strictly increasing sequence of subgroups of  $G$ . Let  $\sigma$  be a 2-cocycle of  $G$ . Let  $\mathbb{L}_H$  be a length function over  $G$  such that  $\lim_{n \rightarrow \infty} \text{Haus}_{\mathbb{L}_H}(G_n, G) = 0$  and  $\mathbb{L}_H$  is proper on each  $G_n$ . If there exists a strictly increasing function  $\text{scale} : \mathbb{N} \rightarrow \mathbb{R}$  such that, setting  $\mathbb{F} : g \in G \mapsto \text{scale}(\min\{n : g \in G_n\})$ , the length  $\max\{\mathbb{L}_H, \mathbb{F}\}$  has the *doubling property*, then

$$\lim_{n \rightarrow \infty} \Lambda^{\text{spec}}((C^*(G, \sigma), \ell^2(G) \otimes \mathbb{C}^2, \mathbb{D}), \\ (C^*(G_n, \sigma), \ell^2(G_n) \otimes \mathbb{C}^2, \mathbb{D}|_{\ell^2(G_n) \otimes \mathbb{C}^2})) = 0$$

where

$$\mathbb{D} := \text{mul}_{\mathbb{L}_H} \otimes \gamma_1 + \text{mul}_{\mathbb{F}} \otimes \gamma_2.$$

Applications: noncommutative solenoids, Bunce-Deddens algebras

# Convergence of the Spectrum

## Theorem (L., 23)

If  $(\mathfrak{A}_n, \mathcal{H}_n, \mathcal{D}_n)_{n \in \mathbb{N}}$  is a sequence of metric spectral triples converging to  $(\mathfrak{A}_\infty, \mathcal{H}_\infty, \mathcal{D}_\infty)$  for the spectral propinquity, then:

$$\mathrm{Sp}(\mathcal{D}_\infty) = \left\{ \lambda \in \mathbb{R} : \forall_{\mathbb{N}} n \quad \exists \lambda_n \in \mathrm{Sp}(\mathcal{D}_n) \quad \lambda = \lim_{n \rightarrow \infty} \lambda_n \right\}.$$

Moreover, under reasonable assumptions, the *multiplicity of the eigenvalues converge as well*.

# Convergence of multiplicities

## Theorem (L., 23)

If  $(\mathfrak{A}_n, \mathcal{H}_n, D_n)_{n \in \mathbb{N}}$  is a sequence of metric spectral triples *converging, for the spectral propinquity*, to a metric spectral triple  $(\mathfrak{A}_\infty, \mathcal{H}_\infty, D_\infty)$ , and if  $\lambda \in \text{Sp}(D_\infty)$ , such that:

- 1 there exists  $\delta > 0$  and  $N \in \mathbb{N}$  such that, for all  $n \in \mathbb{N}$ ,  $n > N$ , the intersection  $\text{Sp}(D_n) \cap (\lambda - \delta, \lambda + \delta)$  is a *singleton*  $\{\lambda_n\}$ ,
- 2 if  $(\text{multiplicity}(\lambda_n | D_n))_{n \in \mathbb{N}}$  is *bounded*,

then

$$\lim_{n \rightarrow \infty} \text{multiplicity}(\lambda_n | D_n) = \text{multiplicity}(\lambda | D_\infty).$$

# Convergence of multiplicities

## Theorem (L., 23)

If  $(\mathcal{A}_n, \mathcal{H}_n, \mathcal{D}_n)_{n \in \mathbb{N}}$  is a sequence of metric spectral triples *converging, for the spectral propinquity*, to a metric spectral triple  $(\mathcal{A}_\infty, \mathcal{H}_\infty, \mathcal{D}_\infty)$ , and if  $\lambda \in \text{Sp}(\mathcal{D}_\infty)$ , such that:

- 1 there exists  $\delta > 0$  and  $N \in \mathbb{N}$  such that, for all  $n \in \mathbb{N}$ ,  $n > N$ , the intersection  $\text{Sp}(\mathcal{D}_n) \cap (\lambda - \delta, \lambda + \delta)$  is a *singleton*  $\{\lambda_n\}$ ,
- 2 if  $(\text{multiplicity}(\lambda_n | \mathcal{D}_n))_{n \in \mathbb{N}}$  is *bounded*,

then

$$\lim_{n \rightarrow \infty} \text{multiplicity}(\lambda_n | \mathcal{D}_n) = \text{multiplicity}(\lambda | \mathcal{D}_\infty).$$

From this, we can conclude, under similar assumptions, that  $\text{trace}(f(\mathcal{D}_n))$  converges to  $\text{trace}(f(\mathcal{D}))$  for any  $f \in C_0(\mathbb{R})$ ; thus we obtain convergence for spectral actions.

# Thank you!

- *The Quantum Gromov-Hausdorff Propinquity*, F. Latrémolière, *Transactions of the AMS* **368** (2016) 1, pp. 365–411.
- *The Dual Gromov-Hausdorff Propinquity*, F. Latrémolière, *Journal de Mathématiques Pures et Appliquées* **103** (2015) 2, pp. 303–351, ArXiv: 1311.0104
- *The dual modular propinquity and completeness*, F. Latrémolière, *J. Noncomm. Geometry* **15** (2021) no. 1, 347–398.
- *Metric approximations of spectral triples on the Sierpiński gasket and other fractal curves*, T. Landry, M. Lapidus, F. Latrémolière, *Adv. Math.* **385** (2021), paper no. 107771, 43 pp.
- *Convergence of Spectral Triples on Fuzzy Tori to Spectral Triples on Quantum Tori*, F. Latrémolière, *Comm. Math. Phys.* **388** (2021) no. 2, 1049–1128.
- *The Gromov-Hausdorff propinquity for metric spectral triples*, F. Latrémolière, *Adv. Math.* **404** (2022), paper no. 108393, 56 pp.
- *Continuity of the spectrum of Dirac operators of spectral triples for the spectral propinquity*, F. Latrémolière, *Math. Ann.* (2023), <https://doi.org/10.1007/s00208-023-02659-x>