

# Math265 Real Analysis Class Notes

Based on lectures by Prof. Huang

Notes taken by Kaifeng Lu

Errata: [klu15 at u dot rochester dot edu](mailto:klu15@u.rochester.edu)

Fall 2024



# **1 Preliminaries**

Still working on it...(missed 1st month due to late enrollment)

# **2 The Real Numbers**

Still working on it...

### 3 Sequences and Series

#### 3.1 Sequences and their limits

**Definition (Sequence).** A sequence of real numbers is a *function* from  $\mathbb{N}$  to  $\mathbb{R}$ .

We adopt the notation with a *sequence*:

$$a : \mathbb{N} \rightarrow \mathbb{R}$$

where instead of writing  $a(1), a(2), \dots$ , we write it as  $a_1, a_2, \dots$  which we called them terms or elements of the sequence.

**Notation.**

$$(a_n)_{n=1}^{\infty} \text{ or } (a_n)_{n \in \mathbb{N}} \text{ or } (a_n) \text{ or } (a_n | n \in \mathbb{N})$$

**Definition (Converge to x).** A sequence  $(x_n) \in \mathbb{R}$  converges to  $x \in \mathbb{R}$  if

$$\forall \varepsilon > 0, \exists N_\varepsilon \in \mathbb{N} \text{ such that } n \geq N_\varepsilon \rightarrow |x_n - x| < \varepsilon$$

We write

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} (x_n) = x.$$

**Definition (Convergent & Divergent).** A sequence is convergent if it has a limit in  $\mathbb{R}$ , and is divergent if it has no limit in  $\mathbb{R}$ .

**Theorem (Uniqueness of Limit).** A sequence in  $\mathbb{R}$  can have at most one limit. Or, the limit of a sequence is unique if the limit exists

*Proof.* Let  $(x_n)$  be a sequence of real numbers. Suppose  $x, x'$  are limits of  $(x_n)$ . We want to prove  $x = x'$  by contradiction.

Assume  $|x - x'| > 0$ . If we consider  $\varepsilon := \frac{1}{3}|x - x'| > 0$ , then

The existence of  $\lim_{x_n \rightarrow x}$  implies that  $\exists N_1 \in \mathbb{N}$  such that  $|x_n - x| < \varepsilon$  if  $n \geq N_1$ .

Similarly, existence of  $\lim_{x_n \rightarrow x'}$  implies that  $\exists N_2 \in \mathbb{N}$  such that  $|x_n - x'| < \varepsilon$  if  $n \geq N_2$ .

Thus,

$$\begin{aligned} |x - x'| &\leq |x - x_{N_1+N_2} + x_{N_1+N_2} - x'| \\ &\leq |x - x_{N_1+N_2}| + |x_{N_1+N_2} - x'| \text{ by triangle inequality} \\ &< \varepsilon + \varepsilon \\ &= \frac{2}{3}|x - x'| \end{aligned}$$

Then,

$$\frac{1}{3}|x - x'| < 0, \text{ which is a contradiction}$$

we thereby prove by contradiction that

$$|x - x'| = 0, \text{ which is equivalent to } x = x'$$

■

**Example.**

$$\lim_{n \rightarrow \infty} \left(\frac{1}{n}\right) = 0$$

Goal:  $\forall \varepsilon > 0$ , want to find  $N_\varepsilon$  such that  $\left|\frac{1}{n} - 0\right| < \varepsilon$  for  $n > N$ , so it suffices to show that

$$\frac{1}{n} < \varepsilon \Leftrightarrow \frac{1}{\varepsilon} < n$$

*Proof.* Let  $\varepsilon > 0$ . Apply Archimedean's property to  $\frac{1}{\varepsilon}$ , then

$$\begin{aligned} &\exists N \in \mathbb{N} \text{ such that } \frac{1}{\varepsilon} < N \\ \Rightarrow &\forall n \geq N, \left|\frac{1}{n} - 0\right| = \frac{1}{n} \leq \frac{1}{N} < \varepsilon. \\ \Rightarrow &\lim_{n \rightarrow \infty} \frac{1}{n} = 0 \end{aligned}$$

■

**Theorem.** Let  $(x_n)$  be a sequence of real numbers, and let  $x \in \mathbb{R}$ . The following are equivalent:

1.  $x_n \rightarrow x$
2.  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$  such that  $|x_n - x| < \varepsilon$ , for  $n \geq N$
3.  $\dots x - \varepsilon < x_n < x + \varepsilon \dots$
4.  $\forall \varepsilon$ -neighborhood  $V_\varepsilon(x), \exists N \in \mathbb{N}$  such that  $x_n \in V_\varepsilon(x)$  for  $n \geq N$

**Sketch of proof:**

$$(1) \Leftrightarrow (2) \Leftrightarrow (3) \Leftrightarrow (4)$$

**Proposition.**

$$\lim_{n \rightarrow \infty} (2\sqrt{2n+1} - \sqrt{2n}) = 0$$

*Proof.* Let  $\varepsilon > 0$ . Consider

$$N = \left\lceil \frac{1}{2} \left( \frac{1}{2\varepsilon} \right)^2 \right\rceil \in \mathbb{N}$$

$$n > N \Rightarrow n > \frac{1}{2} \left( \frac{1}{2\varepsilon} \right)^2 \Rightarrow \frac{1}{2\sqrt{2n}} < \varepsilon \Rightarrow \left| \sqrt{2n+1} - \sqrt{2n} \right| = \dots = \frac{1}{\sqrt{2n+1} + \sqrt{2n}} < \varepsilon$$

■

**Remark.**

$$\lim_{n \rightarrow \infty} (-1)^n \text{ does not exist.}$$

**Definition (m-tail).** If  $(x_n)$  is a sequence of real numbers and  $m \in \mathbb{N}$ , then the **m-tail** of  $(x_n)$  is the sequence

$$\{x_{n+m} : n \in \mathbb{N}\} = \{x_{m+1}, x_{m+2}, \dots\}$$

**Theorem.** Let  $(x_n)$  be a sequence and  $m \in \mathbb{N}$ . Then  $(x_n)$  is **convergent** iff  $(x_{n+m})$  is **convergent**. Moreover,

$$\lim_{n \rightarrow \mathbb{N}} (x_n) = \lim_{n \rightarrow \mathbb{N}} (x_{n+m})$$

*Proof.* ( $\Rightarrow$ )

Suppose  $x_n \rightarrow x$ . Let

$$\varepsilon > 0, \exists N_\varepsilon > 0, \text{ such that } |x_n - x| < \varepsilon \text{ for } n \geq N_\varepsilon$$

Consider  $N'_\varepsilon := N_\varepsilon + m$  then

$$n + m \geq N'_\varepsilon \Rightarrow n \geq N_\varepsilon \Rightarrow |x_{n+m} - x| < \varepsilon$$

It follows that

$$n \geq N_\varepsilon \Rightarrow n + m \geq N'_\varepsilon \Rightarrow |x_{n+m} - x| < \varepsilon$$

( $\Leftarrow$ )

Suppose  $x_{n+m} \rightarrow x$ .

$$\forall \varepsilon > 0, \exists N_\varepsilon > 0 \text{ such that } |x_{n+m} - x| < \varepsilon, \forall n \geq N_\varepsilon$$

Consider  $N := N_\varepsilon + m$ . Then

$$\begin{aligned} n &\geq N = N_\varepsilon + m \\ \Rightarrow n - m &\geq N_\varepsilon \\ \Rightarrow |x_{(n-m)+m} - x| &< \varepsilon \\ \Rightarrow |x_n - x| &< \varepsilon \end{aligned}$$

■

**Remark.** We say that a sequence  $(x_n)$  ***ultimately*** has a property if that property holds for some tail of  $(x_n)$

**Theorem.** Let  $x_n$  be a sequence of real numbers. Let  $a_n$  be a sequence of positive real numbers such that  $\lim_{n \rightarrow \infty} a_n = 0$ . If  $\exists c > 0, m \in \mathbb{N}, x \in \mathbb{R}$  such that

$$|x_n - x| \leq c \cdot a_n, \forall n \geq m$$

then

$$x_n \rightarrow x$$

*Proof.* We know that

$$\forall \varepsilon > 0, \exists N \geq 0 \text{ s.t. } |a_n| < \frac{\varepsilon}{c}, \forall n \geq N$$

Consider  $N' = \max\{N, m\}, \forall n \geq N'$ . Then

$$\begin{aligned} |x_n - x| &\leq C a_n = c |a_n| < c \cdot \frac{\varepsilon}{c} = \varepsilon \\ \Rightarrow x_n &\rightarrow x \end{aligned}$$

■

**Proposition.**

$$\lim_{n \rightarrow \infty} \frac{17}{2 + 3n} = 0$$

*Proof.*

$$\left| \frac{17}{2 + 3n} - 0 \right| = \frac{17}{2 + 3n} \leq \frac{13}{3n} = \frac{17}{3} \cdot \frac{1}{n}$$

Apply the theorem above with

$$a_n = \frac{1}{n}, c = \frac{17}{3}, m = 1$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{17}{2 + 3n} = 0, \text{ since } \lim_{n \rightarrow \infty} \frac{1}{n} = 0$$

■

**Proposition.**

$$\forall c > 0, \lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$$

*Proof.* **Case 1:  $c = 1$**

$$\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$$

**Case 2:  $c > 1$**

Let  $d_n = c^{\frac{1}{n}} - 1$ . Then  $\forall n, d_n > 0$ . It follows that

$$\begin{aligned} (d_n + 1) &= c^{\frac{1}{n}} \Rightarrow c = (1 + d_n)^n \geq 1 + n \cdot d_n \text{ by Bernoulli's inequality} \\ &\Rightarrow d_n \leq (c - 1) \cdot \frac{1}{n} \\ &\Rightarrow \left| c^{\frac{1}{n}} - 1 \right| = d_n \leq (c - 1) \cdot \frac{1}{n} \end{aligned}$$

Apply the theorem with

$$C = c - 1, a_n = \frac{1}{n}, m = 1, x = 1$$

$$\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$$

**Case 3:  $c < 1$  (Note that we cannot use Bernoulli inequality here)**

Define  $e_n$  to be a sequence that satisfies

$$c^{\frac{1}{n}} = \frac{1}{1 + e_n}$$

Then  $e_n > 0 \forall n$ .

$$\begin{aligned} c &= \frac{1}{(1 + e_n)^n} \leq \frac{1}{1 + n \cdot e_n} < \frac{1}{n \cdot e_n} \\ &\Rightarrow e_n < \frac{1}{c} \cdot \frac{1}{n} \\ 1 - c^{\frac{1}{n}} &= 1 - \frac{1}{1 + e_n} = \frac{e_n}{1 + e_n} < e_n < \frac{1}{c} \cdot \frac{1}{n} \end{aligned}$$

Apply the theorem with

$$a_n = \frac{1}{n}, m = 1, C = \frac{1}{c}, x = 1$$

$$\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$$


■

### 3.2 Limit Theorems

**Definition** (Bounded sequence). A sequence  $(x_n) \in \mathbb{R}$  is **bounded** if

$$\exists M > 0 \text{ s.t. } \forall n \in \mathbb{N}, |x_n| \leq M$$

**Theorem.** A convergent sequence  $(x_n) \in \mathbb{R}$  is bounded.



*Proof.* By definition of convergent sequence, let  $\varepsilon = 1$ :

$$\exists N > 0 \text{ s.t. } \forall n \geq N, |x_n - x| < 1$$

Thus we have

$$\begin{aligned} & -1 < x_n - x < 1 \\ \Rightarrow & -1 + x < x_n < x + 1, \forall n \geq N \end{aligned}$$

Then define

$$M := \max \{|x_1|, |x_2|, |x_3|, \dots, |-1 + x|, |x + 1|\}$$

$$|x_n| \leq M$$

■

**Remark.** By contrapositive, an unbounded sequence is divergent.

**Definition.** Given sequence  $(x_n), (y_n) \in \mathbb{R}$ , we define following operations of sequence:

- **Sum**  $(x_n + y_n)$
- **Difference**  $(x_n - y_n)$
- **Product**  $(x_n \cdot y_n)$
- **Quotient**  $(\frac{x_n}{y_n})$  if  $\forall n \in \mathbb{N}, y_n \neq 0$
- **Multiple**  $(c \cdot x_n)$

**Theorem (Limit Laws).** Let  $(x_n), (y_n) \in \mathbb{R}$  be sequences of real numbers with  $x_n \rightarrow x, y_n \rightarrow y$ , and let  $c \in \mathbb{R}$ . Then

- $x_n + y_n \rightarrow x + y$
- $x_n - y_n \rightarrow x - y$
- $x_n \cdot y_n \rightarrow x \cdot y$
- $c \cdot x_n \rightarrow c \cdot x$
- If  $\forall n \in \mathbb{N}, y_n \neq 0$  and  $y \neq 0$ , then  $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$



*Proof of Sum. :*

$\forall \varepsilon > 0,$

$$\exists N_1 \in \mathbb{N} \text{ s.t. } \forall n \geq N_1, |x_n - x| < \frac{\varepsilon}{2}$$

$$\exists N_2 \in \mathbb{N} \text{ s.t. } \forall n \geq N_2, |y_n - y| < \frac{\varepsilon}{2}$$

Consider

$$N := \max \{N_1, N_2\}$$

Then

$$\begin{aligned} \forall n \geq N, |(x_n + y_n) - (x + y)| &= |x_n - x + y_n - y| \\ &\leq |x_n - x| + |y_n - y| \\ &= \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

■

*Proof of Difference. :*

Similarly,

$$\begin{aligned} \forall n \geq N, |(x_n - y_n) - (x - y)| &\leq |x_n - x| + |y_n - y| \end{aligned}$$

■

*Proof of Product. :* Since  $(x_n)$  is convergent, it is also bounded. Thus,

$$\exists M \in \mathbb{N} \text{ s.t. } \forall n \in \mathbb{N}, |x_n| \leq M$$

By definition of convergence,  $\forall \varepsilon > 0$ :

$$\exists N_1 > 0 \text{ s.t. } \forall n \geq N, |x_n - x| < \frac{\varepsilon}{2|y|}$$

$$\exists N_2 > 0 \text{ s.t. } \forall n \geq N, |y_n - y| < \frac{\varepsilon}{2M}$$

Then,  $\exists N = \max \{N_1, N_2\}$  s.t.  $\forall n \geq N,$

$$\begin{aligned} |x_n y_n - xy| &= |x_n y_n - x_n y + x_n y - xy| \\ &= |x_n(y_n - y) - y(x_n - x)| \\ &\leq |x_n| |y_n - y| + |y| |x_n - x| \\ &\leq M \cdot |y_n - y| + |y| |x_n - x| \\ &< M \cdot \frac{\varepsilon}{2M} + |y| \cdot \frac{\varepsilon}{2|y|} = \varepsilon \end{aligned}$$

■

*Proof of Multiply. :*

*Exercise*

■

*Proof of Quotient.*

$$\begin{aligned} \left| \frac{x_n}{y_n} - \frac{x}{y} \right| &= \left| \frac{x_n y - y_n x}{y_n y} \right| \\ &= \left| \frac{x_n y - x_n y_n + x_n y_n - y_n x}{y_n y} \right| \\ &\leq \left| \frac{1}{y_n y} (|x_n| |y_n - y| + |y_n| |x_n - x|) \right| \end{aligned}$$

Since  $y_n \rightarrow y$

$$\exists N_1 \in \mathbb{N} \text{ s.t. } \forall n \geq N, ||y_n| - |y|| \leq |y_n - y| < \frac{|y|}{2}$$

Then

$$\begin{aligned} \Rightarrow & -\frac{|y|}{2} < |y| - |y_n| < \frac{|y|}{2} \\ \Rightarrow & \frac{|y|}{2} < |y_n| \\ \Rightarrow & \frac{1}{|y_n|} < \frac{2}{|y|} \\ \Rightarrow & \frac{1}{|y_n y|} < \frac{2}{|y^2|} \end{aligned}$$

Define  $F := \frac{2}{|y^2|}$ .

Since  $(x_n)$  and  $(y_n)$  are bounded,

$\exists G \in \mathbb{N}$  s.t.  $\forall n \in \mathbb{N}$

$$|x_n| \leq G, |y_n| \leq G$$

Thus  $\forall n > 0$

$$\exists N_2 > 0 \text{ s.t. } \forall n \geq N_2, |x_n - x| < \frac{\varepsilon}{2GF}$$

$$\exists N_3 > 0 \text{ s.t. } \forall n \geq N_3, |y_n - y| < \frac{\varepsilon}{2GF}$$


Define  $N := \max \{N_1, N_2, N_3\}$ , then  $\exists N \in \mathbb{N}$  s.t.  $\forall n > N$ ,

$$\left| \frac{x_n}{y_n} - \frac{x}{y} \right| < F(G \cdot \frac{\varepsilon}{2GF} + G \cdot \frac{\varepsilon}{2GF}) = \varepsilon$$

■

**Theorem.** If  $(x_n)$  is a convergent sequence in  $\mathbb{R}$  of non-negative terms with  $(x_n) \rightarrow x$ , then  $x \geq 0$ .

*Sketch.* : Assume  $x < 0$ , choose  $\varepsilon \leq |x|$ , then  $\forall n > N, x_n < 0$ , which is a contradiction.



■

**Theorem.** If  $x_n \rightarrow x, y_n \rightarrow y$  are sequences in  $\mathbb{R}$  such that  $\forall n \in \mathbb{N}, x_n \leq y_n$ , then  $x \leq y$ .

*Proof.* Consider  $z_n := y_n - x_n$ . Then  $z_n \geq 0$  and  $z_n \rightarrow y - x$  by the limit law.

$$\Rightarrow y - x \geq 0$$

$$y \geq x$$

■

**Theorem.** If  $x_n \rightarrow x$  is a sequence in  $\mathbb{R}$ , and let  $a, b \in \mathbb{R}$  such that  $\forall n \in \mathbb{N}, a \leq x_n \leq b$ , then  $a \leq x \leq b$ .

*Proof.* Consider constant sequence  $a_n = a$  and  $b_n = b$ . Then this is true by the last theorem. ■

**Theorem (Squeeze Theorem).** Suppose  $(x_n), (y_n), (z_n)$  are sequences of real numbers such that  $\forall n \in \mathbb{N}, x_n \leq y_n \leq z_n$ . If  $\lim(x_n) = \lim(z_n)$ , then  $(y_n)$  is convergent and

$$\lim(x_n) = \lim(y_n) = \lim(z_n)$$

*Proof.* Let  $\forall \varepsilon > 0$ .

Write  $L = \lim(x_n)$ . Then by definition of limit,

$\exists N_1 \in \mathbb{N}$  s.t.  $\forall n \geq N$ ,

$$|x_n - L| < \varepsilon, |z_n - L| < \varepsilon$$

$$\Rightarrow -\varepsilon < x_n - L \leq y_n - L \leq z_n - L < \varepsilon$$

$$\Rightarrow |y_n - L| < \varepsilon$$

Thus  $(y_n)$  converges to  $L$  by definition of limit. ■

**Proposition.**  $\ln(n)$  is divergent.

*Proof.* Since  $\ln(n)$  is unbounded, it is divergent. ■

**Exercise.**  $\lim_{n \rightarrow \infty} \frac{3n^2 + 2n + 1}{5n^2 - 4} = \frac{3}{5}$

$$\lim_{n \rightarrow \infty} \frac{3n^2 + 2n + 1}{5n^2 - 4} = \lim_{n \rightarrow \infty} \frac{3 + 2\frac{1}{n} + \frac{1}{n^2}}{5 - 4\frac{1}{n^2}}$$

Notice that each term of  $\frac{1}{n}$  and  $\frac{1}{n^2}$  converges to 0. Thus by limit law,

$$\lim_{n \rightarrow \infty} \frac{3 + 2\frac{1}{n} + \frac{1}{n^2}}{5 - 4\frac{1}{n^2}} = \frac{\lim\{3 + 2 \cdot 0 + 0\}}{\lim\{5 - 4 \cdot 0\}} = \frac{3}{5}$$

**Proposition.**  $(-1)^n$  is divergent.

*Proof: Exercise.* ■

**Theorem.** If  $x_n \rightarrow x$ , then  $|x_n| \rightarrow |x|$

*Sketch of the proof. :*

$$||x_n| - |x|| \leq |x_n - x|$$

■

**Theorem.** Suppose  $(x_n)$  is a sequence of non-negative real numbers, satisfying  $x_n \rightarrow x$ . Then  $\sqrt{x_n} \rightarrow \sqrt{x}$ .

*Proof.* Let  $\forall \varepsilon > 0$

**Case1:**  $x = 0$

$\exists N \in \mathbb{N}$  s.t.  $\forall n \geq N$ ,

$$\begin{aligned} |x_n - 0| &< \varepsilon^2 \\ \Rightarrow |\sqrt{x_n} - 0| &< \varepsilon \end{aligned}$$

**Case2:**  $x > 0$

$\exists N$  s.t.  $\forall n > N$ ,

$$|x_n - x| < \sqrt{x} \cdot \varepsilon$$

Notice that

$$\begin{aligned} |\sqrt{x_n} - \sqrt{x}| &= \left| \frac{x_n - x}{\sqrt{x_n} + \sqrt{x}} \right| \\ &\leq \frac{|x_n - x|}{\sqrt{x}} < \varepsilon \end{aligned}$$

■

**Theorem.** Let  $(x_n)$  be a sequence of positive real numbers such that  $L = \lim(\frac{x_{n+1}}{x_n})$  exists. If  $L < 1$ , then  $(x_n)$  converges to 0.

*Proof.*  $\exists N_1 \in \mathbb{N}$  s.t.  $\forall n > N_1$

$$\left| \frac{x_{n+1}}{x_n} - L \right| \leq \left| \frac{x_{n+1}}{x_n} - L \right| < \frac{1-L}{2}$$

Thus

$$\left| \frac{x_{n+1}}{x_n} \right| < \frac{1+L}{2}$$

Note that  $\frac{1+L}{2} < 1$ , write  $r = \frac{1+L}{2}$ . Then  $\forall m \in \mathbb{N}$ ,

$$x_{N_1+m} < x_{N_1+(m-1)}r < x_{N_1+(m-2)}r^2 < \cdots < x_{N_1}r^m$$

Consider  $(y_m) = (x_{N_1+m})$ ,  $(z_m) = (x_{N_1}r^m)$ . Then

$$0 \leq y_m \leq z_m$$

Since  $z_m \rightarrow 0$

$(y_m) \rightarrow 0$  by squeeze theorem

Thus we conclude that  $(x_n) \rightarrow 0$  by m-th tail theorem.

■

### 3.3 Monotone sequence

**Definition.** Let  $(x_n)$  be a sequence of real. We say  $(x_n)$  is ...

- **increasing** if  $\forall n \in \mathbb{N}, x_{n+1} \geq x_n$ .
- **strictly increasing** if  $\forall n \in \mathbb{N}, x_{n+1} > x_n$ .
- **decreasing** if  $\forall n \in \mathbb{N}, x_{n+1} \leq x_n$ .
- **strictly decreasing** if  $\forall n \in \mathbb{N}, x_{n+1} < x_n$ .

**Theorem (Monotone Convergence Theorem).** A monotone sequence of real numbers is **convergent** iff it is bounded. Moreover, if  $(x_n)$  is increasing, then

$$\lim(x_n) = \sup \{x_n : n \in \mathbb{N}\}$$

If  $(x_n)$  is decreasing, then

$$\lim(x_n) = \inf \{x_n : n \in \mathbb{N}\}$$

*Proof.* ( $\Rightarrow$ )

A convergent sequence is always bounded.

( $\Leftarrow$ )

Suppose  $(x_n)$  is a monotone and bounded sequence.

**Case 1:**  $(x_n)$  is increasing.

Write  $x = \sup \{x_n : n \in \mathbb{N}\}$ .

Let  $\varepsilon > 0$ . Since  $x = \sup(x_n)$ :

$$x - \varepsilon \text{ is NOT an upper bound of } (x_n)$$

Then

$$\exists N \in \mathbb{N} \text{ s.t. } x_N > x - \varepsilon$$

Since  $(x_n)$  is increasing,

$$\forall n \geq N, x_n > x - \varepsilon$$

On the other hand,  $x + \varepsilon$  is an upper bound since  $x$  is an upper bound. Thus,

$$x_n < x + \varepsilon$$

$$\Rightarrow \forall n \geq N, x - \varepsilon < x_n < x + \varepsilon$$

$$\Rightarrow |x_n - x| < \varepsilon \Rightarrow (x_n) \rightarrow x$$

**Case 2:**  $(x_n)$  is decreasing.

Write  $y = \inf(x_n)$ . Let  $\varepsilon > 0$

Since  $y = \inf(x_n)$ ,

$$y + \varepsilon \text{ is NOT an upper bound of } (x_n)$$

Thus

$$\exists N \in \mathbb{N} \text{ s.t. } x_n < y + \varepsilon$$

Since  $(x_n)$  is decreasing,

$$\forall n > N, x_n < y + \varepsilon$$

On the other hand,  $y - \varepsilon$  is a lower bound since  $y$  is a lower bound. Hence,

$$\forall n \in \mathbb{N}, y - \varepsilon < x_n$$

$$\begin{aligned} \forall n \geq N, y - \varepsilon < x_n < y + \varepsilon &\Rightarrow |x_n - y| < \varepsilon \\ &\Rightarrow (x_n) \rightarrow y \end{aligned}$$

**Remark.** One can prove case 2 by following:

$(-x_n)$  is increasing and converges to  $\sup(-x_n)$  by case 1. Also note that

$$(x_n) = (-(-x_n)) \rightarrow -\sup(-x_n)$$

by limit law. So it is easy to prove that

$$-\sup(-x_n) = \inf(x_n)$$

■

**Example.** Consider the sequence  $(x_n)$  is given by

$$\begin{cases} x_0 = \frac{1}{2} \\ x_{n+1} = \frac{3}{2}x_n(1 - x_n) \end{cases}$$

$(x_n)$  is decreasing and bounded.

**Thoughts:** Assume  $(x_n)$  converges, then by limit law,

$$x = \frac{3}{2}x(1 - x) \text{ where } x = \lim(x_n) \Rightarrow x = 0 \text{ or } \frac{2}{3}$$

then, by proof of contradiction, it is not convergent.

*Proof.* **Claim:**  $\frac{1}{3} < x_{n+1} < x_n \leq \frac{1}{2}, \forall n \in \mathbb{N} \cup \{0\}$

**Proof of the claim by induction:**

When  $n = 0$ :

$$\begin{aligned} x_0 &= \frac{1}{2}, x_1 = \frac{3}{2} \cdot \frac{1}{2} \left(1 - \frac{1}{2}\right) = \frac{3}{8} \\ \frac{1}{3} &< \frac{3}{8} < \frac{1}{2} \leq \frac{1}{2} \end{aligned}$$

Suppose this is true for  $n=k$ :

$$\frac{1}{3} < x_{k+1} < x_k \leq \frac{1}{2}$$

Goal:

$$\begin{aligned}\frac{1}{3} < x_{k+2} < x_{k+1} &\leq \frac{1}{2} \\ x_{k+1} &= \frac{3}{2}x_k(1-x_k) \\ \frac{1}{3} < x_k \leq \frac{1}{2} &\Rightarrow \frac{2}{3} > 1-x_k \geq \frac{1}{2} \\ x_{k+1} &< \frac{3}{2} \cdot \frac{1}{2} \cdot \frac{2}{3} = \frac{1}{2}\end{aligned}$$

Complete the square:

$$\begin{aligned}x_{k+1} - \frac{1}{3} &= \frac{3}{2}x_k(1-x_k) - \frac{1}{3} \\ &= -\frac{3}{2}\left[\left(x_k - \frac{1}{2}\right)^2 - \frac{1}{36}\right]\end{aligned}$$

So

$$\begin{aligned}\frac{1}{3} < x_k \leq \frac{1}{2} &\Rightarrow \left|x_k - \frac{1}{2}\right| < \frac{1}{6} \\ &\Rightarrow \left(x_k - \frac{1}{2}\right)^2 < \frac{1}{36} \\ &\Rightarrow x_{k+1} - \frac{1}{3} > 0 \\ &\Rightarrow \frac{1}{3} < x_{k+1} \leq \frac{1}{2}\end{aligned}$$

With the similar process, we can derive that

$$\begin{aligned}\frac{1}{3} < x_{k+2} &\leq \frac{1}{2} \\ x_{k+2} &= \frac{3}{2}x_{k+1}(1-x_{k+1}) < \frac{3}{2}x_{k+1} \cdot \frac{3}{2} = x_{k+1}\end{aligned}$$

Therefore, this claim is also true for  $n=k+1$ :

$$\frac{1}{3} < x_{k+2} < x_{k+1} \leq \frac{1}{2}$$

We thereby prove the theorem by induction:

$$\frac{1}{3} < x_{n+1} < x_n \leq \frac{1}{2}$$

■

**Exercise:** Textbook p.75: A sequence that converges to  $\sqrt{a}$  for  $a > 0$ .

**Definition (Euler's Number).**

$$e = \lim(1 + (\frac{1}{n})^n)$$

**Goal:**  $(x_n)$  is convergent where  $x_n = (1 + \frac{1}{n})^n$

$$\begin{aligned} x_n &= (1 + \frac{1}{n})^n = 1 + nC1 \cdot \frac{1}{n} + nC2 \cdot \frac{1}{n^2} + \cdots + nCn \frac{1}{n^n} \\ &= 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdot 3 \cdot 2 \cdot 1}{n!} \cdot \frac{1}{n^2} \\ &= 1 + 1 + \frac{1}{2}(1 - \frac{1}{n}) + \frac{1}{6}(1 - n)(1 - \frac{2}{n}) + \cdots + \frac{1}{n!}(1 - \frac{1}{n})(1 - \frac{2}{n}) \cdots \frac{2}{n} \cdot \frac{1}{n} \end{aligned}$$

Write  $x_{n+1}$  in a similar way, we observe that

$$x_n < x_{n+1}$$

**Facts**  $2^{m-1} \leq m!$  for  $m \in \mathbb{N} \Rightarrow \frac{1}{m!} \leq \frac{1}{2^{m-1}}$

$$x_n < 1 + 1 + \frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^{n-1}} = 1 + \frac{1(1 - (\frac{1}{2})^n)}{1 - \frac{1}{2}} < 3$$

$\Rightarrow (x_n)$  is increasing and bounded



### 3.4 Subsequence and the Bolzano-Weierstrass Theorem

**Example.**

$$\begin{aligned}(x_n) &= ((-1)^n) \\ x_{2n} &= (-1)^{2n} \\ x_{2n+1} &= (-1)^{2n-1}\end{aligned}$$

So  $a_n = x_{2n}$  is a sequence, while  $b_n = x_{2n+1}$  is a subsequence of  $x_n$ .

**Definition (Subsequences).** Let  $(x_n)$  be a real sequence and consider a strictly increasing sequence of natural numbers  $n_1 < n_2 < n_3 < \dots$ . The sequence

$$(x_{n_k} : k \in \mathbb{N})$$

is called a **subsequence** of  $(x_n)$

**Example.** Any tails of a sequence is a subsequence:  $(x_n)$   $n$ -th tail:  $(x_{m+k})$ , where  $n = m + k, k \in \mathbb{N}$ , is a subsequence.

**Theorem.** Suppose  $(x_n)$  converges to  $x$ . Then  $x_{n_k} \rightarrow x$  for any subsequence of  $(x_n)$ .

*Proof.* Let  $\varepsilon > 0$ , then

$$\exists N_\varepsilon > 0 \text{ s.t. } |x_n - x| < \varepsilon \text{ for } n > N_\varepsilon.$$

Note that

$$n_k \geq k, \forall k \in \mathbb{N}.$$

**Exercise.** By induction,  $n_1 \geq 1, n_2 \geq n_1 \geq 1 \Rightarrow n_2 \geq 2$

When  $k > N_\varepsilon$ ,  $n_k > N_\varepsilon$ , thus

$$|x_{n_k} - x| < \varepsilon$$

Therefore

$$(x_{n_k}) \rightarrow x$$

■

**Theorem.** Let  $(x_n)$  be a sequence of real numbers, and let  $x \in \mathbb{R}$ . Then the following are equivalent:

1.  $(x_n)$  does not converge to  $x$ .
2.  $\exists \varepsilon_0 > 0$ , s.t.  $\forall k \in \mathbb{N}, \exists n_k \in \mathbb{N}$  s.t.

$$n_k \geq k \text{ \& } |x_{n_k} - x| > \varepsilon_0$$

3.  $\exists \varepsilon_0 > 0$  and a subsequence  $(x_{n_k})$  s.t.

$$|x_{n_k} - x| > \varepsilon_0, \forall k \in \mathbb{N}$$

**Example.**

$$x_n = (-1)^n \Rightarrow (x_n) \text{ does not converge to } 1$$

*Proof.*

$3 \rightarrow 1$ : by the contrapositive statement of definition

$3 \rightarrow 2$ : 3 is a stronger statement of 2

$1 \rightarrow 2$ : left as exercise

■

**Theorem.** If  $(x_n)$  satisfies either of the following property, then it is **divergent**:

1. There exists two subsequence  $(x_{n_k})$  &  $(x_{m_k})$  whose limits are NOT equal.
2.  $(x_n)$  is unbounded.

**Example.**

1.  $(-1)^n$
2.  $(n)$
3.  $(x_n)$  such that

$$\begin{aligned} x_{2k} &= k \\ x_{2k+1} &= (-1)^k \end{aligned}$$

*Proof. exercise*

■

**Theorem (Bolzano-Weierstrass Theorem).** A bounded sequence of real numbers has a **convergent subsequence**.

**Example.**


$$x_n = (-1)^n$$

*Proof.*

**Lemma.** If  $(x_n)$  is a sequence of real numbers, there exists a subsequence of  $(x_n)$  which is monotone.

**proof of lemma:**

Call the  $m$ -th term  $x_m$  a "*peek*" if  $x_m$  is at least as large as any term after it in the sequence.



**Case 1:**  $(x_n)$  has infinitely many peaks

List the peaks of  $(x_n)$  in order of increasing index

$$x_{n_1}, x_{n_2}, \dots$$

$\Rightarrow (x_{n_i})$  is a decreasing sequence.

**Case 2:**  $(x_n)$  has a finite number of peaks

Let  $s_1$  be the first index after the last peak of  $(x_n)$ . Then for every  $n \geq s_1$ ,  $\exists m \in \mathbb{N}$  such that  $x_m > x_n$ .

Choose

$$s_2 \geq s_1 \text{ such that } x_{s_2} > x_{s_1}, s_2 \geq s_1 \text{ such that } x_{s_2} > x_{s_1}, \dots$$

$$\Rightarrow (x_{s_i}) \text{ is an increasing sequence.}$$

**Remark.** Lemma + monotone convergent theorem implies Bolzano-Weierstrass theorem.

### Second proof

Suppose  $(x_n)$  is a bounded sequence.

$$\Rightarrow \exists I_1 = [a_1, b_1] \text{ such that } (x_n) \in I_1$$

Consider

$$I'_2 = [a_1, \frac{a_1 + b_1}{2}], I''_2 = [\frac{a_1 + b_1}{2}, b_1]$$

Let  $I_2 = [a_2, b_2]$  be one of  $I'_2, I''_2$  such that  $I_2$  contains infinitely many terms of  $(x_n)$ .

For  $n \in \mathbb{N}$ , define  $I_n = [a_n, b_n]$  in a similar way.

For  $i \in \mathbb{N}$ , choose a term  $x_{n_i}$  such that  $x_{n_i} \in I_i$  and  $n_i > n_{i-1}$

Then

$$\begin{array}{ll} i = 1, & n_i = 1 \\ i = 2, & \text{choose } n_2 \in \mathbb{N} \text{ such that } n_2 > n_1 \text{ \& } x_{n_2} \in I_2 \\ \vdots & \vdots \end{array}$$

- $\forall i \in \mathbb{N}, a_i \leq x_{n_i} \leq b_i$
- $(a_i)$  increases, bounded above by  $b_1 \Rightarrow (a_i) \rightarrow \sup(a_i)$
- $(b_i)$  decreases, bounded below by  $a_1 \Rightarrow (b_i) \rightarrow \inf(b_i)$
- $\inf |a_i - b_i| = \inf \frac{b_1 - a_1}{2^n} = n \Rightarrow \sup(a_i) = \inf(b_i)$

Thus  $(x_{n_i})$  is convergent by squeeze theorem. ■

**Theorem.** Let  $(x_n)$  be a bounded sequence, and  $x \in \mathbb{R}$  has the property that every convergent subsequence of  $(x_n)$  converges to  $x$ . Then  $x_n$  converges to  $x$

*Proof.* Let  $\forall \varepsilon > 0$

By Bolzano-Weierstrass theorem,  $\exists$  a convergent subsequence  $(x_{n_i})$  such that

$$\exists N_\varepsilon \in \mathbb{N} \text{ s.t. for } i > N_\varepsilon, |x_{n_i} - x| < \varepsilon$$

Assume  $(x_n)$  does not converge to  $x$ . Then by previous theorem of subsequence,

$$\exists \varepsilon_0 > 0 \text{ and a subsequence } (x_{n_k}) \text{ s.t. } |x_{n_k} - x| > \varepsilon_0, \forall k \in \mathbb{N}$$

Since  $(x_n)$  is bounded,  $(x_{n_k})$  is also bounded. Thus there exists a convergent subsequence of  $(x_{n_k})$  as  $(x_{n_{k_i}})$ .

Note that  $(x_{n_{k_i}})$  is a convergent subsequence of  $(x_n)$ , thus

$$(x_{n_{k_i}}) \rightarrow x \text{ which is contradiction to previous assumption}$$

■

**Definition.** Let  $(x_n)$  be a sequence of real numbers. A point is called a **subsequential limit** of  $(x_n)$  if it is the limit of a subsequence of  $(x_n)$ .

$$S = \{\alpha \in \mathbb{R} : \alpha \text{ is a subsequential limit}\} \text{ NOTE: may be infinite set}$$

**Definition.** Let  $(x_n)$  be a sequence of real numbers.

- The **limit superior** of  $(x_n)$  is the infimum of the set of  $v \in \mathbb{R}$  s.t.  $v < x_n$  for at most a finite number of  $n \in \mathbb{N}$ . We write it as

$$\limsup(x_n) \text{ or } \limsup x_n \text{ or } \overline{\lim} x_n$$

- The **limit inferior** of  $(x_n)$  is the supremum of the set of  $v \in \mathbb{R}$  s.t.  $w > x_n$  for at most a finite number of  $n \in \mathbb{N}$ . We write it as

$$\liminf(x_n) \text{ or } \liminf x_n \text{ or } \underline{\lim} x_n$$

### **Intuition**

- Suppose  $v < x_n$  for at most finitely many  $n \in \mathbb{N}$ , then

**Theorem.** Let  $(x_n)$  be a bounded sequence, and  $x \in \mathbb{R}$  has the property that every convergent subsequence of  $(x_n)$  converges to  $x$ . Then  $x_n$  converges to  $x$

*Proof.* Let  $\forall \varepsilon > 0$

By Bolzano-Weierstrass theorem,  $\exists$  a convergent subsequence  $(x_{n_i})$  such that

$$\exists N_\varepsilon \in \mathbb{N} \text{ s.t. for } i > N_\varepsilon, |x_{n_i} - x| < \varepsilon$$

Assume  $(x_n)$  does not converge to  $x$ . Then by previous theorem of subsequence,

$$\exists \varepsilon_0 > 0 \text{ and a subsequence } (x_{n_k}) \text{ s.t. } |x_{n_k} - x| > \varepsilon_0, \forall k \in \mathbb{N}$$

Since  $(x_n)$  is bounded,  $(x_{n_k})$  is also bounded. Thus there exists a convergent subsequence of  $(x_{n_k})$  as  $(x_{n_{k_i}})$ .

Note that  $(x_{n_{k_i}})$  is a convergent subsequence of  $(x_n)$ , thus

$$(x_{n_{k_i}}) \rightarrow x \text{ which is contradiction to previous assumption}$$

■

**Definition.** Let  $(x_n)$  be a sequence of real numbers. A point is called a **subsequential limit** of  $(x_n)$  if it is the limit of a subsequence of  $(x_n)$ .

$$S = \{\alpha \in \mathbb{R} : \alpha \text{ is a subsequential limit}\} \text{ (NOTE: may be infinite set)}$$

**Example.** Consider  $(x_n) = \{(-1)^n | n \in \mathbb{N}\}$ . Then


$$S \supseteq \{1, -1\}$$

**Definition** (lim sup and lim inf). Let  $(x_n)$  be a sequence of real numbers.

- The **limit superior** of  $(x_n)$  is the infimum of the set of  $v \in \mathbb{R}$  s.t.  $v < x_n$  for at most a finite number of  $n \in \mathbb{N}$ . We write it as

$$\limsup(x_n) = \limsup x_n = \overline{\lim} x_n = \inf \{v \in \mathbb{R} | v < x_n \text{ for at most a finite number of } n\}$$

**Example.** Consider  $(x_n) = \frac{1}{n}$



Let


$$X = \{v \in \mathbb{R} | v < x_n \text{ for at most a finite number of } n\}$$

- $-1 \notin X$  because there are infinitely many  $x_n$  such that  $v < x_n$ .
- $\frac{1}{2} \in X$  because there are finitely many  $x_n$  such that  $v < x_n$ .
- $2 \in X$  because there is no  $x_n$  such that  $v < x_n$ , which is smaller than finite and thereby satisfies the definition.

We thus conclude that

$$(0, \infty) \subset X \text{ and } \limsup x_n = \inf X$$

**Example.** Consider  $(x_n) = (-1)^n$ :



- $1 \in X$  because there is no  $x_n$  such that  $v < x_n$ .
- $2 \in X$  because there is no  $x_n$  such that  $v < x_n$ .
- $0, -1 \notin X$  because there are infinitely many  $x_n$  such that  $v < x_n$ .

We thus conclude that

$$[1, \infty) \subset X \text{ (in fact they are equal)}$$

- The **limit inferior** of  $(x_n)$  is the supremum of the set of  $w \in \mathbb{R}$  s.t.  $w > x_n$  for at most a finite number of  $n \in \mathbb{N}$ . We write it as

$$\liminf(x_n) \text{ or } \liminf x_n \text{ or } \overline{\lim} x_n = \sup \{w \in \mathbb{R} | w > x_n \text{ for at most a finite number of } n\}$$

**Intuition**

- Suppose  $v < x_n$  for at most finitely many  $n \in \mathbb{N}$ , then for all large  $n$ ,  $v \geq x_n$ .  
 $\Rightarrow$  No subsequential limit of  $(x_n)$  can possibly exceed  $v$ .
- Similar observation for  $\underline{\lim} x_n$

**Theorem.** Let  $(x_n)$  be a bounded sequence of real numbers, and let  $x^* \in \mathbb{R}$ . Then TFAE:

1.  $x^* = \limsup(x_n)$
2. If  $\varepsilon > 0$ , there are at most a finite number of  $n \in \mathbb{N}$  s.t.  $x^* + \varepsilon < x_n$ , but infinitely many  $n$  for which  $x^* - \varepsilon < x_n$
3. If  $u_m = \sup \{x_n | n \geq m\}$  (sup of  $(m-1)$ -th tail), then  $x^* = \inf \{u_m | m \in \mathbb{N}\} = \lim u_m$
4. If  $S$  is the set of subsequential limits of  $x_n$ , then  $x^* = \sup S$ .

**Remark.** .

- $u_m$  is decreasing.
- There is a similar such list of equivalent properties for  $\liminf$ .

**Corollary.** A bounded sequence  $(x_n)$  is convergent iff  $\overline{\lim} x_n = \lim x_n$

*Proof.* A direct result of the theorem:

$$\overline{\lim} x_n = \sup S \text{ and } \underline{\lim} x_n = \inf S$$

■

*Proof of thm. (a)  $\Rightarrow$  (b).* Let  $\varepsilon > 0$ . Then

$$x^* + \varepsilon > x^* = X = \inf \{v \in \mathbb{R} | v < x_n \text{ for at most a finite number of } n.\}$$

$$\Rightarrow \exists v \in \mathbb{R} \text{ s.t. } x^* \leq v < x^* + \varepsilon$$

and there are only finitely many  $n$  with  $v < x_n$ .

For any  $n$  for which  $x^* + \varepsilon < x_n$ ,  $v < x_n$ . Thus there are only finitely many such  $n$ .

If  $x^* - \varepsilon \notin X$ , then there are infinitely many  $n$  such that  $x^* - \varepsilon < x_n$

■

*Proof of thm. (b)  $\Rightarrow$  (c).* Fix  $\varepsilon > 0$ .

By (b), there are only finitely many  $n$  with  $x^* + \varepsilon < x_n$ .

Take  $N \in \mathbb{N}$  large enough such that

$$\begin{aligned} & x^* + \varepsilon \geq x_n && \forall n \geq N \\ \Rightarrow & x^* + \varepsilon \geq u_N \\ \Rightarrow & x^* + \varepsilon \geq \lim u_n \\ \Rightarrow & x^* \geq \lim u_n && \forall n \geq N \end{aligned}$$

On the other hand, there are infinitely many  $n$  with  $x^* - \varepsilon < x_n \leq u_n$ .

Thus, there exists a subsequence of  $u_n$ , say  $u_{n_k}$ , satisfies

$$\begin{aligned} & x^* - \varepsilon \leq u_{n_k} \\ \Rightarrow & x^* - \varepsilon \leq \lim u_{n_k} = \lim (u_n) \\ \Rightarrow & x^* \leq \lim (u_n) \end{aligned}$$

■

*Proof of thm. (c)  $\Rightarrow$  (d).* **Goal:**

$$\begin{aligned} & x^* = \lim (u_m), u_m = \sup \{x_n | n \geq m\} \\ \Rightarrow & x^* = \sup S \text{ where } S \text{ is the set of subsequential limits.} \end{aligned}$$

Let  $(x_{n_k})$  be a convergent subsequence of  $(x_n)$ . Notice that  $\lim (x_{n_k}) \in S$ .

$$\begin{aligned} & n \geq k \\ \Rightarrow & x_{n_k} \leq \sup \{x_n | n \geq k\} = u_k \\ \Rightarrow & \lim (x_{n_k}) \leq \lim (u_k) = x^* \\ \Rightarrow & x^* \text{ is an upper bound of } S. \end{aligned}$$

For 1,  $\exists n_1 \in \mathbb{N}$  s.t.

$$u_1 - 1 \leq x_{n_1} \leq u_1$$

For  $\frac{1}{2}$ ,  $\exists n_2 \in \mathbb{N}$  s.t.

$$\begin{aligned} u_2 - \frac{1}{2} & \leq x_{n_2} \leq u_2 \\ & \dots \end{aligned}$$

For  $\frac{1}{k}$ ,  $\exists n_k \in \mathbb{N}$  s.t.

$$u_k - \frac{1}{k} \leq x_{n_k} \leq u_k$$

When  $k \rightarrow \infty$ ,

$$x^* - 0 \leq \lim (x_{n_k}) \leq x^*$$

By squeeze theorem,

$$\lim (x_{n_k}) = x^*$$

■

*Proof of thm. (d)  $\Rightarrow$  (a). **Goal:***

$$x^* = \sup S$$

$$\Rightarrow x^* = \limsup x_n = \inf \{v \in \mathbb{R} | v < x_n \text{ for at most finite many of } n\}$$

Fix  $\varepsilon > 0$

There is no subsequence of  $x_n$  which has a limit exceeding  $x^* + \varepsilon$ .

$\Rightarrow$  There is only finitely many  $n$  with  $x_n > x^* + \varepsilon$ .

$\Rightarrow$

$$x^* + \varepsilon \in X$$

$\Rightarrow$

$$\inf X \leq x^* + \varepsilon$$

$\Rightarrow$

$$\limsup x_n \leq x^* + \varepsilon$$

$\Rightarrow$

$$\limsup x_n \leq x^*$$

Next, consider  $x^* - \varepsilon$

Then, there exists a subsequential limit of  $x_n$  which is greater or equal to  $x^* - \frac{1}{2}\varepsilon$ .

There exists a convergent subsequence of  $(x_n)$ , say  $(x_{n_k})$ , such that

$$\lim (x_{n_k}) \geq x^* - \frac{1}{2}\varepsilon$$

$\Rightarrow$

There are infinitely many  $n$  with  $x_n > x^* - \varepsilon$ .

$\Rightarrow$

$$\forall a \in X, x^* - \varepsilon \leq a$$

$\Rightarrow$

$$x^* - \varepsilon \leq \inf X$$

$\Rightarrow$

$$\limsup x_n \geq x^* - \varepsilon$$

$\Rightarrow$

$$\limsup x_n \geq x^*$$

In conclusion,

$$\limsup x_n = x^*$$

■



### 3.5 Cauchy Criterion

**Definition (Cauchy Sequence).** A sequence  $(x_n)$  is Cauchy sequence if

$$\forall \varepsilon > 0, \exists H \in \mathbb{N} \text{ s.t. } \forall n, m \in \mathbb{N}, n > 0, m > 0,$$

$$|x_n - x_m| < \varepsilon$$

**Example.**  $(\frac{1}{n})$  is a Cauchy sequence.

*Proof.* Observe that  $\forall n, m \in \mathbb{N}, n \geq m$ ,

$$\left| \frac{1}{n} - \frac{1}{m} \right| = \left| \frac{m - n}{mn} \right| = \frac{n - m}{mn} \leq \frac{n}{mn} < \frac{1}{m}$$

Choose  $H = \lceil \frac{1}{\varepsilon} \rceil + 1$ , Then

$$\forall n, m \geq H,$$

$$\left| \frac{1}{n} - \frac{1}{m} \right| \leq \frac{1}{m} < \varepsilon$$

■

**Example.**  $(-1)^n$  is not a Cauchy sequence.

*Proof.* Choose  $\varepsilon_0 = \frac{1}{2}$ ,  $\forall H \in \mathbb{N}$ , choose  $n, m \geq H$  s.t.  $n$  is even,  $m$  is odd. Then

$$|(-1)^n - (-1)^m| = |1 - (-1)| = 2 > \frac{1}{2}$$

Thus  $(-1)^n$  does not satisfies the definition of Cauchy sequence.

■

**Theorem.** If  $(x_n)$  is convergent, then it is Cauchy sequence.

*Proof.* Let  $\lim (x_n) = x$ .

Goal:

$$|x_n - x| < \varepsilon \Rightarrow |x_n - x_m| < \varepsilon$$

$$\forall n, m \geq N_\varepsilon,$$

$$\begin{aligned} |x_n - x_m| &= |(x_n - x) + (x - x_m)| \\ &\leq |x_n - x| + |x - x_m| \text{ by triangle inequality} \\ &< \frac{1}{2}\varepsilon + \frac{1}{2}\varepsilon = \varepsilon \end{aligned}$$

■

**Lemma.** If  $(x_n)$  is Cauchy, then it is bounded.

*Proof.* Choose  $\varepsilon = 1, \exists H \geq 0$  s.t.  $\forall n, m \geq H$ ,

$$|x_n - x_m| < 1$$

Since the choice of  $m$  satisfies  $m \geq H$ , we may choose  $m = H$  s.t.

$$|x_n - x_H| < 1$$

It follows that

$$|x_n| - |x_H| \leq |x_n - x_H| < 1$$

$$|x_n| < |x_H| + 1, \forall n \geq H$$

Let

$$M := \max \{|x_1|, |x_2|, |x_3|, \dots, |x_{H-1}|, |x_H| + 1\}$$

Thus  $\forall n \in \mathbb{N}, |x_n| \leq M$  ■

**Theorem (Cauchy Convergence Theorem).** A sequence of real numbers is convergent if and only if it is Cauchy sequence.

$(\Rightarrow)$ . Done in the previous theorem. ■

$(\Leftarrow)$ . Suppose  $(x_n)$  is Cauchy. By lemma, it is bounded.

By Bolzano-Weierstrass theorem, there exists a convergent subsequence  $(x_{n_k})$ .

Let  $\lim(x_{n_k}) = x$ .

**Goal:**  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$  s.t.  $\forall n \geq N$

$$|x_n - x| < \varepsilon$$

We will use the trick of insert subsequence:

By definition of convergence,  $\forall \varepsilon > 0, \exists N_\varepsilon \in \mathbb{N}$  s.t.  $\forall k \geq N_\varepsilon$

$$|x_{n_k} - x| < \frac{1}{2}\varepsilon$$

By definition of Cauchy sequence,  $\forall \varepsilon > 0, \exists H \in \mathbb{N}, H \geq 0$  s.t.  $\forall n, m \geq H$

$$|x_n - x_m| < \frac{1}{2}\varepsilon$$

Thus  $\forall k \geq \max\{H, N_\varepsilon\}$ ,

$$\begin{aligned} |x_k - x| &= |(x_k - x_{n_k}) + (x_{n_k} - x)| \\ &\leq |x_k - x_{n_k}| + |x_{n_k} - x| \quad \text{since } n_k \geq k \geq \max\{H, N_\varepsilon\} \\ &< \frac{1}{2}\varepsilon + \frac{1}{2}\varepsilon = \varepsilon \end{aligned}$$

■

**Example.** Let  $(h_n)$  be the sequence of harmonic series such that

$$h_n = \sum_{i=1}^n \frac{1}{i}, \text{ and } \lim(h_n) = \sum_{n=1}^{\infty} \frac{1}{n}$$

**Claim:**  $(h_n)$  is divergent

**Goal:** show that it is NOT Cauchy.

*Proof.*  $\forall m, n \in \mathbb{N}$ , WLOG suppose  $m \geq n, h_m > h_n$ , we have

$$\begin{aligned} |h_m - h_n| &= \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{m-1} + \frac{1}{m} \\ &\geq \frac{m-n}{n} \end{aligned}$$

since there are  $(m-n)$ 's terms on the right side of the equation.

So we choose  $\varepsilon_0 = \frac{1}{2}, \forall H \geq 0$ , choose  $n = H, m = 2H$ , then

$$|h_m - h_n| \geq \frac{m-n}{m} = \frac{1}{2}$$

Thus,  $(h_n)$  is NOT Cauchy. ■

### 3.6 Property of Divergent Sequence

**Example** (divergent sequence). :

- $(n) \rightarrow \infty$
- $(-n) \rightarrow -\infty$
- $(-1)^n \cdot n$  is divergent and unbounded.
- $(-1)^n$  is divergent and bounded.

**Definition (Properly Divergent).** Let  $(x_n)$  be a sequence of real numbers. We say that

1.  $(x_n)$  **tends to**  $\infty$ , or  $\lim(x_n) = \infty$  if  $\forall a \in \mathbb{R}, \exists N \in \mathbb{N}$  s.t.  $\forall n \geq N$ ,

$$x_n > a$$

2. Similarly,  $(x_n)$  **tends to**  $-\infty$ , or  $\lim(x_n) = -\infty$  if  $\forall a \in \mathbb{R}, \exists N \in \mathbb{N}$  s.t.  $\forall n \geq N$ ,

$$x_n < a$$

In either case, we say that  $(x_n)$  is **properly divergent**.

**Example.** Let  $C > 0$ .  $(C^n)$  is properly divergent. We write  $\lim(C^n) = \infty$ .

*Proof.* Notice that

$$C^n = [1 + (C - 1)]^n \geq 1 + n(C - 1) \text{ by Bernoulli's inequality}$$

**Goal:**  $\forall a \in \mathbb{R}, \exists N \in \mathbb{N}$  s.t.  $\forall n \geq N$ ,

$$1 + n(C - 1) > a_n \Leftrightarrow n > \frac{a_n - 1}{C - 1}$$

Then we choose  $N = \lceil \frac{a-1}{C-1} \rceil + 1$ . Thus  $\forall n \geq N, C^n > a$ . ■

**Theorem.** A monotone sequence is **divergent** if and only if it is **bounded**. ■

*Proof. Exercise.* ■

**Theorem (Comparison Test).** Let  $(x_n)$  and  $(y_n)$  be two sequences. Suppose  $\forall n \in \mathbb{N}, x_n \geq y_n$ . Then,

1. If  $(x_n) \rightarrow \infty$ , then  $(y_n) \rightarrow \infty$ .
2. If  $(y_n) \rightarrow -\infty$ , then  $(x_n) \rightarrow -\infty$ .

*Proof. Exercise.* ■

**Theorem (Limit Comparison Test).** Let  $(x_n)$  and  $(y_n)$  be two sequences of positive real numbers. Suppose  $\exists L \in \mathbb{R}, L > 0$  s.t.

$$\exists \lim\left(\frac{x_n}{y_n}\right) = L$$

then  $\lim(x_n) = \infty$  if and only if  $\lim(y_n) = \infty$

*Proof.* **Claim:** For  $N$  large enough,

$$\frac{1}{2}L \cdot y_n < x_n < 2L \cdot y_n$$

**Goal:** Claim+Comparison Theorem=Proof.

By definition of limit,  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$  s.t.  $\forall n > N$ ,

$$L - \varepsilon < \frac{x_n}{y_n} < L + \varepsilon$$

Choose  $\varepsilon = \frac{1}{2}L$ , we have

$$\frac{1}{2}L \cdot y_n < x_n < \frac{3}{2}L < 2L \cdot y_n$$

since  $L > 0$  and these are positive sequences.

Thus by Comparison Theorem,

$$\lim\left(\frac{1}{2}L \cdot y_n\right) = \infty \Rightarrow \lim(x_n) = \infty$$

■

### 3.7 Introduction to Infinite Series

**Definition (Infinite Series).** If  $(x_n)$  is a sequence of real numbers, then the **infinite series** generated by  $(x_n)$  is the sequence  $(s_k)$  defined by

$$s_k = \sum_{i=1}^k x_i$$

Terms  $s_k$  are called **partial sums**.

**Notation.**  $\sum x_i$  to mean this series or its limit at infinity  $\lim(x_k)$ .

**Theorem (Cauchy Criterion for Series).** The series  $\sum x_i$  **converges** if and only if  $\forall \varepsilon > 0, \exists M \in \mathbb{N}$  s.t.  $\forall n, m \in \mathbb{N}, n > m \geq M$ ,

$$|x_{m+1} + x_{m+2} + \cdots + x_{n-1} + x_n| < \varepsilon$$

Or we write it as

$$|s_k - s_m| < \varepsilon$$

*Proof: Exercise.* ■

**Theorem (Montone Convergence for Series).** Let  $(x_n)$  be a sequence of non-negative real numbers. Then the series  $\sum x_n$  **converges** if and only if  $(s_k)$  is bounded.

*Proof. Exercise.* ■

**Example.**  $\sum \frac{1}{n^2}$  is convergent.

*Proof. Goal:* find convergent subsequence  $(s_{k_j})$ .

Consider subsequence  $(s_{k_j})$  where  $k_j = 2^j - 1$ .

Observe that:

$$\begin{aligned} s_{k_1} &= 1 \\ s_{k_2} &= 1 + \frac{1}{2^2} + \frac{1}{3^2} \\ &< s_{k_1} + 2 \cdot \frac{1}{2^2} = 1 + \frac{1}{2} \\ s_{k_3} &= 1 + \left(\frac{1}{2^2} + \frac{1}{3^2}\right) + \left(\frac{1}{4^2} + \frac{1}{5^2} + \cdots + \frac{1}{9^2}\right) \\ &< s_{k_1} + 2 \cdot \frac{1}{2^2} = 1 + \frac{1}{2} \end{aligned}$$

By induction(details are left as exercise), one can show that

$$s_{k_j} < \sum_{n=0}^{j-1} \frac{1}{2^n} < \sum_{n=0}^{\infty} \frac{1}{2^n} = 2 \text{ (by limit of geometric series.)}$$

Thus  $(s_{k_j})$  is bounded.

By theorem proved in homework, an increasing sequence with bounded(and thus convergent) subsequence implies that the sequence is convergent. ■

## 4 Limits

### 4.1 Limits of Functions

Let  $f : A \rightarrow B$  be a function where  $A, B \subseteq \mathbb{R}$ . Let  $a \in A, L \in B$ .

**Goal:** Define

$$\lim_{x \rightarrow a} f(x) = ?$$

**Intuition:** Define *closeness* on real line.


**Definition (Cluster Point).** Let  $A \subseteq \mathbb{R}$ . A point  $c \in \mathbb{R}$  is a **cluster point** of  $A$  if

$\forall \delta > 0, \exists x \in A, x \neq c$  such that

$$|x - c| < \delta$$

Or

$$V_\delta(c) \cap (A \setminus \{c\}) \neq \emptyset$$



**Theorem.**  $c \in \mathbb{R}$  is a cluster point if and only if

there exists a sequence  $(a_n) \in A$  such that

$$\lim(a_n) = c \text{ and } \forall n \in \mathbb{N}, a_n \neq c$$

*Sketch of Proof.* ( $\Rightarrow$ )

$$\delta = 1, \exists a_1 \in A \setminus \{c\} \text{ s.t. } |a_1 - c| < 1$$

$$\delta = \frac{1}{2}, \exists a_2 \in A \setminus \{c\} \text{ s.t. } |a_2 - c| < \frac{1}{2}$$

... observe that:

$$\forall \delta > 0, \exists a_n \in A \setminus \{c\} \text{ s.t. } |a_n - c| < \frac{1}{n} < \delta$$

$$\Rightarrow \lim(a_n) = c \text{ by squeeze theorem.}$$

( $\Leftarrow$ )

$\forall \delta > 0, \exists N_\varepsilon > 0$  s.t.

$$|a_{N_\varepsilon} - c| < \delta$$

Note that  $a_{N_\varepsilon} \in A \setminus \{c\}$  where  $c$  is a cluster point of  $A$ . ■

**Example.** Let  $X$  be the set of cluster points of  $A$ .

- $A = (1, 2) \cup (3, 4) \Rightarrow X = [1, 2] \cup [3, 4]$ . *Proof: Exercise.*
- $A = \{0\} \cup (1, 2) \Rightarrow X = \phi$ .

*Sketch of (2). :*

1. Let  $x \in [1, 2]$ . Prove that  $x \in X$ . Thus  $[1, 2] \in X$ .
2. Prove that  $0 \notin X$ .
3. Prove that  $x \in X$  if  $x \notin \{0\} \cup [1, 2]$ .

■

**Remark.**  $A$  may not be a subset of the set of cluster points of  $A$ .

**Example. :**

- $A = \mathbb{Z} \Rightarrow X = \phi$ .
- $A = \{\frac{1}{n} | n \in \mathbb{N}\} \Rightarrow X = \{0\}$  *Proof: Exercise.*

**Definition (Delta-Epsilon Definition of Limit).** Let  $A \subseteq \mathbb{R}$ ,  $c$  is a cluster point of  $A$ ,  $f : A \rightarrow \mathbb{R}$ . A real number  $L$  is the **limit of  $f$  at  $c$**  if

$\forall \varepsilon > 0, \exists \delta > 0$  s.t.  $\forall x \in A$ ,

$$0 < |x - c| < \delta \rightarrow |f(x) - L| < \varepsilon$$

**Theorem (Uniqueness of Limit).** If  $f : A \rightarrow \mathbb{R}$  and  $c$  is a cluster point of  $A$ , then  $f$  has at most 1 limit at  $c$ .

*Proof.* We will prove this by contradiction.

Let  $L_1$  and  $L_2$  be limits of  $f$  at  $c$ . Assume  $L_1 \neq L_2$ . Choose  $\varepsilon = \frac{|L_1 - L_2|}{2} > 0$ . Then,

$$\exists \delta_1 \text{ s.t. } 0 < |x - c| < \delta_1 \Rightarrow |f(x) - L_1| < \frac{\varepsilon}{2}$$

$$\exists \delta_2 \text{ s.t. } 0 < |x - c| < \delta_2 \Rightarrow |f(x) - L_2| < \frac{\varepsilon}{2}$$

Consider  $\delta := \min\{\delta_1, \delta_2\}$ .

Since  $c$  is a cluster point,  $\exists x_0 \in A$  s.t.

$$0 < |x_0 - c| < \delta$$

Since

$$|f(x_0) - L_1| < \frac{\varepsilon}{2}, |f(x_0) - L_2| < \frac{\varepsilon}{2}$$

We have

$$\begin{aligned} |L_1 - L_2| &\leq |L_1 - f(x_0)| + |f(x_0) - L_2| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \\ &= \frac{|L_1 - L_2|}{2} \end{aligned}$$

which is a contradiction.

■



**Notation.**

$$L = \lim_{x \rightarrow c} f(x) \text{ or } L = \lim_{x \rightarrow c} f$$


And we say that  $f(x)$  approaches to  $L$  as  $x$  approaches to  $c$ .

**Remark (Divergence of function).** If the limit of  $f(x)$  at  $c$  does not exist, we say that  $f$  diverges at  $c$ .

**Theorem.** Let  $f : A \rightarrow \mathbb{R}$  and  $c$  be a cluster point of  $f$ . The following are equivalent:

1.  $\lim_{x \rightarrow c} f(x) = L$
2.  $\forall V_\varepsilon(L)$   $\varepsilon$ -neighborhood of  $L$ ,  $\exists V_\delta(c)$   $\delta$ -neighborhood of  $c$  s.t.

$$x \in V_\delta(c) \cap (A \setminus \{c\}) \Rightarrow f(x) \in V_\varepsilon(L)$$



**Example.**  $\lim_{x \rightarrow c} f(x) = a$ .

*Proof.*  $\forall \varepsilon > 0, \exists \delta = 1$  s.t.

$$x \in V_\delta(c) \cap (\mathbb{R} \setminus \{c\}) \Rightarrow f(x) = a \in V_\varepsilon(L)$$

■

**Example.**  $\lim_{x \rightarrow c} f(x) = c$ .

*Proof.*  $\forall \varepsilon > 0, \exists \delta = \varepsilon$  s.t.

$$\begin{aligned} & x \in V_\delta(c) \cap (A \setminus \{c\}) \\ \Rightarrow & f(x) = x \in V_\delta(c) \cap (A \setminus \{c\}) \\ \Rightarrow & f(x) \in V_\varepsilon(c) \cap (A \setminus \{c\}) \subseteq V_\varepsilon(c) \end{aligned}$$

■

**Example.**

$$\lim_{x \rightarrow c} x^2 = c^2$$

*Proof.* **Goal:**  $\forall \varepsilon > 0$ , find a  $\delta(c, \varepsilon) > 0$  s.t.

$$\text{if } 0 < |x - c| < \delta, \text{ then } |x^2 - c^2| < \varepsilon$$

which is equivalent to show

$$|x + c| |x - c| < \varepsilon$$

Since the choice of  $\delta$  is dependent on  $\varepsilon$  and  $c$  only,  $|x - c|$  can be easily confined with some constant. Let's assume that

$$|x - c| < 1$$

Now, the only task left is to find a way to confine  $|x + c|$  with a constant by manipulating  $|x - c| < 1$ .

Here we will apply a common trick that estimates addition  $|x + c|$  with subtraction  $|x - c|$ :

$$|x| - |c| \leq |x - c| < 1 \text{ by triangle inequality.}$$

Rearranging the inequality, we have:

$$|x| < |c| + 1$$

Adding the second  $|c|$  to both side of inequality, then, we apply triangle inequality again:

$$|x + c| \leq |x| + |c| < 2|c| + 1$$

$$|x + c| < 2|c| + 1$$

Notice that this is equivalent to show

$$|x + c| |x - c| < (2|c| + \delta) |x - c| < \varepsilon$$

Rearrange the constant factor

$$|x - c| < \frac{\varepsilon}{2|c| + 1}$$

Thus, our choice of  $\delta$  must satisfies two conditions at the same time:

$$\begin{cases} |x - c| < 1 \\ |x - c| < \frac{\varepsilon}{2|c|+1} \end{cases}$$

We achieve this by simply choosing

$$\delta = \min \left\{ \frac{\varepsilon}{2|c| + 1}, 1 \right\}$$

■

**Remark (*The Toolbox of Proofs*).** The readers should develop their "toolbox" of proof techniques. That is, the ***estimation against a constant*** + manipulation of ***triangle inequality*** + choice of  $\delta$  that satisfies ***multiple conditions***.

**Example** (Harder).

$$\lim_{x \rightarrow c} \frac{1}{x} = \frac{1}{c}$$

*Proof.* Observe that

$$\left| \frac{1}{x} - \frac{1}{c} \right| = \frac{|x - c|}{|cx|}$$

Here, we cannot choose  $|x - c|$  smaller than some constant (why? try it on your own). Instead, we choose

$$|x - c| < \frac{|c|}{2}$$

By triangle inequality,

$$\frac{|c|}{2} < |x| < \frac{3|c|}{2}$$

Multiply  $|c|$  to each term of the inequality,

$$\frac{c^2}{2} < |cx| < \frac{3c^2}{2}$$

Thus

$$\frac{1}{|cx|} < \frac{2}{c^2}$$

It follows that

$$\frac{|x - c|}{|cx|} < \frac{2}{c^2} |x - c|$$

In order to make the left term of the inequality less than  $\varepsilon$ , it suffices to confine

$$\frac{2}{c^2} |x - c| < \varepsilon$$

$$|x - c| < \frac{c^2}{2} \varepsilon$$

Thus, our choice of  $\delta$  must satisfy two conditions at the same time:

$$\begin{cases} |x - c| < \frac{|c|}{2} \\ |x - c| < \frac{c^2}{2} \varepsilon \end{cases}$$

We achieve this by simply choosing

$$\delta = \min \left\{ \frac{|c|}{2}, \frac{c^2}{2} \varepsilon \right\}$$

■

**Remark** (**Factoring**  $|x - c|$ ). Since we have the premise of  $|x - c| < \delta$  for free, it would be much easier for us to confine  $|f(x) - L|$  if we factor  $|x - c|$  from the difference.

**Example** (Much Harder).

$$\forall n \in \mathbb{N}, \lim_{x \rightarrow c} x^n = c^n$$

*Proof.* By difference of  $n$ -th powers factorization

$$|x^n - c^n| = |x - c| \left| \sum_{i=0}^{n-1} x^i c^{n-1-i} \right| \leq |x - c| \cdot \sum_{i=0}^{n-1} |x|^i |c|^{n-1-i}$$

If

$$|x - c| < 1$$

then by triangle inequality,

$$|x| < |c| + 1$$

It follows that

$$|x^n - c^n| < |x - c| \cdot \sum_{i=0}^{n-1} (|c| + 1)^i |c|^{n-1-i}$$

Similar to the previous example, it suffices to confine

$$|x - c| \cdot \sum_{i=0}^{n-1} (|c| + 1)^i |c|^{n-1-i} < \varepsilon$$

$$|x - c| < \frac{\varepsilon}{\sum_{i=0}^{n-1} (|c| + 1)^i |c|^{n-1-i}}$$

Thus, choose

$$\delta = \min\left\{1, \frac{\varepsilon}{\sum_{i=0}^{n-1} (|c| + 1)^i |c|^{n-1-i}}\right\}$$

■

**Remark** (*Estimate  $x$  against  $c$* ). This is another trick by triangle inequality:

If  $\exists k \in \mathbb{R}, k > 0$  s.t.  $|x - c| < k$ , then  $|x| < |c| + k$

**Remark** (*difference of  $n$ -th powers factorization*).

$$(x^n - c^n) = (x - c)(x^{n-1} + cx^{n-2} + \cdots + c^{n-2}x + c^{n-1}) = (x - c) \sum_{i=0}^{n-1} x^i c^{n-1-i}$$

**Example (*Some Tedious Factorization*).**

$$\lim_{x \rightarrow 2} \frac{x^3 + 2x - 1}{6x^2 - 5} = \frac{11}{19}$$

*Proof.*

$$\left| \frac{x^3 + 2x - 1}{6x^2 - 5} - \frac{11}{19} \right| = \left| \frac{19x^3 - 66x^2 + 38x + 36}{19(6x^2 - 5)} \right|$$

By some tedious factorization,

$$\dots = |x - 2| \frac{|19x^3 - 28x - 18|}{19|6x^2 - 5|}$$

We estimate  $|x - 2|$  with constant 1

$$|x - 2| < 1$$

$$1 < x < 3 \text{ or } |x| < 3$$

It follows that

$$1 < x^2 < 9$$

$$1 < 6x^2 - 5 < 49$$

$$1 > \frac{1}{6x^2 - 5} > \frac{1}{49}$$

So

$$\frac{1}{19|6x^2 - 5|} < \frac{1}{19}$$

Similarly,

$$|19x^3 - 66x^2 + 38x + 36| \leq 19|x|^2 + 28|x| + 18 < 19 \cdot 3^2 + 28 \cdot 3 + 18 = 273$$

Thus

$$\dots \leq |x - 2| \cdot \frac{273}{19} < \varepsilon$$

$$|x - 2| < \frac{19}{273} \varepsilon$$

We conclude that it suffices to choose

$$\delta = \min \left\{ 1, \frac{19}{273} \varepsilon \right\}$$

■

**Theorem (Sequential Criterion of Limits).** Let  $f : A \rightarrow \mathbb{R}$  and let  $c$  be a cluster point of  $A$ .  $\lim_{x \rightarrow c} f = L$  if and only if for all sequence  $(x_n) \in A$  that converge to  $c$  and  $(x_n) \neq c, \forall n \in \mathbb{N}$ ,  $(f(x_n))$  converges to  $L$ .

*Proof.*  $(\Rightarrow)$ . By definition of convergent sequence,  $\forall \delta > 0, \exists K \in \mathbb{N}$  s.t.  $\forall n \in \mathbb{N}, n \geq K$ ,

$$|x_n - c| < \delta$$

By definition of limit,  $\forall \varepsilon > 0, \exists \delta > 0$  s.t.  $\forall x \in A$ ,

$$|x - c| < \delta \Rightarrow |f(x) - L| < \varepsilon$$

Thus, choose  $\delta$  given  $\varepsilon$ , and choose  $K$  given  $\delta$ , we have:

$$|x_n - c| < \delta \Rightarrow |f(x_n) - L| < \varepsilon$$

■

$(\Leftarrow)$ . We will prove this by contrapositive.

Assume that there exists  $\varepsilon_0 > 0$  and a  $(x_n) \in A$  converges to  $c$  with  $(x_n) \neq c$  such that for all  $n \in \mathbb{N}$

$$0 < |x_n - c| < \frac{1}{n} \Rightarrow |f(x_n) - L| \geq \varepsilon_0$$

Thus the function does not have a limit at  $c$ . We thereby conclude the converse of the statement.

■

**Theorem (Sequential Criterion of Divergence).** Let  $A \subseteq \mathbb{R}, f : A \rightarrow \mathbb{R}$ ,  $c$  be a cluster point of  $A$ ,  $(x_n) \rightarrow c$  s.t.  $(x_n) \neq c$ , and  $L \in \mathbb{R}^1$ .

1.  $L$  is NOT the limit of  $f$  at  $c \iff f(x_n)$  does NOT converge to  $L$ .
2.  $f$  diverges  $\iff f(x_n)$  does NOT converge.

*Proof.* Exercise.

■

**Example.**

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does NOT exist in } \mathbb{R}$$

*Proof.* Let  $(x_n) = \frac{1}{n} \rightarrow 0$ . Then

$$f(x_n) = \frac{1}{\frac{1}{n}} = n \rightarrow \infty$$


■

---

<sup>1</sup>The following theorems are NOT equivalent!!

**Example.**

$$\lim_{x \rightarrow 0} \sin\left(\frac{1}{x}\right) \text{ DNE in } \mathbb{R}$$



*Proof.* Let  $(x_n) = \frac{1}{2n\pi} \rightarrow 0$  and  $(y_n) = \frac{1}{\frac{1}{2}\pi + 2n\pi} \rightarrow 0$ . Then

$$f(x_n) = \sin(2n\pi) = 0$$

$$f(y_n) = \sin\left(\frac{1}{2}\pi + 2n\pi\right) = 1$$

Thus limit of  $f$  at 0 does NOT exist in  $\mathbb{R}$ . ■

**Example.**

**Definition (Signum Function).** Let  $A \subseteq \mathbb{R}$  and  $\text{sgn}(x) : A \rightarrow \mathbb{R}$  s.t.

$$\text{sgn}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0} \text{sgn}(x) \text{ DNE in } \mathbb{R}$$

*Proof.* Let  $(x_n) = \frac{1}{n} \rightarrow 0$  and  $(y_n) = \frac{1}{-n} \rightarrow 0$ . Then

$$f(x_n) = \text{sgn}(n) = 1$$

$$f(y_n) = \text{sgn}(-n) = -1$$

Thus limit of  $f$  at 0 does NOT exist in  $\mathbb{R}$ . ■

## 4.2 Limit Theorem

**Definition (Bounded Neighborhood of  $c$ ).** Let  $A \subseteq \mathbb{R}$ ,  $f : A \rightarrow \mathbb{R}$ ,  $c$  be a cluster point of  $A$ . Then we say  $f$  is a **bounded Neighborhood of  $c$**  if there exists  $\delta > 0$  and  $\exists M > 0$  s.t.  $\forall x \in A \cap V_\delta(c)$

$$|f(x)| \leq M$$

**Theorem (Existence of Bounded Neighborhood at Limit).** If  $f$  has a limit at  $c$ , then  $f$  is bounded on some neighborhood of  $c$ .

*Proof.* By definition of limit,  $\forall \epsilon > 0, \exists \delta > 0$  s.t.  $\forall x \in A \setminus \{c\}$

$$0 < |x - c| < \delta \Rightarrow |f(x) - L| < \epsilon$$

By triangle inequality,

$$|f(x)| - |L| \leq |f(x) - L| < \epsilon$$

$$|f(x)| < |L| + \epsilon$$

If  $f(c)$  is not defined on  $A$ , then let  $M = |L| + \epsilon$ . If  $f(c)$  is defined on  $A$ , then let  $M = \max\{f(c), |L| + \epsilon\}$ . Since the choice of  $\epsilon$  is arbitrary, we will choose  $\epsilon = 1$ . Thus,

$$f(x) \leq M$$

■

**Definition.** Let  $A \subseteq \mathbb{R}$ ;  $f, g : A \rightarrow \mathbb{R}$ ,  $c$  be a cluster point of  $A$ . Then  $\forall x \in A$

- **Sum** of function:  $(f + g)(x) = f(x) + g(x)$ .
- **Difference**:  $(f - g)(x) = f(x) - g(x)$ .
- **Multiple**:  $(bf)(x) = b \cdot f(x)$  for some  $b \in \mathbb{R}$ .
- **Product**:  $(f \cdot g)(x) = f(x) \cdot g(x)$ .
- **Quotient**:  $(\frac{f}{g})(x) = \frac{f(x)}{g(x)}$  if  $g(x) \neq 0$ .

**Theorem (Limit Theorem).** If  $\exists L, M \in \mathbb{R}$  s.t.  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$ , then

- $\lim_{x \rightarrow c} (f + g)(x) = L + M$ .
- $\lim_{x \rightarrow c} (f - g)(x) = L - M$ .
- $\lim_{x \rightarrow c} (bf)(x) = b \cdot L$ .
- $\lim_{x \rightarrow c} (f \cdot g)(x) = L \cdot M$ .
- $\lim_{x \rightarrow c} (\frac{f}{g})(x) = \frac{L}{M}$  if  $\lim_{x \rightarrow c} g(x) \neq 0$ .

*Proof. Exercise.*

■

**Remark.** Always check conditions before applying: this is true only if both  $f$  and  $g$  has a limit at  $c$  !



**Example.**  $\lim_{x \rightarrow 4} \frac{(x-4)(x+3)}{4(x-4)(x-5)} = \lim_{x \rightarrow 4} \frac{x+3}{4(x-5)} = -\frac{7}{4}$

**Corollary (Polynomial Function).**

$$\lim_{x \rightarrow c} p(x) = \lim_{x \rightarrow c} \sum_{i=0}^n a_i \cdot x^i = \sum_{i=0}^n a_i \cdot \lim_{x \rightarrow c} x^i = \sum_{i=0}^n a_i \cdot c^i = p(c)$$

**Corollary (Rational Function).** For polynomial functions  $p(x), q(x)$  s.t.  
 $\lim_{x \rightarrow c} p(x) \rightarrow p(c), \lim_{x \rightarrow c} q(x) \rightarrow q(c) \neq 0,$

$$\lim_{x \rightarrow c} \frac{p(x)}{q(x)} = \frac{p(c)}{q(c)}$$

**Theorem.** Let  $A \subseteq \mathbb{R}; f, g : A \rightarrow \mathbb{R}, c$  be a cluster point of  $A$ .

If  $\exists a, b \in \mathbb{R}, \forall x \in A, x \neq c$  satisfying

$$a \leq f(x) \leq b \text{ and } \lim_{x \rightarrow c} f = L$$

then

$$a \leq \lim_{x \rightarrow c} f \leq b$$

*Proof.*  $\forall (x_n) \in A \setminus \{c\}, (x_n) \rightarrow c,$

$$a \leq f(x_n) \leq b \Rightarrow a \leq L \leq b$$

■

**Theorem (Squeeze Theorem).** Let  $A \subseteq \mathbb{R}; f, g, h : A \rightarrow \mathbb{R}, c$  be a cluster point of  $A$ .

If  $\forall x \in A, x \neq c,$

$$f(x) \leq g(x) \leq h(x) \text{ and } \lim_{x \rightarrow c} f = L = \lim_{x \rightarrow c} h$$

then

$$\lim_{x \rightarrow c} g = L$$

*Proof. Exercise. Try squeeze theorem of sequence.*

■

**Example.**  $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) = 0$

*Proof.* Notice that

$$-1 \leq \sin\left(\frac{1}{x}\right) \leq 1 \Rightarrow -x \leq x \sin\left(\frac{1}{x}\right) \leq x$$

Since

$$\lim_{x \rightarrow 0} (-x) = 0 = \lim_{x \rightarrow 0} (x)$$

$$\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) = 0$$

■

**Example.** Let  $b \in \mathbb{R}, b > 0$ . Then  $\lim_{x \rightarrow 0} x^b = 0$ . Notice that when  $x \in [0, 1]$

$$x^{\lceil b \rceil} \leq x^b < x^{\lfloor b \rfloor}$$

The rest is left as exercise.

**Theorem.** Let  $A \subseteq \mathbb{R}$ ;  $f, g, h : A \rightarrow \mathbb{R}$ ,  $c$  be a cluster point of  $A$ . If  $\lim_{x \rightarrow c} f > 0$ , then  $\exists \delta > 0$  s.t.  $V_\delta(c)$  s.t.  $\forall x \in A \cap V_\delta(c) \setminus \{c\}$ ,

$$f(x) > 0$$

*Proof. Exercise.*

