

Universidade Católica Portuguesa

Escola Superior de Biotecnologia

MSc Biomedical Engineering

**AI-Based Characterization of Clinical Alzheimer's
Disease Stages from Structural Brain MRI Using
Foundation Model Representations**

Biomedical Project

Author

Frederik Dürr // Lucca Moreira // Laurindo

Supervisor

Prof. Dr. Pedro Rodrigues

Co-supervisor

Dr. Ines Barbosa

Porto, Portugal

2027

Abstract

[Replace this placeholder: Write a concise abstract here. A strong scientific abstract usually contains the background/problem, objective, methods, key results, and conclusion.]

Keywords: *[Replace this placeholder: keyword 1; keyword 2; keyword 3; keyword 4; keyword 5]*

Contents

Abstract	i
List of Figures	iv
List of Tables	v
List of Abbreviations	vi
1. Introduction	1
1.1. Clinical Context of Alzheimers disease	1
1.1.1. Diagnosing Alzheimers Disease	2
1.2. Foundation Models	2
1.3. Structural MRI acquisition	2
1.4. MRI properties for analysis	3
1.4.1. MRI in diagnosing Alzheimer’s disease	3
1.5. Foundation Models	4
1.6. Aim and Research Questions	4
2. Materials and Methods	5
2.1. Dataset	5
2.3. Image Preprocessing	5
2.4. Feature Extraction	6
2.5. Machine Learning / Deep Learning	6
2.6. Statistical Analysis	6
2.7. Software and Reproducibility	6
3. Results	7
3.1. Cohort / Dataset Characteristics	7
3.2. Feature Results	7
3.3. Model Performance	7
4. Discussion	8
4.1. Main Findings	8
4.2. Comparison with Previous Literature	8
4.3. Limitations	8

4.4. Implications and Future Work	8
5. Conclusion	9
References	10
A. Additional Material	11

List of Figures

List of Tables

3.1. Example model-performance table. Replace all values before submission. .	7
---	---

List of Abbreviations

Magnetic resonance Imaging (MRI)

Computed Tomography (CT)

Alzheimer's disease (AD)

1. Introduction

1.1. Clinical Context of Alzheimers disease

Alzheimer's disease (AD) is a progressive neurodegenerative disorder and the most common cause of dementia worldwide. Clinically, AD is characterized by a progressive decline in memory, cognitive abilities, behaviour, and the ability to perform everyday activities. Dementia represents a major global health challenge: in 2021, approximately 57 million people were living with dementia worldwide, with nearly 10 million new cases occurring each year. Alzheimer's disease accounts for an estimated 60–70

A major challenge in the management of AD is the long period between the initiation of pathological processes and the appearance of clinically recognizable symptoms. The pathological changes associated with AD can begin approximately 15–20 years before the onset of obvious cognitive symptoms [1]. During this preclinical phase, pathological alterations can accumulate while the affected individual may remain clinically asymptomatic. Consequently, substantial neuronal and structural changes may already be present when cognitive impairment becomes apparent. This long latent phase highlights the importance of developing sensitive methods for the early detection and characterization of AD, as early intervention may provide a greater opportunity to slow disease progression [1].

In addition to the challenge of early detection, the development of effective therapies for AD remains difficult. One of the major obstacles to pharmacological treatment is the blood–brain barrier (BBB), a highly selective physiological barrier that protects the central nervous system from potentially harmful substances in the systemic circulation. While the BBB is essential for maintaining brain homeostasis, it also restricts the transport of many therapeutic compounds into the brain, thereby limiting their bioavailability at the intended site of action [2]. This is particularly relevant for therapeutic approaches involving large biological molecules and other compounds with limited BBB permeability.

Consequently, improving the delivery of therapeutic agents to the central nervous system has become an important research objective in AD treatment. Several strategies have been investigated to overcome or circumvent the BBB, including receptor-mediated transport, focused ultrasound, intranasal administration, liposomal systems, and nanoparticle-based drug delivery systems [2, 3]. Nanoparticle-based approaches are particularly promising because their physicochemical properties can be engineered to improve drug solubility and stability, control drug release, enhance transport across the BBB, and potentially increase accumulation

at specific target sites while reducing systemic exposure [4, 5].

The investigation of novel drug delivery systems is therefore important for enabling therapeutic agents to reach the brain at concentrations sufficient to exert their intended effects. This is particularly relevant as increasingly sophisticated disease-modifying therapies are being developed, for which efficient and targeted delivery to the central nervous system remains a significant challenge. However, despite promising preclinical results, nanoparticle-based and other advanced delivery systems still face important translational challenges, including long-term safety, biological variability, manufacturing scalability, and regulatory requirements [5]. Addressing these limitations is therefore essential for translating advanced drug delivery concepts into clinically effective therapies for Alzheimer's disease.

Taken together, the combination of a long preclinical disease phase, the progressive and irreversible nature of neurodegeneration, and the restricted accessibility of the brain to many therapeutic compounds demonstrates the need for both improved early disease characterization and more effective drug delivery strategies. Investigating advanced drug delivery systems capable of overcoming the limitations imposed by the BBB may therefore contribute to the development of more targeted and effective therapeutic approaches for AD.

Labels of interest in the context of Alzheimer's Disease.

1.1.1. Diagnosing Alzheimers Disease

Atrophy due to gray matter loss with anatomical magnetic resonance imaging or hypometabolism with 18F-fluorodeoxyglucose positron emission tomography (FDG PET).

This diagnostic status is established by clinicians based on cognitive scores such as the Mini-Mental State Evaluation (MMSE), and the Clinical Dementia Rating. Even though assessment protocols are standardized, these scores can be highly variable based on the examiner and the physical/psychological condition of the patient during the test.

1.2. Foundation Models

1.3. Structural MRI acquisition

Image Acquisition: A strong magnetic field is applied B_0 , normally 1.5-3T. That forces the hydrogen nuclei of the abundant water molecules in the soft tissue in the body to align with the magnetic field. A gradient coil applies a second magnetic field, which leads to a different precession frequency of molecules in the field of interest. A radio-frequency (RF) impulse is applied with the same frequency as the gradient coil made the molecules spin. This pulse which tilts these nuclei from their alignment along the B_0 . The nuclei then precess back to the alignment. The precessing nuclei emit a signal, which is registered by the receiver coils in the scanner. Each signal that is produced by the realignment of the hydrogen nuclei to

the magnetic field B_0 has two components. First the longitudinal z-axis, which is along the scanner's magnetic field. The second is the transversal signal. This describes the xy-plane orthogonal to the scanner's magnetic field.

Initially the longitudinal signal is weak, due to the tilt of the nuclei. But over time it increases again due to the realignment of the hydrogen nuclei. The time constant that dictates the speed of realignment is denoted by T_1 . The transversal signal on the other hand is strong due to phase coherence. The signal decays as the nuclei dephase and realign. This decay is denoted as the T_2 time constant.

Tissues have different T_1 and T_2 relaxation times, which are used to build contrast in the MRI image. In the final image the contrast depends on the receiving time of a signal, the echo time (TE) and how fast you repeat the tilt-relax-process, repetition-time (TR).

1.4. MRI properties for analysis

Magnetic resonance Imaging (MRI) is a widely used imaging modality in radiology. Even though MR has a long data acquisition time, and complex scanning protocols, it is preferred more than other imaging modalities such as Computed Tomography (CT) because it lacks the use of ionizing radiation and has a better visualization potential.

1.4.1. MRI in diagnosing Alzheimer's disease

Alzheimer's is a complex little researched neurodegenerative disease, which can not be diagnosed within a single test. However, brain imaging plays an important role in diagnosing Alzheimer's disease. The 3D image of the brain provides a great opportunity for physicians to spot abnormalities. These abnormalities can be markers of Alzheimer's disease, but could also be symptoms of other cognitive conditions [6]. Alzheimer's is primarily connected to atrophy in key regions like the hippocampus, the core memory area of the brain, and parietal lobe. *[Replace this placeholder:*

1. *K-space: It is a collection of spatial frequencies captured after applying the pulse sequences. The (x,y) coordinates in this space represent the frequency and phase for every pixel of the image generated after applying inverse Fourier transform to the captured frequency data.*
2. *Fourier transform: is a mathematical technique that allows a signal to be decomposed into a sum of sine or cosine waves of different frequencies, phases and amplitudes. This link can be followed for further reading about Fourier transform*

]

1.5. Foundation Models

Foundation models can be used as a Feature extractor the workflow would be MRI-Images-> Foundation Models -> embedding matrix (Feature representation) -> Classification model -> 6 classes **Three ways of foundation model approaches**

1. Frozen Foundation model. You don't change the weights of the model itself, instead you fine-tune the classifier.
2. Fine tune the foundation model. Fine tune is much more computationally expensive, and can be prone to overfitting in smaller data sets
3. Zero-shot classification. No classification needed.

1.6. Aim and Research Questions

The aim of this project is to analyse whether pretrained foundation models can turn structural MRI slices into useful numerical representations, then test whether those representations separate six clinical groups of alzheimer's disease.

[Replace this placeholder: State the main aim and, where appropriate, explicit research questions or hypotheses.]

2. Materials and Methods

2.1. Dataset

[Replace this placeholder: Describe the dataset, inclusion/exclusion criteria, sample size, acquisition protocol, preprocessing, and relevant ethics/data-governance aspects.]

2.2.

2.3. Image Preprocessing

[Replace this placeholder: This part we can reference to the paper they have published in the beginning of this year] Image processing is a highly relevant procedure for deploying statistical pipelines. Without a solid and robust preprocessing the downstream task will not produce any relevant information. For this project we gathered pre-processed sMRI images. The fully description of the pre-processing steps can be investigated in J. Menzes et al. [7]. Here we are going to explain the pre-processing in a short version.

One of the methods of preprocessing in MRI images is the bias field correction. The bias fields is a low-frequency spatially varying MRI artifact resulting from spatial inhomogeneity of the magnetic field, variations in the sensitivity of the reception coil, and the interaction between the magnetic field and the human body. The bias fields is dependent on the strength of the magnetic field. By increasing the magnetic field, which increases the spatial resolution also increases the bias field, resulting in a higher Signal to noise ratio.

Followed by the Image registration. This step is fundamental for further analysis. It aligns all images, in space positioning and inter-subject anatomically variability.

another important pre-processing method is the brain extraction, skull-stripping. Image contrast from non-brain tissue such as fat, skull, or neck can cause issues with downstream analysis starting with the brain tissue segmentation. The brain extraction generates a mask that identifies brain voxels comprising gray-matter, white-matter, and Cerebrospinal fluid (CSF) of the cerebral cortex and subcortical structures, including the brain stem and the cerebellum.

Temporal Signal-to-Noise Ratio: Signal to noise ratio is evaluated by the mean signal intensity in each voxel divided by the standard deviation of the signal across time points. **Spatial Normalisation:**

Image Registration: A process that aligns an image from one coordinate space to another.

E.g. images taken on different days or locations. Differences can appear due to different spatial resolutions and field of view. **Segmentation and Parcellation:** Brain tissues are divided into:

1. gray matter (GM), containing neuronal cell bodies
2. white matter (WM), including neuron connection fibres wrapped in a special signal-acceleration substance called myelin.
3. Cerebrospinal fluid (CSF), a protecting fluid mostly around the brain but also within the brain cavities called ventricles.
- 4.

[Replace this placeholder: Describe registration, skull stripping, normalization, segmentation, resampling, denoising, bias-field correction, or other preprocessing steps as applicable.]

2.4. Feature Extraction

[Replace this placeholder: Describe the features, extraction method, software, and parameters.]

2.5. Machine Learning / Deep Learning

[Replace this placeholder: Describe model architecture, train/validation/test split, preprocessing, class balancing, hyperparameter selection, training procedure, and reproducibility.]

2.6. Statistical Analysis

[Replace this placeholder: Describe statistical tests, assumptions, confidence intervals, effect sizes, and significance thresholds.]

2.7. Software and Reproducibility

The analysis was implemented in Python. Record the relevant package versions and the operating environment here.

3. Results

3.1. Cohort / Dataset Characteristics

[Replace this placeholder: Present descriptive statistics and dataset characteristics.]

3.2. Feature Results

[Replace this placeholder: Present extracted features and important observations.]

3.3. Model Performance

[Replace this placeholder: Report performance metrics with uncertainty and validation details.]

Table 3.1.: Example model-performance table. Replace all values before submission.

Model	Accuracy	Precision	Recall
Baseline	0.00	0.00	0.00
Model A	0.00	0.00	0.00
Model B	0.00	0.00	0.00

As shown in table 3.1, this is only a formatting example.

4. Discussion

4.1. Main Findings

[Replace this placeholder: Interpret the principal findings without repeating the Results chapter.]

4.2. Comparison with Previous Literature

[Replace this placeholder: Compare your findings with the relevant literature and explain similarities, differences, and plausible methodological reasons.]

4.3. Limitations

[Replace this placeholder: Discuss limitations in data, methodology, generalisability, statistical power, and potential bias.]

4.4. Implications and Future Work

[Replace this placeholder: Describe what the findings imply and what should be investigated next.]

5. Conclusion

[Replace this placeholder: State the main conclusion directly and concisely. Answer the research question without introducing new results or references.]

References

- [1] M. S. Rafii and P. S. Aisen. “Detection and treatment of Alzheimer’s disease in its preclinical stage”. In: *Nature Aging* 3 (2023), pp. 520–531. doi: 10.1038/s43587-023-00410-4.
- [2] I. Iqbal, F. Saqib, Z. Mubarak, M. F. Latif, M. Wahid, B. Nasir, H. Shahzad, J. Sharifi-Rad, and M. S. Mubarak. “Alzheimer’s disease and drug delivery across the blood–brain barrier: approaches and challenges”. In: *European Journal of Medical Research* 29.313 (2024). DOI: 10.1186/s40001-024-01915-3.
- [3] “Drug delivery strategies to cross the blood-brain barrier in Alzheimer’s disease: a comprehensive review on three promising strategies”. In: *Frontiers in Neuroscience* (2025).
- [4] A. K. Roghani, R. I. Garcia, A. Roghani, A. Reddy, S. Khemka, R. P. Reddy, V. Pattoor, M. Jacob, P. H. Reddy, and U. Sehar. “Treating Alzheimer’s disease using nanoparticle-mediated drug delivery strategies/systems”. In: *Ageing Research Reviews* (2024). DOI: 10.1016/j.arr.2024.102291.
- [5] K. Roamcharern and K. Yubolphan. “Overcoming the blood–brain barrier in Alzheimer’s disease: translational perspectives on advanced drug delivery platforms”. In: *Frontiers in Neuroscience* 20 (2026). DOI: 10.3389/fnins.2026.1810486.
- [6] Envision Radiology. *Role of MRIs in Detecting Alzheimer’s*. URL: <https://www.envrad.com/role-of-mris-in-detecting-alzheimers/> (visited on 09/25/2026).
- [7] J. Menezes, M. I. Barbosa, and P. M. Rodrigues. “ β -amyloid–stratified six-stage Alzheimer’s disease discrimination via multiband MRI histogram and GLCM features”. In: *Biomedical Signal Processing and Control* 127 (2026), p. 111143. DOI: <https://doi.org/10.1016/j.bspc.2026.111143>. URL: <https://www.sciencedirect.com/science/article/pii/S1746809426016976>.

A. Additional Material

[Replace this placeholder: Use appendices for detailed material that is useful for transparency but would interrupt the main narrative, such as extra tables, questionnaires, parameter lists, or code.]