

1 (In)determinacy in woody plants: limits and opportunities for
2 timing growth in a changing climate

3 Frederik Baumgarten¹⁻³, Sally Aitken¹, Yann Vitasse³, Robert D Guy¹, EM Wolkovich¹

4 May 2, 2026

5 ¹ Department of Forest and Conservation Sciences, Faculty of Forestry, University of British Columbia,
6 2424 Main Mall Vancouver, BC Canada V6T 1Z4.

7 ² Department of Environmental Sciences-Botany, University of Basel, Schönbeinstrasse 6, Basel 4056,
8 Switzerland.

9 ³ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse. 111, Bir-
10 mensdorf 8903, Switzerland

11 **Corresponding Author:** Frederik Baumgarten; frederik.baumgarten@unibas.ch;
12 Tel.: +41 61 207 23 37

13 **E-Mail-Addresses:** sally.aitken@ubc.ca; yann.vitasse@wsl.ch; rob.guy@ubc.ca; e.wolkovich@ubc.ca

14 **Type of article:** Perspective

15 **Short running title:** (In)determinate growth strategies of trees

16 **Keywords:** plant growth, tree phenology, shoot extension, indeterminate growers, carbon sequestra-
17 tion, growing season length, drought, genetic programming, phenotypic plasticity, forest productivity

18 **wordnumbers:** abstract: 200; main text: 4005; box1: 404; number of figures: 4; number of tables: 0

19 **number of references:** 95

20 **Authors contribution:** FB and EM initiated this project. FB took a lead in drafting, designed
21 figures and performed phylogenetic analyses. All authors contributed substantially to the writing of
22 the article.

23 **Data accessibility statement:** The data supporting the phylogeny are archived in the following
24 public repository. Link: <https://doi.org/10.5281/zenodo.18637216>.

25 **Conflict of interest statement:** The authors declare no conflict of interest

Abstract

When and how much organisms grow given environmental constraints are fundamental questions in biology, and especially pressing given the need to predict biomass production and carbon sequestration with climate change. While environmental factors such as temperature and water availability directly regulate plant growth, the timing and rate of growth are also governed by a plant's phenological sequence—its genetically programmed developmental stages. This sequence is critical for predicting climate responses but often underappreciated.

Here, we leverage the concept of (in-)determinacy in growth and development—which captures how flexibly plants produce organs from preformed cells versus initiating them *de novo*—to propose a new framework for predicting tree growth responses to climate change.

We propose that: 1) determinate growth is an adaptation to strongly seasonal climates, restricting tissue investment to a short favorable period before dormancy; 2) indeterminate species are more exposed to increasing climatic extremes (spring/fall frost, drought) but may recover more effectively from resulting damage; and 3) species with greater indeterminacy are more likely to benefit from lengthening frost-free growing seasons. Future carbon sequestration may thus depend not only on environmental changes, but also on the degree of determinacy set by intrinsic genetic programming.

Introduction

Investing the right amount of resources in growth at the right time is of critical importance to the survival and fitness of any living organism. The topic of growth strategies and habits has a long history in science, spanning the fields of genetics, genomics, physiology and ecology across the animal and plant kingdoms (Stearns, 1998). At its core lies the concept of determinacy—the classification of organisms as either reaching a fixed size at maturity or continuing to grow throughout their lives. Like mollusks, fish and reptiles, plants add biomass to their primary bodies as long as they live and are therefore considered ‘indeterminate growers’ (Ejzmond *et al.*, 2010).

Various terms have emerged to describe this phenomenon across spatial and temporal scales, e.g. from a single cell to the whole organism, and from a single season to an entire lifetime (McDaniel, 1992; Karkach, 2006). In plants, organs can develop from preformed embryonic tissue (e.g. primordia in overwintering buds), form *de novo*, or both, depending on species and life form. While these underlying developmental and meristematic constraints apply broadly across the plant kingdom, they play a particularly central role in the life cycle of woody perennials, especially trees growing in extra-tropical regions, which must balance growth, survival, and reproduction over long lifespans in highly seasonal environments. As we use it here, (in)determinacy in woody perennials refers to the ability to:

- a) preform organs in one year as a future investment, which is deployed rapidly in the following spring with no further primary shoot growth during that season (determinate strategy);
- b) maintain a continuous or episodic meristem activity by forming new shoot tissue during the current growing season (indeterminate strategy).

In this Perspective, we revisit classical concepts of determinacy as they relate to shoot growth in woody perennials, place them in the context of climate change, and explore to what extent related developmental principles may also inform growth processes in other meristematic tissues. Although we focus on woody perennials, the concept of (in)determinacy is broadly relevant across vascular plants, including herbaceous species; trees provide a particularly informative model system because the ecological and climatic consequences of growth timing are especially pronounced in long-lived woody plants. To provide context, we first outline the fundamental limitations to plant growth in extra-tropical regions by linking external environmental drivers with internal growth control mechanisms. In this context,

our aim is to provide a conceptual and methodological basis for analysing (in)determinacy as a trait relevant to plant responses to climatic variability and change, to clarify how the concept can be defined and measured, and to identify key research gaps and directions for future comparative and trait-based analyses.

The further one travels from the equator towards the poles, the more strongly plants are confined to a shrinking time window of opportunity for growth set primarily by low temperatures. Below c. 5 °C, metabolic activity slows to an extent where growth and development largely cease (Schenker *et al.*, 2014; Rossi *et al.*, 2008; Körner, 2008). More importantly, sub-freezing temperatures can cause severe damage to plant tissue if they occur at vulnerable developmental stages, e.g. during or after leaf unfolding, prior to fruit maturation, or before cold acclimation in fall (Sakai & Larcher, 1987; Baumgarten *et al.*, 2023). While annual plant species accommodate their entire life cycle within this window, perennial (woody) species of extra-tropical regions are forced to partition their growing phase seasonally, with periods of activity alternating with periods of dormancy (Delpierre *et al.*, 2016). This is referred to as intermittent or rhythmic (as opposed to continuous) growth.

High temperatures can directly inhibit growth by exceeding species-specific physiological thresholds (O’Sullivan *et al.*, 2017). In addition, increased evaporative demand and reduced precipitation can lower soil moisture, leading to water stress that slows or stalls growth and, in extreme cases, causes leaf scorching or premature senescence (Hsiao, 1973; Pugnaire *et al.*, 1999; Etzold *et al.*, 2021; Estiarte & Peñuelas, 2015). Together, these temperature and soil moisture limitations act as environmental filters, narrowing the period during which growth is possible (Figure 1). Notably, comparable (or even stronger) seasonal constraints occur in seasonally dry forests as is typical in Mediterranean climates, where temperatures may remain suitable for growth year-round but multi-month dry seasons impose pronounced water limitation. Across species, leaf flush and renewed growth commonly track the onset of rains and access to plant-available water (including deep soil moisture), making soil water availability a primary driver of seasonal activity in these systems (Borchert, 1994; Ramos *et al.*, 2023; Alvarado *et al.*, 2025).

Light also influences plant growth through two key processes. First, light intensity provides the primary energy source that drives photosynthesis and, consequently, source activity. Second, light carries a

105 wealth of information through its quality and the length of daily light and dark periods (photoperiod),
 106 which mediate many physiological processes, including bud set and dormancy transitions (Wang *et al.*,
 107 2024). Both factors may contribute to the common observation that tree growth often peaks near the
 108 summer solstice in extra-tropical regions (Rossi *et al.*, 2006; Etzold *et al.*, 2021; Luo *et al.*, 2018) or that
 109 a window of temperature sensitivity ‘opens’ around the summer solstice to synchronize reproductive
 110 efforts over very large spatial scales (Journé *et al.*, 2024).

111 **Internal programming of plants**

112 Given our relatively robust understanding of environmental influences on plant growth, it may seem
 113 straightforward to predict how plants will grow under current and future climatic conditions. Yet,
 114 this has proven challenging, particularly when predicting responses under scenarios with extended, cli-
 115 matically favorable growing seasons (Zohner *et al.*, 2021). Here we propose that accurate predictions
 116 require incorporating another key, but often overlooked, factor: internal growth control—the geneti-
 117 cally encoded developmental program implemented through hormonal and gene-regulatory networks,
 118 that may prevent further growth *despite* favorable environmental conditions.

119
 120 While plants have evolved numerous morphological adaptations to tolerate or avoid harmful environ-
 121 mental conditions, most temperate and boreal woody perennials cope with seasonal fluctuations in
 122 temperature and moisture by temporally escaping these conditions. They do this by aligning their
 123 life-history events (phenology) with a cycle of growth and dormancy that balances survival, reproduc-
 124 tion, and growth over the long lifespan typical of many tree species (Delpierre *et al.*, 2016). Hence,
 125 the phenological sequence can impose abrupt internal switches in resource allocation—from vegetative
 126 growth to reproduction (flowering, fruit maturation) and storage (Stearns, 1989; Chapin *et al.*, 1990).
 127 These transitions act as additional internal filters, further narrowing the effective window during which
 128 vegetative growth can occur, regardless of environmental potential (Figure 1).

130 **The concept of (in)determinate growth**

131 In annual plants, growth ends with the production of flowers to form fruits and seeds: a signal in the
 132 apical meristem triggers a sudden switch in resource investment from vegetative growth to building
 133 a reproductive structure with no point of return, signaling the end of the life-cycle (Poethig, 2003;

134 Huijser & Schmid, 2011). A shoot axis that terminates in a flower or other modified structure and
135 ceases meristematic activity is considered ‘determinate’ (Barthélémy & Caraglio, 2007). In woody
136 perennial species, however, flowers are typically produced on lateral shoots or buds, preserving the
137 main vegetative axis and allowing continued structural expansion. However, ‘determinate growth’ also
138 refers to a temporal suspension of the meristem, which leads to a time lag between the formation of
139 organs and their expansion (Kozlowski, 2012; Hallé *et al.*, 1978).

140
141 In their first year, all seedlings exhibit indeterminate growth. They begin growth from an embry-
142 onic shoot meristem and typically prioritize rapid leaf and canopy establishment, reflecting a juvenile,
143 largely opportunistic phase of development (Borchert, 1976). In subsequent years, many temperate
144 and boreal tree species preform their new shoot increments during the previous growing season, over-
145 wintering them as primordial shoots with leaves and internodes in hardened buds to be ‘ready-to-go’
146 when spring arrives. Once the leaves flush and the preformed internodes elongate, species differ in
147 how long the shoot apical meristem remains competent to continue organ production (Figure 2). Some
148 species exhibit strongly determinate shoot extension, in which the terminal meristem is intrinsically
149 downregulated soon after the spring flush and remains inactive for the rest of the season, consistent
150 with a prolonged internal suppression that is hormonally mediated (paradormancy; Lang *et al.*, 1987;
151 Figure 4; e.g. *Fraxinus* spp., *Acer* spp. and *Prunus* spp.). At the other extreme, highly indetermi-
152 nate species maintain an active shoot apical meristem and can continue producing new leaves and
153 internodes on top of the preformed shoot as long as conditions remain favorable, sometimes stretching
154 their indeterminate growth well into autumn until low temperature eventually forces them to stop
155 (e.g. *Sequoia* spp., *Cupressus* spp. and *Alnus* spp.). Between these extremes, many species occupy an
156 intermediate position: they continue to produce neoformed leaves and internodes after the preformed
157 increment, but growth cessation is often coordinated by seasonal cues—frequently photoperiod acting
158 through hormonal regulation (e.g. *Pinus* spp., *Betula* spp. and *Populus* spp.; Maurya & Bhalerao,
159 2017; Triozzi *et al.*, 2018; Böhlenius *et al.*, 2006). Indeterminate growth can occur either through
160 continuous activity of the shoot apical meristem forming new leaves and internodes without buds (free
161 growth), or through the formation and subsequent flushing of buds mid-season without a period of
162 dormancy (second flushing or lammas growth; Figure 2).

163
164 Although this concept is often presented dichotomously (Kozlowski & Pallardy, 1997; Lechowicz, 1984),

but see Kikuzawa, 1983; Damascos *et al.*, 2005), with species classified as either determinate or indeterminate growers, most species likely exist along a gradient with numerous intermediate forms. For instance, many temperate oaks (*Quercus spp.*) are considered determinate growers, but can exhibit multiple flushes. Similarly, Douglas-fir (*Pseudotsuga menziesii*) gradually adopts a more determinate growth habit with maturity (Borchert, 1976; Heuret *et al.*, 2006). Importantly, the degree of determinacy can be quantified at the species level, at least for the shoot apical meristem as the ratio of leaves produced during the relevant growth period to the number of leaf primordia preformed in the bud, with values above 1 indicating increasing neoformation (Box 1). Figure 4 provides a literature-based classification of determinacy for some common species, many of which fall into an intermediate position.

Evolutionary trade-offs: the cost of certainty and flexibility

In most temperate and boreal ecosystems, determinate and indeterminate growth strategies co-occur, indicating that neither strategy is universally superior. Instead, they reflect contrasting solutions to a fundamental trade-off between growth certainty and growth flexibility under seasonal and variable environments.

Indeterminate species are characterized by continued or episodic activity of the shoot apical meristem during the growing season. This does not necessarily imply faster intrinsic growth rates, but rather a greater capacity to extend growth in time when conditions remain favorable. Such temporal flexibility allows individuals to exploit transient resource availability and recover from partial damage, traits that are advantageous in disturbed or early-successional environments. Accordingly, indeterminate growth is frequently associated with early successional species (Marks, 1975). However, maintaining active meristems later into the season also increases exposure to environmental stressors such as summer drought and early autumn frost, thereby raising the risk of tissue loss or incomplete maturation (Figure 4). As a result, indeterminate growth strategies often involve higher interannual variability in performance and, in some cases, shorter tissue or organismal lifespans (Millet *et al.*, 1999; Brien *et al.*, 2020).

In contrast, determinate species preform a substantial portion of their shoot growth in advance and

195 deploy this tissue during a restricted period of the growing season. This strategy limits short-term
 196 plasticity, because the year’s shoot-growth potential is set during bud formation in the preceding year
 197 and is largely realized by expanding preformed organs (Gordon *et al.*, 2006). At the same time, it
 198 reduces exposure of sensitive tissues to late-season stress and synchronizes growth with the most pre-
 199 dictable and favorable part of the season. As a consequence, determinate growth strategies are often
 200 associated with stronger local adaptation and reduced sensitivity to short-term climatic fluctuations
 201 (Leites & Benito Garzón, 2023).

202
 203 These contrasting strategies help explain the coexistence of determinate and indeterminate species
 204 within the same landscapes. For example, western larch (*Larix occidentalis*) and Douglas-fir (*Pseudot-*
 205 *suga menziesii*) frequently co-occur despite marked differences in growth determinacy. Western larch
 206 exhibits more flexible, indeterminate shoot growth and performs well in environments shaped by fre-
 207 quent disturbance, whereas Douglas-fir shows more determinate growth and stronger local adaptation,
 208 conferring stability under predictable seasonal regimes (Roskilly & Aitken, 2024). Such differences
 209 suggest niche differentiation along gradients of disturbance frequency, environmental stress, and cli-
 210 matic predictability, rather than outcomes driven by straightforward competitive dominance.

211
 212 Together, these patterns indicate that growth determinacy reflects a trade-off between flexibility and
 213 reliability in seasonal environments. Under ongoing climate change—marked by longer growing seasons,
 214 greater climatic variability, and increasing environmental stress in many regions—it remains uncertain
 215 whether determinate or indeterminate growth strategies will be favored. Resolving both how trees
 216 regulate their degree of (in)determinacy and which levels along this trait continuum are advantageous
 217 under future conditions will be critical for predicting shifts in species performance, coexistence, and
 218 forest composition.

219 **The performance of (in)determinacy with climate change**

220 Spring warming has advanced the onset of leaf emergence by up to a month compared to pre-industrial
 221 times (Vitasse *et al.*, 2022), reflecting an earlier reactivation of shoot apical and cambial meristems
 222 under warmer conditions. In contrast, the cessation of growth in autumn has shifted far less than
 223 expected given the extension of climatically favorable conditions (Zani *et al.*, 2020; Zohner *et al.*,
 224 2023). As a result, phenological sequences are increasingly observed to shift as a whole toward spring

rather than being symmetrically extended at both ends of the growing season (Keenan & Richardson, 2015; Meng *et al.*, 2024). This asymmetric response highlights a key constraint relevant to growth (in)determinacy: while growth onset is highly plastic and environmentally responsive, the duration of meristematic activity later in the season appears more tightly regulated by plant internal processes and differs markedly among species. Such constraints on growth cessation help explain why longer growing seasons do not necessarily translate into increased growth or biomass production (Zani *et al.*, 2020; Bose *et al.*, 2025; Wolkovich *et al.*, 2025).

Along the continuum of growth determinacy, we hypothesize that species positioned toward the more determinate end will largely maintain their fixed seasonal growth schedules under climate warming, with limited capacity to extend growth later into the season. In regions where climate change primarily increases heat and water stress during mid to late summer, such species are expected to show reduced annual biomass production, because their restricted growth window increasingly overlaps with unfavorable conditions (Figure 3).

In contrast, we predict that species toward the indeterminate end of the continuum will be more capable of prolonging meristematic activity into late summer and autumn when conditions become favorable again, potentially resulting in higher annual production under some future climate scenarios (Caffarra & Donnelly, 2011; Richardson *et al.*, 2009; Zohner *et al.*, 2020, Figure 3). This capacity for temporal extension of growth is particularly common among early successional species, which tend to leaf out earlier in spring and senesce later in autumn than late-successional species. Their earlier leaf-out is consistent with generally lower chilling and forcing requirements for bud burst and dormancy release (Heide, 1993; Laube *et al.*, 2014; Hu *et al.*, 2022; Baumgarten *et al.*, 2021; Basler & Körner, 2012), and in many cases weaker photoperiod sensitivity (Basler & Körner, 2012).

Extreme climatic events and increasing environmental stress provide an important context for evaluating these predictions. More determinate growth strategies concentrate growth earlier in the season and may therefore escape unfavorable growing conditions by operating within a relatively narrow spring to early-summer window that often coincides with comparatively favorable temperature and soil moisture conditions. Under historical climatic regimes, this strategy provided ample safety margins against late-season stress. Where climate change introduces novel or intensified drought stress within this

256 window, however, determinate species are likely to experience reduced annual production and limited
257 capacity for recovery within the same growing season once canopy development has been compromised
258 (Silvestro *et al.*, 2023, Figure 3).

259

260 Species with more indeterminate growth habits often remain green and physiologically active for longer
261 periods, leafing out early in spring and senescing late in autumn, with deciduous species occasionally
262 retaining functional foliage until the first freezing events (Figure 3). As a consequence, a substantial
263 fraction of their growth period coincides with parts of the growing season during which drought stress
264 is projected to intensify under future climates, particularly in water-limited regions (Figure 3). While
265 this extended activity increases exposure to environmental extremes, it also creates opportunities for
266 compensatory growth when conditions become favorable again.

267 In essence, the flexible growth schedule of indeterminately growing species may allow plants to: (1)
268 produce tissue better adapted to harsh environmental conditions as it is formed in the current season
269 (e.g., smaller leaves with higher specific leaf area), and (2) catch up and compensate later in the season
270 by another production boost (Stevens & Lindroth, 2005). Mediterranean ecosystems provide a clear
271 example of this flexibility, where many species suspend growth during summer drought but resume
272 meristematic activity in autumn or winter when conditions allow (Volaire *et al.*, 2023). Evidence that
273 trees can take advantage of such a ‘second growing season’ after summer drought has been found in
274 pines in Mediterranean climates through polycyclic flushing—a form of indeterminate growth (Girard
275 *et al.*, 2011, Figure 3,).

276

277 We therefore hypothesize that climate change will increasingly disrupt established phenological cy-
278 cles, favoring species and genotypes that are more plastic in rearranging their activities by resuming
279 growth and/or restoring reserves (e.g., starch, nutrients) later in the year, thereby recovering from
280 and compensating for stress-induced damage and losses. Such differential capacity to exploit newly
281 available growth opportunities or to recover after stress is expected to intensify competition among
282 co-occurring species by altering the timing and magnitude of resource uptake, potentially leading to
283 the reassembly of forest communities toward species occupying more indeterminate positions along the
284 growth continuum. This shift may be reinforced if climate change increases not only mean stress but
285 also climatic variability (van der Wiel & Bintanja, 2021), because flexible (plastic) growth schedules
286 can capitalize on transient favorable windows and compensate after setbacks (Richter *et al.*, 2012;

Chevin *et al.*, 2010; Botero *et al.*, 2015).

(In)determinacy beyond primary shoot growth

Although (in)determinacy is most clearly defined *sensu stricto* at the shoot apical meristem, its relevance may extend beyond primary shoot growth because shoot extension and cambial growth are structurally and functionally coordinated within the plant. Primary growth governs crown expansion and annual leaf deployment, thereby influencing total leaf area and canopy photosynthetic capacity (Girard *et al.*, 2011; Soolanayakanahally *et al.*, 2015). Primary shoot growth also establishes the structural framework within which later radial growth can occur by determining the extent and geometry of axes along which cambial activity can be expressed. Empirical studies showing close within-season associations between shoot extension and stem or radial growth therefore point less to simple control by one process over the other than to coordinated development between apical and cambial meristems (Cuny *et al.*, 2012; Zhang *et al.*, 2019).

At the same time, we do not suggest that cambial growth is simply controlled by newly formed leaves or by current assimilate supply. Primary and secondary growth have partially distinct controls and constraints, and cause-and-effect relationships between them are not straightforward (Hartmann & Trumbore, 2016; Trugman & Anderegg, 2025). Rather, from a whole-plant perspective, sustained production of additional leaves, xylem, and phloem is likely to be coordinated because each entails construction and maintenance costs that are only advantageous if matched by corresponding hydraulic and carbon-transport demands. In this sense, variation in shoot (in)determinacy may provide insight into how woody plants coordinate continued axis extension with continued secondary tissue production over the course of a growing season. In highly indeterminate species, continued primary growth later in the season may therefore be viewed not only as prolonged leaf or shoot production, but as part of a broader developmental syndrome in which the extension of axes and the continuation of cambial activity remain jointly permissive for longer.

Although vascular development is not preformed in the same mechanistic sense as shoot organs enclosed in buds, seasonal carry-over in vascular tissues—particularly where conductive phloem persists through winter—may represent a functional analogue of preformation, in that part of the transport system available early in the next season was established previously (Savage, 2020; Savage & Chuine, 2021). This suggests that concepts related to developmental carry-over may extend beyond primary

shoot growth, while still remaining distinct from preformation *sensu stricto* at the shoot apical meristem. The timing and duration of cambial cell division constrain the rate and seasonal extent of secondary growth by limiting the number of cells entering differentiation, while subsequent cell maturation imposes additional carbon and resource demands that can further restrict wood formation (Lupi *et al.*, 2010). At the same time, cambial cell production can cease even when environmental conditions remain favorable and the canopy is still photosynthetically active, pointing to internal regulation of secondary growth (Buttò *et al.*, 2020; Arend *et al.*, 2024).

Below-ground meristems appear to follow a more opportunistic strategy. Root apical meristems can remain active whenever local conditions permit, with rhizotron studies documenting root growth during winter under suitable soil temperatures (Lyford & Wilson, 1966), suggesting that roots often lack a clearly defined dormant phase (Radville *et al.*, 2016; Marchand *et al.*, 2025). However, the extent to which below-ground growth dynamics reflect direct environmental control, coordination with above-ground growth, or genetically constrained strategies remains unresolved (Abramoff & Finzi, 2015; Makoto *et al.*, 2020; Silvestro *et al.*, 2025; Campioli *et al.*, 2024).

Together, these observations suggest that although (in)determinacy is most rigorously defined at the shoot apical meristem, variation in primary shoot growth may still have important downstream consequences for whole-plant performance. In this sense, shoot-level determinacy may provide a useful lens for studying coordination among plant growth processes, even where other organs operate through different developmental mechanisms.

Future directions

A central challenge emerging from this perspective is to better understand how variation along the continuum of growth (in)determinacy shapes tree performance under changing climatic conditions. Rather than representing discrete strategies, degrees of determinacy define bounded trait spaces that differ among species but can vary with age, provenance, and environmental context. While the degree of determinacy is largely constrained at the species level, evidence from provenance trials shows meaningful intraspecific variation; for example, populations from lower latitudes often exhibit higher frequencies of second flushes than those from northern origins (Rudolph, 1964). Clarifying how these species-specific

347 bounds and population-level adjustments interact with warming, drought, and increasing climatic vari-
348 ability will be critical for understanding how forest communities respond and reassemble under climate
349 change. Extending this framework beyond woody perennials to include herbaceous species and peren-
350 nial grasses may further help test the generality of (in)determinacy as a developmental constraint on
351 growth timing, while highlighting differences arising from life form and the absence of secondary growth.

352

353 One key knowledge gap concerns how different degrees of (in)determinacy help trees avoid or buffer
354 environmental stress while still maintaining the capacity to resume growth, repair tissues, replenish
355 reserves, and compensate for earlier losses within the same season. Resolving the trade-off between
356 escaping unfavorable conditions and exploiting longer or intermittently favorable growing periods is
357 essential for anticipating how forest communities will change over the coming decades.

358

359 Understanding whether shoot-level (in)determinacy is associated with analogous patterns of develop-
360 mental carry-over or seasonal flexibility in other organs may provide insight into the evolution of this
361 trait, especially through cross-species analyses. Both growth forms seem to occur across most species
362 of the same family or clade and hence appear to have evolved repeatedly in different groups (Figure 4),
363 making it difficult to speculate on which strategy is ancestral (but see Hariharan *et al.*, 2016) and how
364 rapidly (in)determinacy can evolve. As data on the degree of (in)determinacy across species accumu-
365 lates (see Box 1), such analyses could potentially provide insights into this and also aid forecasting for
366 unsampled species.

367

368 Finally, future work should aim to link the temporal dynamics of primary (shoot and root) and sec-
369 ondary (cambial) meristems with patterns of reserve storage and reproductive allocation. Integrating
370 shoot increment formation, wood production, non-structural carbohydrate dynamics, and reproduc-
371 tive effort within a common temporal framework will be essential for understanding whole-tree carbon
372 allocation. Correlating annual tree-ring formation with shoot increments—while accounting for the
373 timing of inter-annual phytomer production and distinguishing between preformed and neoformed tis-
374 sues (Figure 2)—offers a promising path toward such integration and toward improving mechanistic
375 predictions of tree growth and forest carbon dynamics under future climates.

376 **Acknowledgments**

377 This text emerged from a grant of the Swiss National Foundation to F.B. (grant number P500PB_210943)
378 and was supported by CFI and NSERC Discovery funding to EMW. We thank J. Ngo for designing
379 figure 2.

Bibliography

- Abramoff, R.Z. & Finzi, A.C. (2015). Are above- and below-ground phenology in sync? *New Phytologist*, 205, 1054–1061.
- Alvarado, M.V., Terrazas, T. & Rojas-Leal, A. (2025). Cambial activity in five species of the tropical dry forest: Dynamics of seasonal changes in the cambial region. *Trees*, 39, 51.
- Arend, M., Hoch, G. & Kahmen, A. (2024). Stem growth phenology, not canopy greening, constrains deciduous tree growth. *Tree Physiology*, 44, tpad160.
- Barthélémy, D. & Caraglio, Y. (2007). Plant Architecture: A Dynamic, Multilevel and Comprehensive Approach to Plant Form, Structure and Ontogeny. *Annals of Botany*, 99, 375–407.
- Basler, D. & Körner, C. (2012). Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology*, 165, 73–81.
- Baumgarten, F., Gessler, A. & Vitasse, Y. (2023). No risk—no fun: Penalty and recovery from spring frost damage in deciduous temperate trees. *Functional Ecology*, 37, 648–663.
- Baumgarten, F., Zohner, C.M., Gessler, A. & Vitasse, Y. (2021). Chilled to be forced: The best dose to wake up buds from winter dormancy. *New Phytologist*, 230, 1366–1377.
- Böhlenius, H., Huang, T., Charbonnel-Campaa, L., Brunner, A.M., Jansson, S., Strauss, S.H. & Nilsson, O. (2006). CO/FT Regulatory Module Controls Timing of Flowering and Seasonal Growth Cessation in Trees. *Science*, 312, 1040–1043.
- Borchert, R. (1976). The concept of juvenility in woody plants. *Acta Horticulturae*, pp. 21–36.
- Borchert, R. (1994). Soil and Stem Water Storage Determine Phenology and Distribution of Tropical Dry Forest Trees. *Ecology*, 75, 1437–1449.
- Bose, A.K., Etzold, S., Meusburger, K., Gessler, A., Baltensweiler, A., Braun, S., Buchmann, N., Camarero, J.J., Haeni, M., Kahmen, A., Peters, R.L., Sterck, F.J., Tresch, S., Walthert, L. & Zweifel, R. (2025). Decreasing Stem Growth in Common European Tree Species Despite Earlier Growth Onset. *Global Change Biology*, 31, e70318.

405 Botero, C.A., Weissing, F.J., Wright, J. & Rubenstein, D.R. (2015). Evolutionary tipping points in
 406 the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences*,
 407 112, 184–189.

408 Brienens, R.J.W., Caldwell, L., Duchesne, L., Voelker, S., Barichivich, J., Baliva, M., Ceccantini, G.,
 409 Di Filippo, A., Helama, S., Locosselli, G.M., Lopez, L., Piovesan, G., Schöngart, J., Villalba, R.
 410 & Gloor, E. (2020). Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature*
 411 *Communications*, 11, 4241.

412 Buttò, V., Shishov, V., Tychkov, I., Popkova, M., He, M., Rossi, S., Deslauriers, A. & Morin, H. (2020).
 413 Comparing the Cell Dynamics of Tree-Ring Formation Observed in Microcores and as Predicted by
 414 the Vaganov–Shashkin Model. *Frontiers in Plant Science*, 11.

415 Caffarra, A. & Donnelly, A. (2011). The ecological significance of phenology in four different tree
 416 species: Effects of light and temperature on bud burst. *International Journal of Biometeorology*, 55,
 417 711–721.

418 Campioli, M., Marchand, L.J., Zahnd, C., Zuccarini, P., McCormack, M.L., Landuyt, D., Lorer, E.,
 419 Delpierre, N., Gričar, J. & Vitasse, Y. (2024). Environmental Sensitivity and Impact of Climate
 420 Change on leaf-, wood- and root Phenology for the Overstory and Understory of Temperate Decid-
 421 uous Forests. *Current Forestry Reports*, 11, 1.

422 Chapin, F.S., Schulze, E. & Mooney, H.A. (1990). The Ecology and Economics of Storage in Plants.
 423 *Annual Review of Ecology and Systematics*, 21, 423–447.

424 Chevin, L.M., Lande, R. & Mace, G.M. (2010). Adaptation, Plasticity, and Extinction in a Changing
 425 Environment: Towards a Predictive Theory. *PLOS Biology*, 8, e1000357.

426 Cuny, H.E., Rathgeber, C.B., Lebourgeois, F., Fortin, M. & Fournier, M. (2012). Life strategies in
 427 intra-annual dynamics of wood formation: Example of three conifer species in a temperate forest in
 428 north-east France. *Tree Physiology*, 32, 612–625.

429 Damascos, M.A., Prado, C.H.B.A. & Ronquim, C.C. (2005). Bud Composition, Branching Patterns
 430 and Leaf Phenology in Cerrado Woody Species. *Annals of Botany*, 96, 1075–1084.

431 Delpierre, N., Vitasse, Y., Chuine, I., Guillemot, J., Bazot, S., Rutishauser, T. & Rathgeber, C.B.K.
 432 (2016). Temperate and boreal forest tree phenology: From organ-scale processes to terrestrial ecosys-
 433 tem models. *Annals of Forest Science*, 73, 5–25.

434 Ejsmond, M.J., Czarnoński, M., Kapustka, F. & Kozłowski, J. (2010). How to Time Growth and
 435 Reproduction during the Vegetative Season: An Evolutionary Choice for Indeterminate Growers in
 436 Seasonal Environments. *The American Naturalist*, 175, 551–563.

437 Estiarte, M. & Peñuelas, J. (2015). Alteration of the phenology of leaf senescence and fall in winter
 438 deciduous species by climate change: Effects on nutrient proficiency. *Global Change Biology*, 21,
 439 1005–1017.

440 Etzold, S., Sterck, F., Bose, A.K., Braun, S., Buchmann, N., Eugster, W., Gessler, A., Kahmen, A.,
 441 Peters, R.L. & Vitasse, Y. (2021). Number of growth days and not length of the growth period
 442 determines radial stem growth of temperate trees. *Ecology Letters*.

443 Girard, F., Vennetier, M., Ouarmim, S., Caraglio, Y. & Misson, L. (2011). Polycyclism, a fundamental
 444 tree growth process, decline with recent climate change: The example of *Pinus halepensis* Mill. in
 445 Mediterranean France. *Trees*, 25, 311–322.

446 Gordon, D., Damiano, C. & DeJong, T.M. (2006). Preformation in vegetative buds of *Prunus persica*:
 447 Factors influencing number of leaf primordia in overwintering buds. *Tree Physiology*, 26, 537–544.

448 Guédon, Y., Puntieri, J.G., Sabatier, S. & Barthélémy, D. (2006). Relative Extents of Preformation
 449 and Neoformation in Tree Shoots: Analysis by a Deconvolution Method. *Annals of Botany*, 98,
 450 835–844.

451 Hallé, F., Oldeman, R.A.A. & Tomlinson, P.B. (1978). *Tropical Trees and Forests*. Springer, Berlin,
 452 Heidelberg.

453 Hariharan, I.K., Wake, D.B. & Wake, M.H. (2016). Indeterminate growth: Could it represent the
 454 ancestral condition? *Cold Spring Harbor Perspectives in Biology*, 8, a019174.

455 Hartmann, H. & Trumbore, S. (2016). Understanding the roles of nonstructural carbohydrates in
 456 forest trees – from what we can measure to what we want to know. *New Phytologist*, 211, 386–403.

457 Heide, O.M. (1993). Daylength and thermal time responses of budburst during dormancy release in
 458 some northern deciduous trees. *Physiologia Plantarum*, 88, 531–540.

459 Heuret, P., Meredieu, C., Coudurier, T., Courdier, F. & Barthélémy, D. (2006). Ontogenetic trends
 460 in the morphological features of main stem annual shoots of *Pinus pinaster* (Pinaceae). *American*
 461 *Journal of Botany*, 93, 1577–1587.

- 462 Hover, A., Buissart, F., Caraglio, Y., Heinz, C., Pailler, F., Ramel, M., Vennetier, M., Prévosto, B.
463 & Sabatier, S. (2017). Growth phenology in *Pinus halepensis* Mill.: Apical shoot bud content and
464 shoot elongation. *Annals of Forest Science*, 74, 39.
- 465 Hsiao, T.C. (1973). Plant responses to water stress. *Annual review of plant physiology*, 24, 519–570.
- 466 Hu, Z., Wang, H., Dai, J., Ge, Q. & Lin, S. (2022). Stronger Spring Phenological Advance in Future
467 Warming Scenarios for Temperate Species With a Lower Chilling Sensitivity. *Frontiers in Plant*
468 *Science*, 13.
- 469 Huijser, P. & Schmid, M. (2011). The control of developmental phase transitions in plants. *Develop-*
470 *ment*, 138, 4117–4129.
- 471 Jin, S., Zhang, W., Shao, J., Wan, P., Cheng, S., Cai, S., Yan, G. & Li, A. (2022). Estimation of Larch
472 Growth at the Stem, Crown, and Branch Levels Using Ground-Based LiDAR Point Cloud. *Journal*
473 *of Remote Sensing*, 2022.
- 474 Journé, V., Szymkowiak, J., Foest, J., Hacket-Pain, A., Kelly, D. & Bogdziewicz, M. (2024). Summer
475 solstice orchestrates the subcontinental-scale synchrony of mast seeding. *Nature Plants*, 10, 367–373.
- 476 Karkach, A.S. (2006). Trajectories and models of individual growth. *Demographic Research*, 15, 347–
477 400.
- 478 Keenan, T.F. & Richardson, A.D. (2015). The timing of autumn senescence is affected by the timing
479 of spring phenology: Implications for predictive models. *Global Change Biology*, 21, 2634–2641.
- 480 Kikuzawa, K. (1983). Leaf survival of woody plants in deciduous broad-leaved forests. 1. Tall trees.
481 *Canadian Journal of Botany*, 61, 2133–2139.
- 482 Körner, C. (2008). Winter crop growth at low temperature may hold the answer for alpine treeline
483 formation. *Plant Ecology & Diversity*, 1, 3–11.
- 484 Kovaleski, A.P., Londo, J.P. & Finkelstein, K.D. (2019). X-ray phase contrast imaging of *Vitis* spp.
485 buds shows freezing pattern and correlation between volume and cold hardiness. *Scientific Reports*,
486 9, 14949.
- 487 Kozłowski, T.T. (2012). *Seed Germination, Ontogeny, and Shoot Growth*. Elsevier.
- 488 Kozłowski, T.T. & Pallardy, S.G. (1997). *Growth Control in Woody Plants*. Elsevier.

489 Kuster, T.M., Dobbertin, M., Günthardt-Goerg, M.S., Schaub, M. & Arend, M. (2014). A Phenological
490 Timetable of Oak Growth under Experimental Drought and Air Warming. *PLoS ONE*, 9, e89724.

491 Lang, G.A., Early, J.D., Martin, G.C. & Darnell, R.L. (1987). Endo-, Para-, and Ecodormancy:
492 Physiological Terminology and Classification for Dormancy Research. *HortScience*, 22, 371–377.

493 Laube, J., Sparks, T.H., Estrella, N., Hoffer, J., Ankerst, D.P. & Menzel, A. (2014). Chilling outweighs
494 photoperiod in preventing precocious spring development. *Global Change Biology*, 20, 170–82.

495 Lechowicz, M.J. (1984). Why Do Temperate Deciduous Trees Leaf Out at Different Times? Adaptation
496 and Ecology of Forest Communities. *The American Naturalist*.

497 Leites, L. & Benito Garzón, M. (2023). Forest tree species adaptation to climate across biomes:
498 Building on the legacy of ecological genetics to anticipate responses to climate change. *Global
499 Change Biology*, 29, 4711–4730.

500 Luo, T., Liu, X., Zhang, L., Li, X., Pan, Y. & Wright, I.J. (2018). Summer solstice marks a seasonal
501 shift in temperature sensitivity of stem growth and nitrogen-use efficiency in cold-limited forests.
502 *Agricultural and Forest Meteorology*, 248, 469–478.

503 Lupi, C., Morin, H., Deslauriers, A. & Rossi, S. (2010). Xylem phenology and wood production:
504 Resolving the chicken-or-egg dilemma. *Plant, Cell & Environment*, 33, 1721–1730.

505 Lyford, W. & Wilson, B. (1966). Controlled growth of forest tree roots: Technique and application.
506 *Harvard Forest Paper*.

507 Makoto, K., Wilson, S.D., Sato, T., Blume-Werry, G. & Cornelissen, J.H.C. (2020). Synchronous and
508 asynchronous root and shoot phenology in temperate woody seedlings. *Oikos*, 129, 643–650.

509 Marchand, L.J., Gričar, J., Zuccarini, P., Dox, I., Mariën, B., Verlinden, M., Heinecke, T., Prislan,
510 P., Marie, G. & Lange, H. (2025). No winter halt in below-ground wood growth of four angiosperm
511 deciduous tree species. *Nature Ecology & Evolution*, pp. 1–9.

512 Marks, P.L. (1975). On the Relation between Extension Growth and Successional Status of Deciduous
513 Trees of the Northeastern United States. *Bulletin of the Torrey Botanical Club*, 102, 172–177.

514 Maurya, J.P. & Bhalerao, R.P. (2017). Photoperiod- and temperature-mediated control of growth
515 cessation and dormancy in trees: A molecular perspective. *Annals of Botany*, 120, 351–360.

516 McDaniel, C.N. (1992). Induction and Determination: Developmental Concepts. *Flowering Newsletter*,
517 pp. 3–6.

518 Meng, F., Felton, A.J., Mao, J., Cong, N., Smith, W.K., Körner, C., Hu, Z., Hong, S., Knott, J., Yan,
519 Y., Guo, B., Deng, Y., Leisz, S., Dorji, T., Wang, S. & Chen, A. (2024). Consistent time allocation
520 fraction to vegetation green-up versus senescence across northern ecosystems despite recent climate
521 change. *Science Advances*, 10, eadn2487.

522 Millet, J., Bouchard, A. & Édelin, C. (1999). Relationship between architecture and successional status
523 of trees in the temperate deciduous forest. *Écoscience*, 6, 187–203.

524 Moore, E. (1909). The Study of Winter Buds with Reference to Their Growth and Leaf Content.
525 *Bulletin of the Torrey Botanical Club*, 36, 117–145.

526 O’Sullivan, O.S., Heskell, M.A., Reich, P.B., Tjoelker, M.G., Weerasinghe, L.K., Penillard, A., Zhu,
527 L., Egerton, J.J.G., Bloomfield, K.J., Creek, D., Bahar, N.H.A., Griffin, K.L., Hurry, V., Meir, P.,
528 Turnbull, M.H. & Atkin, O.K. (2017). Thermal limits of leaf metabolism across biomes. *Global*
529 *Change Biology*, 23, 209–223.

530 Poethig, R.S. (2003). Phase Change and the Regulation of Developmental Timing in Plants. *Science*,
531 301, 334–336.

532 Pugnaire, F.I., Serrano, LUIS. & Pardos, JOSE. (1999). Constraints by water stress on plant growth.
533 *Handbook of plant and crop stress*, 2, 271–283.

534 Radville, L., McCormack, M.L., Post, E. & Eissenstat, D.M. (2016). Root phenology in a changing
535 climate. *Journal of Experimental Botany*, 67, 3617–3628.

536 Ramos, D.M., Andrade, J.M., Alberton, B.C., Moura, M.S.B., Domingues, T.F., Neves, N., Lima,
537 J.R.S., Souza, R., Souza, E., Silva, J.R., Espírito-Santo, M.M., Morellato, L.P.C. & Cunha, J.
538 (2023). Multiscale phenology of seasonally dry tropical forests in an aridity gradient. *Frontiers in*
539 *Environmental Science*, 11.

540 Richardson, A.D., Hollinger, D.Y., Dail, D.B., Lee, J.T., Munger, J.W. & O’keefe, J. (2009). Influence
541 of spring phenology on seasonal and annual carbon balance in two contrasting New England forests.
542 *Tree Physiology*, 29, 321–331.

543 Richter, S., Kipfer, T., Wohlgemuth, T., Calderón Guerrero, C., Ghazoul, J. & Moser, B. (2012). Phe-
544 notypic plasticity facilitates resistance to climate change in a highly variable environment. *Oecologia*,
545 169, 269–279.

546 Roskilly, B. & Aitken, S. (2024). Weak Local Adaptation to Climate in Seedlings of a Deciduous
547 Conifer Suggests Limited Benefits and Risks of Assisted Gene Flow. *Evolutionary Applications*, 17,
548 e70001.

549 Rossi, S., Deslauriers, A., Anfodillo, T., Morin, H., Saracino, A., Motta, R. & Borghetti, M. (2006).
550 Conifers in cold environments synchronize maximum growth rate of tree-ring formation with day
551 length. *New Phytologist*, 170, 301–310.

552 Rossi, S., Deslauriers, A., Gricar, J., Seo, J.W., Rathgeber, C.B.K., Anfodillo, T., Morin, H., Levanic,
553 T., Oven, P. & Jalkanen, R. (2008). Critical temperatures for xylogenesis in conifers of cold climates.
554 *Global Ecology and Biogeography*, 17, 696–707.

555 Rudolph, T.D. (1964). Lammas Growth and Prolepsis in Jack Pine in the Lake States. *Forest Science*,
556 10, a0001–70.

557 Sakai, A. & Larcher, W. (1987). *Freezing Injuries in Plants*, Springer Berlin Heidelberg, Berlin,
558 Heidelberg, vol. 62, pp. 39–58.

559 Savage, J.A. (2020). It’s all about timing—or is it? Exploring the potential connection between phloem
560 physiology and whole plant phenology. *American Journal of Botany*, 107, 848–851.

561 Savage, J.A. & Chuine, I. (2021). Coordination of spring vascular and organ phenology in deciduous
562 angiosperms growing in seasonally cold climates. *New Phytologist*, 230, 1700–1715.

563 Schenker, G., Lenz, A., Körner, C. & Hoch, G. (2014). Physiological minimum temperatures for root
564 growth in seven common European broad-leaved tree species. *Tree Physiology*, 34, 302–313.

565 Silvestro, R., Deslauriers, A., Prislan, P., Rademacher, T., Rezaie, N., Richardson, A.D., Vitasse, Y.
566 & Rossi, S. (2025). From Roots to Leaves: Tree Growth Phenology in Forest Ecosystems. *Current*
567 *Forestry Reports*, 11, 12.

568 Silvestro, R., Zeng, Q., Buttò, V., Sylvain, J.D., Drolet, G., Mencuccini, M., Thiffault, N., Yuan, S.
569 & Rossi, S. (2023). A longer wood growing season does not lead to higher carbon sequestration.
570 *Scientific Reports*, 13, 4059.

Smith, S.A. & Brown, J.W. (2018). Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany*, 105, 302–314.

Soolanayakanahally, R.Y., Guy, R.D., Street, N.R., Robinson, K.M., Silim, S.N., Albrechtsen, B.R. & Jansson, S. (2015). Comparative physiology of allopatric *Populus* species: Geographic clines in photosynthesis, height growth, and carbon isotope discrimination in common gardens. *Frontiers in plant science*, 6, 528.

Stearns, S.C. (1989). Trade-Offs in Life-History Evolution. *Functional Ecology*, 3, 259.

Stearns, S.C. (1998). *The Evolution Of Life Histories*. Oxford University Press.

Stevens, M.T. & Lindroth, R.L. (2005). Induced resistance in the indeterminate growth of aspen (*Populus tremuloides*). *Oecologia*, 145, 297–305.

Triozzi, P.M., Ramos-Sánchez, J.M., Hernández-Verdeja, T., Moreno-Cortés, A., Allona, I. & Perales, M. (2018). Photoperiodic Regulation of Shoot Apical Growth in Poplar. *Frontiers in Plant Science*, 9, 1030.

Trugman, A.T. & Anderegg, L.D.L. (2025). Source vs sink limitations on tree growth: From physiological mechanisms to evolutionary constraints and terrestrial carbon cycle implications. *New Phytologist*, 245, 966–981.

van der Wiel, K. & Bintanja, R. (2021). Contribution of climatic changes in mean and variability to monthly temperature and precipitation extremes. *Communications Earth & Environment*, 2, 1.

Vitasse, Y., Baumgarten, F., Zohner, C.M., Rutishauser, T., Pietragalla, B., Gehrig, R., Dai, J., Wang, H., Aono, Y. & Sparks, T.H. (2022). The great acceleration of plant phenological shifts. *Nature Climate Change*, 12, 300–302.

Volaire, F., Barkaoui, K., Grémillet, D., Charrier, G., Dangles, O., Lamarque, L.J., Martin-StPaul, N. & Chuine, I. (2023). Is a seasonally reduced growth potential a convergent strategy to survive drought and frost in plants? *Annals of Botany*, 131, 245–254.

Wang, Q., Liu, W., Leung, C.C., Tarté, D.A. & Gendron, J.M. (2024). Plants distinguish different photoperiods to independently control seasonal flowering and growth. *Science*, 383, eadg9196.

597 Wolkovich, E.M., Ettinger, A.K., Chin, A.R., Chamberlain, C.J., Baumgarten, F., Pradhan, K., Man-
598 zanedo, R.D. & Hille Ris Lambers, J. (2025). Why longer seasons with climate change may not
599 increase tree growth. *Nature Climate Change*, 15, 1283–1292.

600 Zani, D., Crowther, T.W., Mo, L., Renner, S.S. & Zohner, C.M. (2020). Increased growing-season
601 productivity drives earlier autumn leaf senescence in temperate trees. *Science*, 370, 1066–1071.

602 Zhang, Y., Wu, H. & Yang, W. (2019). Forests Growth Monitoring Based on Tree Canopy 3D Recon-
603 struction Using UAV Aerial Photogrammetry. *Forests*, 10, 1052.

604 Zohner, C.M., Mirzaghali, L., Renner, S.S., Mo, L., Rebindaine, D., Bucher, R., Palouš, D., Vitasse,
605 Y., Fu, Y.H., Stocker, B.D. & Crowther, T.W. (2023). Effect of climate warming on the timing of
606 autumn leaf senescence reverses after the summer solstice. *Science*, 381, eadf5098.

607 Zohner, C.M., Mo, L., Pugh, T.A.M., Bastin, J.F. & Crowther, T.W. (2020). Interactive climate
608 factors restrict future increases in spring productivity of temperate and boreal trees. *Global Change*
609 *Biology*, 26, 4042–4055.

610 Zohner, C.M., Renner, S.S., Sebald, V. & Crowther, T.W. (2021). How changes in spring and autumn
611 phenology translate into growth-experimental evidence of asymmetric effects. *Journal of Ecology*,
612 109, 2717–2728.

Box 1: Metrics and proxies related to (in)determinacy

To advance our understanding of growth strategies under climate change, we need not only improved methods for quantifying the degree of (in)determinacy—but also empirical approaches that move beyond a simple binary classification. An organ-level metric at the shoot apical meristem captures the defining features of (in)determinacy and allows it to be explicitly defined and quantified. Temporal dynamics of shoot growth provide a related but indirect proxy that can be applied from individual shoots to crowns and entire forest stands.

Ratio of preformed to neoformed organs: The ratio of the number of leaves (or leaf scars) produced during a growing season to the number of leaf primordia preformed in dormant buds provides a direct, mechanistic, and quantitative measure of shoot-level (in)determinacy. Values close to one indicate growth constrained to preformed organs, whereas values greater than one reflect the production of neoformed organs and thus indeterminate development. This ratio should be interpreted relative to the relevant growth period considered. In species with more than one distinct growth phase within a year, estimates may need to integrate all flushes contributing to annual shoot development, or be reported separately for distinct flushes. Because the expression of neoformation depends on environmental conditions, indeterminate species may appear more determinate in short or unfavourable years. Preformation and neoformation have been documented for more than a century (Moore, 1909), and several studies have quantified the number of preformed primordia and compared it with the final number of emerged leaves or phytomers (Damascos *et al.*, 2005; Kikuzawa, 1983; Guédon *et al.*, 2006). Quantification typically relies on bud dissection, but emerging non-destructive approaches such as micro-CT imaging offer promising alternatives for assessing bud structure and preformation (Kovaleski *et al.*, 2019). This metric remains the most conceptually robust approach for defining and quantifying the level of (in)determinacy across species.

Seasonal growth dynamics of the shoot apical meristem: Temporal patterns of shoot elongation provide a complementary perspective on (in)determinacy at the shoot apical meristem. Across species, elongation may occur over a brief period, extend continuously through the growing season, or appear in multiple distinct growth phases, reflecting differences in how apical meristem activity is maintained or reactivated. Such dynamics can be quantified using repeated shoot measurements or phenological observations, and increasingly through high-resolution terrestrial LiDAR and photogrammetric techniques that capture shoot elongation and structural change at the scale of entire crowns or stands (Zhang *et al.*, 2019; Jin *et al.*, 2022). While temporal growth dynamics do not distinguish preformed from neoformed organs directly, they remain closely linked to shoot apical meristem activity and provide a scalable proxy for the seasonal expression of (in)determinacy. Repeated measurements also allow shoot-elongation curves to be reconstructed, whose shape differs systematically among species (single, short sigmoid vs. prolonged or multi-phase extension; Hover *et al.* (2017); Kuster *et al.* (2014)). Specifically, curve-derived parameters such as the duration of shoot expansion relative to the growing season provide a quantitative proxy for the degree of (in)determinacy.




Figure 1: Schematic overview of the discrepancy between the potential growing season and the effectively realized period of vegetative growth (shoot and possibly cambial growth). Green indicates the potential growing season, whereas overlaid coloured areas represent portions of this period that are constrained by different external and internal filters. Environmental factors, such as temperature and soil moisture, can narrow the window of opportunity for vegetative growth when conditions fall outside growth-promoting thresholds (potential light and photoperiodic constraints are not shown). Species-specific phenology can impose an additional filter through dormancy and through prioritization of other developmental processes (e.g. reproduction and storage). The degree of (in)determinacy is illustrated here as two extremes, although we propose that this trait is better understood as continuous.




Figure 2: Determinate and indeterminate growth within one growing season for species producing terminal buds. Commonly all tree species deploy buds during their first (spring) flush from prebuilt and overwintering leaf primordia (LP) protected by bud scales (BS; A). Determinate growing species set buds (B) that are under hormonal suppression to inhibit any further activity of the shoot apical meristem – SAM (paradormancy). The basic unit of a shoot is the phytomer which is composed of a node, a leaf, the axillary bud and an internode (see bottom right). Indeterminate growing species continue to produce new phytomers directly (C) or through one (D) to several (E) intermediate bud stage(s). Finally, most species set their terminal buds (F) and enter full dormancy (endodormancy).




Figure 3: Hypothesized predictions of growth rates under current (upper panels) and future (lower panels) climate for determinate and indeterminate growing species. Note that the indeterminate strategy is more exposed to the risk of frost and drought events while the determinate strategy condenses most growth within a safe period. In the current climate the indeterminate strategy is in balance with benefiting from the full climatic growing season in some years with some drawbacks in other years, resulting in the same mean yearly biomass increment, but with a higher variation (right panel). In a future climate the indeterminate strategy might benefit from longer growing seasons, resulting in an overall higher mean annual biomass increment compared to determinate growers, but this will be dependent on the severity of drought that overlaps the potential growth period of determinate species. Triangle outlines represent the activity window for a current (black) versus a future (red) climate.




Figure 4: Phylogenetic tree of plant species colored by growth determinacy trait values, visually representing the evolutionary distribution of growth strategies across taxa. This tree includes species for which shoot growth determinacy data were compiled (mainly from Hallé *et al.* (1978) and Kikuzawa (1983)), with the phylogeny pruned from the comprehensive seed plant tree of Smith & Brown (2018) to match the trait dataset. Growth determinacy was scored as indeterminate (blue), intermediate (yellow) and determinate (red), with mean values calculated per species when multiple entries existed. These values were then mapped as a continuous trait onto the phylogeny using the `contMap()` function from the `phytools` R package.