

1 (In)determinacy in woody perennials: limits and opportunities
2 for timing growth in a changing climate

3 Frederik Baumgarten¹⁻³, Sally Aitken¹, Yann Vitasse³, Robert D Guy¹, EM Wolkovich¹

4 February 4, 2026

5 ¹ Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424
6 Main Mall Vancouver, BC Canada V6T 1Z4.

7 ² Department of Environmental Sciences-Botany, University of Basel, Schönbeinstrasse 6, Basel 4056,
8 Switzerland.

9 ³ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse. 111, Bir-
10 mensdorf 8903, Switzerland

11 **Corresponding Author:** Frederik Baumgarten; frederik.baumgarten@unibas.ch;
12 Tel.: +41 61 207 23 37

13 **E-Mail-Addresses:** sally.aitken@ubc.ca; yann.vitasse@wsl.ch; rob.guy@ubc.ca; e.wolkovich@ubc.ca

14 **Type of article:** Perspective

15 **Short running title:** (In)determinate growth strategies of trees

16 **Keywords:** plant growth, tree phenology, shoot extension, indeterminate growers, carbon sequestra-
17 tion, growing season length, drought, genetic programming, phenotypic plasticity, forest productivity

18 **wordnumbers:** abstract: 200; main text: 4005; box1: 404; number of figures: 4; number of tables: 0

19 **number of references:** 84

20 **Authors contribution:** FB and EM initiated this project. FB took a lead in drafting, designed
21 figures and performed phylogenetic analyses. All authors contributed substantially to the writing of
22 the article.

23 **Data accessibility statement:** The data supporting the phylogeny are archived in the following
24 public repository. Link: <https://doi.org/10.6084/m9.figshare.30294826>.

25 **Conflict of interest statement:** The authors declare no conflict of interest

Abstract

When and how much organisms grow given environmental constraints are fundamental questions in biology, and especially pressing in the context of climate change as we strive to predict biomass production and carbon sequestration. While environmental factors such as temperature and water availability directly regulate plant growth, the timing and rate of growth are also governed by a plant's phenological sequence—its genetically programmed developmental stages. This sequence is critical for predicting climate responses but is often underappreciated.

Here, we leverage the concept of (in-)determinacy in growth and development—which captures how flexibly plants produce organs from preformed cells versus initiating them *de novo*—to propose a new framework for predicting tree growth responses to climate change.

We propose that: 1) determinate growth is an adaptation to strongly seasonal climates, restricting tissue investment to a short favorable period before dormancy; 2) indeterminate species are more exposed to increasing climatic extremes (spring/fall frost, drought) but may recover more effectively from resulting damage; and 3) species with greater indeterminacy are more likely to benefit from lengthening frost-free growing seasons. Thus, future carbon sequestration may depend not only on environmental changes, but also on the degree of determinacy set by intrinsic genetic programming.

Introduction

Investing the right amount of resources in growth at the right time is of crucial importance to the survival and fitness of any living organism. The topic of growth strategies and habits has a long history in science, spanning the fields of genetics, genomics, physiology and ecology across the animal and plant kingdoms (Stearns, 1998). At its core lies the concept of determinacy—the classification of organisms as either reaching a fixed size at maturity or continuing to grow throughout their lives. Like mollusks, fish and reptiles, plants add biomass to their primary bodies as long as they live and are therefore considered ‘indeterminate growers’ (Ejzmond *et al.*, 2010).

Various terms have emerged to describe this phenomenon across spatial and temporal scales, e.g. from a single cell to the whole organism, and from a single season to an entire lifetime (McDaniel, 1992; Karkach, 2006). In plants, organs can develop from preformed embryonic tissue (e.g. primordia in overwintering buds), form *de novo*, or both, depending on species and life form. While these underlying developmental and meristematic constraints apply broadly across the plant kingdom, they play a particularly central role in the life cycle of woody perennials, especially trees, which must balance growth, survival, and reproduction over long lifespans in highly seasonal environments. As we use it here, (in)determinacy in woody perennials refers to the ability to:

- a) preform organs in one year as a future investment, which is deployed rapidly in the following spring with no further primary shoot growth during that season (determinate strategy)
- b) maintain a continuous or episodic meristem activity by forming new shoot tissue during the current growing season (indeterminate strategy).

In this perspective we revisit classical concepts of determinacy and growth patterns of the shoot apical meristem in woody perennials, generalize them across meristem types and position them within the context of climate change. Although we focus here on woody perennials, the concept of (in)determinacy is broadly applicable across vascular plants, including herbaceous species; trees are used as a model system in which the ecological and climatic consequences of determinacy are particularly pronounced. To provide context, we briefly introduce the fundamental limitations to plant growth in extra-tropical regions by linking external environmental drivers and internal growth control mechanisms.

Seasonality of temperature, soil moisture and light

The further one travels from the equator towards the poles, the more strongly plants are confined to a shrinking time window of opportunity for growth set primarily by low temperatures. Below c. 5 °C, metabolic activity slows to an extent where growth and development largely cease (Schenker *et al.*, 2014; Rossi *et al.*, 2008; Körner, 2008). More importantly, sub-freezing temperatures can cause severe damage to plant tissue if they occur at vulnerable developmental stages, e.g. during or after leaf unfolding, prior to fruit maturation, or before cold acclimation in fall (Sakai & Larcher, 1987; Baumgarten *et al.*, 2023). While annual plant species accommodate their entire life cycle within this window, perennial (woody) species are forced to partition their growing phase seasonally, with periods of activity alternating with periods of dormancy (Delpierre *et al.*, 2016). This is referred to as intermittent or rhythmic (as opposed to continuous) growth.

High temperatures can directly inhibit growth by exceeding species-specific physiological thresholds (O’Sullivan *et al.*, 2017). In addition, increased evaporative demand and reduced precipitation can lower soil moisture, leading to water stress that slows or stalls growth and, in extreme cases, causes leaf scorching or premature senescence (Hsiao, 1973; Pugnaire *et al.*, 1999; Etzold *et al.*, 2021; Estiarte & Peñuelas, 2015). Together, these temperature and soil moisture limitations act as environmental filters, narrowing the period during which growth is possible (Figure 1).

Light also influences plant growth through two key processes. First, light intensity provides the primary energy source that drives photosynthesis and, consequently, source activity. Second, light carries a wealth of information through its quality and the length of daily light and dark periods (photoperiod), which mediate many physiological processes, including bud set and dormancy transitions (Wang *et al.*, 2024). Both factors may contribute to the common observation that tree growth often peaks near the summer solstice in extra-tropical regions (Rossi *et al.*, 2006; Etzold *et al.*, 2021; Luo *et al.*, 2018) or that a window of temperature sensitivity ‘opens’ around the summer solstice to synchronize reproductive efforts over very large spatial scales (Journé *et al.*, 2024).

Internal programming of plants

Given our relatively robust understanding of environmental influences on plant growth, it may seem straightforward to predict how plants will grow under current and future climatic conditions. Yet,

this has proven challenging, particularly when predicting responses under scenarios with extended, climatically favorable growing seasons (Zohner *et al.*, 2021). Here we propose that accurate predictions require incorporating another key, but often overlooked, factor: internal growth control—the genetically encoded developmental program that may prevent further growth *despite* favorable environmental conditions.

While plants have evolved numerous morphological adaptations to tolerate or avoid harmful environmental conditions, most temperate and boreal woody perennials cope with seasonal fluctuations in temperature and moisture by temporally escaping these conditions. They do this by aligning their life-history events (phenology) with a cycle of growth and dormancy that balances survival, reproduction, and growth over the long lifespan typical of many tree species (Delpierre *et al.*, 2016). Hence, the phenological sequence can impose abrupt internal switches in resource allocation—from vegetative growth to reproduction (flowering, fruit maturation) and storage (Stearns, 1989; Chapin *et al.*, 1990). These transitions act as additional internal filters, further narrowing the effective window during which vegetative growth can occur, regardless of environmental potential (Figure 1).

The concept of (in)determinate growth

In annual plants, growth ends with the production of flowers to form fruits and seeds: a signal in the apical meristem triggers a sudden switch in resource investment from vegetative growth to building a reproductive structure with no point of return, signaling the end of the life-cycle (Poethig, 2003; Huijser & Schmid, 2011). A shoot axis that terminates in a flower or other modified structure and ceases meristematic activity is considered ‘determinate’ (Barthélémy & Caraglio, 2007). In woody perennial species, however, flowers are typically produced on lateral shoots or buds, preserving the main vegetative axis and allowing continued structural expansion. However, ‘determinate growth’ also refers to a temporal suspension of the meristem, which leads to a time lag between the formation of organs and their expansion (Kozlowski, 2012; Hallé *et al.*, 1978).

In their first year, all seedlings exhibit indeterminate growth. In subsequent years, many temperate and boreal tree species preform their new shoot increments during the previous growing season, overwintering them as primordial shoots with leaves and internodes in hardened buds to be ‘ready-to-go’ when spring arrives. Once the leaves flush and the internodes elongate, some species (e.g. *Fraxinus*

spp., *Prunus* spp. and *Acer* spp.) suppress further activity of the apical meristem for the rest of the season through internal control mechanisms such as paradormancy (Lang *et al.*, 1987), often regulated by photoperiod-induced hormonal changes (Böhlenius *et al.*, 2006). Other species, by contrast, continue to produce neoformed leaves and internodes on top of the preformed shoot, sometimes stretching their indeterminate growth well into autumn until low temperature eventually forces them to stop (e.g. *Alnus* spp., *Betula* spp. and *Sequoia* spp.). Indeterminate growth can occur either through the continuous activity of the shoot apical meristems forming new leaves and internodes without buds, termed free growth, or through the formation and subsequent flushing of buds mid-season without a period of dormancy, called second flushing or lammas growth (Figure 2).

Although this concept is often presented dichotomously (Kozłowski & Pallardy, 1997; Lechowicz, 1984), but see Kikuzawa, 1983; Damascos *et al.*, 2005), with species classified as either determinate or indeterminate growers, most species likely exist along a gradient with numerous intermediate forms. For instance, many oaks (*Quercus* spp.) are considered determinate growers, but can exhibit multiple flushes. Similarly, Douglas-fir (*Pseudotsuga menziesii*) gradually adopts a more determinate growth habit with maturity (Borchert, 1976; Heuret *et al.*, 2006). However, a gradual quantification of the degree of determinacy on the species level is possible at least for the shoot apical meristem (Box 1). See Figure 4 for a literature-based classification of determinacy of some common species with many among them falling into an intermediate position.

Evolutionary trade-offs: the cost of certainty and flexibility

In most temperate and boreal ecosystems, determinate and indeterminate growth strategies co-occur, indicating that neither strategy is universally superior. Instead, they reflect contrasting solutions to a fundamental trade-off between growth certainty and growth flexibility under seasonal and variable environments.

Indeterminate species are characterized by continued or episodic activity of the shoot apical meristem during the growing season. This does not necessarily imply faster intrinsic growth rates, but rather a greater capacity to extend growth in time when conditions remain favorable. Such temporal flexibility allows individuals to exploit transient resource availability and recover from partial damage, traits

that are advantageous in disturbed or early-successional environments. Accordingly, indeterminate growth is frequently associated with early successional species (Marks, 1975). However, maintaining active meristems later into the season also increases exposure to environmental stressors such as summer drought and early autumn frost, thereby raising the risk of tissue loss or incomplete maturation (Figure 4). As a result, indeterminate growth strategies often involve higher interannual variability in performance and, in some cases, shorter tissue or organismal lifespans (Millet *et al.*, 1999; Brien *et al.*, 2020).

In contrast, determinate species preform a substantial portion of their shoot growth in advance and deploy this tissue during a restricted period of the growing season. This strategy limits short-term plasticity, as growth is largely constrained by conditions experienced during bud formation in the previous year. At the same time, it reduces exposure of sensitive tissues to late-season stress and synchronizes growth with the most predictable and favorable part of the season. As a consequence, determinate growth strategies are often associated with stronger local adaptation and reduced sensitivity to short-term climatic fluctuations (Leites & Benito Garzón, 2023).

These contrasting strategies help explain the coexistence of determinate and indeterminate species within the same landscapes. For example, western larch (*Larix occidentalis*) and Douglas-fir (*Pseudotsuga menziesii*) frequently co-occur despite marked differences in growth determinacy. Western larch exhibits more flexible, indeterminate shoot growth and performs well in environments shaped by frequent disturbance, whereas Douglas-fir shows more determinate growth and stronger local adaptation, conferring stability under predictable seasonal regimes (Roskilly & Aitken, 2024). Such differences suggest niche differentiation along gradients of disturbance frequency, environmental stress, and climatic predictability, rather than outcomes driven by straightforward competitive dominance.

Together, these patterns indicate that growth determinacy reflects a trade-off between flexibility and reliability in seasonal environments. Under ongoing climate change—marked by longer growing seasons, greater climatic variability, and increasing environmental stress in many regions—it remains uncertain whether determinate or indeterminate growth strategies will be favored. Resolving both how trees regulate their degree of (in)determinacy and which strategy advantages under future conditions will be critical for predicting shifts in species performance, coexistence, and forest composition.

The performance of (in)determinacy with climate change

Spring warming has advanced the onset of leaf emergence by up to a month compared to pre-industrial times (Vitasse *et al.*, 2022), reflecting an earlier reactivation of shoot apical and cambial meristems under warmer conditions. In contrast, the cessation of growth in autumn has shifted far less than expected given the extension of climatically favorable conditions (Zani *et al.*, 2020; Zohner *et al.*, 2023). As a result, phenological sequences are increasingly observed to shift as a whole toward spring rather than being symmetrically extended at both ends of the growing season (Keenan & Richardson, 2015; Meng *et al.*, 2024). This asymmetric response highlights a key constraint relevant to growth (in)determinacy: while growth onset is highly plastic and environmentally responsive, the duration of meristematic activity later in the season appears more tightly regulated and differs markedly among species. Such constraints on growth cessation help explain why longer growing seasons do not necessarily translate into increased growth or biomass production (Zani *et al.*, 2020; Wolkovich *et al.*, 2025).

Along the continuum of growth determinacy, we hypothesize that species positioned toward the more determinate end will largely maintain their fixed seasonal growth schedules under climate warming, with limited capacity to extend growth later into the season. In regions where climate change primarily increases heat and water stress during mid to late summer, such species are therefore expected to show little change in annual biomass production, as their restricted growth window increasingly overlaps with unfavorable conditions (Figure 3).

In contrast, we predict that species toward the indeterminate end of the continuum will be more capable of prolonging meristematic activity when conditions remain favorable, potentially resulting in higher annual production under some future climate scenarios (Caffarra & Donnelly, 2011; Richardson *et al.*, 2009; Zohner *et al.*, 2020, Figure 3). This capacity for temporal extension of growth is particularly common among early successional species, which tend to leaf out earlier in spring and senesce later in autumn than late-successional species, consistent with their generally lower chilling and forcing requirements for bud burst and dormancy release (Heide, 1993; Laube *et al.*, 2014; Hu *et al.*, 2022; Baumgarten *et al.*, 2021).

Extreme climatic events and increasing environmental stress provide an important context for evaluating these predictions. More determinate growth strategies concentrate growth earlier in the season and may therefore escape unfavorable growing conditions by operating within a relatively narrow

window of potential growth. Under historical climatic regimes, this strategy provided ample safety margins against late-season stress. Where climate change introduces novel or intensified drought stress within this window, however, determinate species are likely to experience reduced annual production and limited capacity for recovery within the same growing season once canopy development has been compromised (Silvestro *et al.*, 2023, Figure 3).

Species with more indeterminate growth habits often remain green and physiologically active for longer periods, leafing out early in spring and senescing late in autumn, occasionally retaining functional foliage until the first freezing events (Figure 3). As a consequence, a substantial fraction of their growth period coincides with parts of the growing season during which drought stress is projected to intensify under future climates, particularly in water-limited regions (Figure 3). While this extended activity increases exposure to environmental extremes, it also creates opportunities for compensatory growth when conditions remain favorable.

In essence, the flexible growth schedule of indeterminately growing species may allow plants to: (1) produce tissue better adapted to harsh environmental conditions as it is formed in the current season (e.g., smaller leaves with higher specific leaf area), and (2) catch up and compensate later in the season by another production boost (Stevens & Lindroth, 2005). Mediterranean ecosystems provide a clear example of this flexibility, where many species suspend growth during summer drought but resume meristematic activity in autumn or winter when conditions allow (Volaire *et al.*, 2023). Evidence that trees can take advantage of such a ‘second growing season’ after summer drought has been found in pines in Mediterranean climates through polycyclic flushing—a form of indeterminate growth (Girard *et al.*, 2011, Figure 3).

We therefore hypothesize that climate change will increasingly disrupt established phenological cycles, favoring species and genotypes that are more plastic in rearranging their activities by resuming growth and/or restoring reserves (e.g., starch, nutrients) later in the year, thereby recovering from and compensating for stress-induced damage and losses. Such differential capacity to exploit newly available growth opportunities or to recover after stress is expected to intensify competition among co-occurring species by altering the timing and magnitude of resource uptake, potentially leading to the reassembly of forest communities toward species occupying more indeterminate positions along the growth continuum.

(In)determinacy beyond leaf tissue

Although (in)determinacy is most clearly defined *sensu stricto* at the shoot apical meristem, its relevance extends beyond primary shoot growth because shoot extension imposes fundamental structural constraints on whole-plant growth. Primary growth governs crown expansion and annual leaf deployment, thereby shaping total leaf area and canopy photosynthetic capacity—key determinants of biomass production that have often been underappreciated in studies of tree growth (Girard *et al.*, 2011; Soolanayakanahally *et al.*, 2015). At the same time, primary shoot growth establishes the vascular framework within which secondary growth occurs, such that the size and extent of a shoot increment determine the length and surface area over which cambial activity can later be expressed. Empirical studies report close coordination between primary shoot growth and stem or radial growth within seasons, indicating that variation in shoot extension is tightly linked to variation in wood formation (Cuny *et al.*, 2012; Zhang *et al.*, 2019).

Despite its central role in tree development, primary shoot growth remains far less frequently measured than radial growth, an imbalance that is striking given that shoot extension occurs at the decimeter scale, while radial growth—expressed at the micrometer scale—can be monitored continuously and with relative ease using dendrometers or microcoring (see Box 1 for alternative approaches to quantify primary growth).

In the vascular cambium, seasonal patterns of cell division and differentiation are temporally structured but cannot be interpreted in terms of preformation or neoformation. Cambial initial cells divide to produce daughter cells that mature into functional xylem and phloem (Valdovinos-Ayala *et al.*, 2022). The timing and duration of cambial cell division constrain the rate and seasonal extent of secondary growth by limiting the number of cells entering differentiation, while subsequent cell maturation imposes additional carbon and resource demands that can further restrict wood formation (Lupi *et al.*, 2010). At the same time, cambial cell production can cease even when environmental conditions remain favorable and the canopy is still photosynthetically active, pointing to internal regulation of secondary growth (Buttò *et al.*, 2020; Arend *et al.*, 2024).

Below-ground meristems appear to follow a more opportunistic strategy. Root apical meristems can remain active whenever local conditions permit, with rhizotron studies documenting root growth during winter under suitable soil temperatures (Lyford & Wilson, 1966), suggesting that roots often lack a clearly defined dormant phase (Radville *et al.*, 2016; Marchand *et al.*, 2025). However, the extent to which below-ground growth dynamics reflect direct environmental control, coordination with above-

287 ground growth, or genetically constrained strategies remains unresolved (Abramoff & Finzi, 2015;
288 Makoto *et al.*, 2020; Silvestro *et al.*, 2025; Campioli *et al.*, 2024).
289 Together, these observations highlight that while (in)determinacy is most rigorously defined at the
290 shoot apical meristem, primary shoot growth exerts a disproportionate influence on whole-plant per-
291 formance by simultaneously shaping canopy source capacity and the structural context for downstream
292 sink activity. As such, determinacy at the shoot level provides a meaningful organizing principle for
293 understanding growth regulation across plant organs, even where analogous processes operate through
294 different mechanisms.

Future directions

A central challenge emerging from this perspective is to better understand how variation along the continuum of growth (in)determinacy shapes tree performance under changing climatic conditions. Rather than representing discrete strategies, degrees of determinacy define bounded trait spaces that differ among species but can vary with age, provenance, and environmental context. While the degree of determinacy is largely constrained at the species level, evidence from provenance trials shows meaningful intraspecific variation; for example, populations from lower latitudes often exhibit higher frequencies of second flushes than those from northern origins (Rudolph, 1964). Clarifying how these species-specific bounds and population-level adjustments interact with warming, drought, and increasing climatic variability will be critical for understanding how forest communities respond and reassemble under climate change. Extending this framework beyond woody perennials to include herbaceous species and perennial grasses may further help test the generality of (in)determinacy as a developmental constraint on growth timing, while highlighting differences arising from life form and the absence of secondary growth.

One key knowledge gap concerns how different degrees of (in)determinacy help trees avoid or buffer environmental stress while still maintaining the capacity to resume growth, repair tissues, replenish reserves, and compensate for earlier losses within the same season. Resolving the trade-off between escaping unfavorable conditions and exploiting longer or intermittently favorable growing periods is essential for anticipating how forest communities will change over the coming decades.

Understanding how universally tissues within species are formed determinately or indeterminately could provide insight into the evolution of this trait, which could be aided by cross-species analyses. Both growth forms seem to occur across most species of the same family or clade and hence appear to have evolved repeatedly in different groups (Figure 4), making it difficult to speculate on which strategy is ancestral (but see Hariharan *et al.*, 2016) and how rapidly (in)determinacy can evolve. As data on the degree of (in)determinacy across species accumulates (see Box 1), such analyses could potentially provide rapid insights into this and also aid forecasting for unsampled species.

Finally, future work should aim to link the temporal dynamics of primary (shoot and root) and secondary (cambial) meristems with patterns of reserve storage and reproductive allocation. Integrating shoot increment formation, wood production, non-structural carbohydrate dynamics, and reproduc-

326 tive effort within a common temporal framework will be essential for understanding whole-tree carbon
327 allocation. Correlating annual tree-ring formation with shoot increments—while accounting for the
328 timing of inter-annual phytomer production and distinguishing between preformed and neoformed tis-
329 sues (Figure 2)—offers a promising path toward such integration and toward improving mechanistic
330 predictions of tree growth and forest carbon dynamics under future climates.

331 Acknowledgments

332 This text emerged from a grant of the Swiss National Foundation to F.B. (grant number P500PB_210943).
333 We thank J. Ngo for designing figure 2.

Bibliography

- Abramoff, R.Z. & Finzi, A.C. (2015). Are above- and below-ground phenology in sync? *New Phytologist*, 205, 1054–1061.
- Arend, M., Hoch, G. & Kahmen, A. (2024). Stem growth phenology, not canopy greening, constrains deciduous tree growth. *Tree Physiology*, 44, tpad160.
- Barthélémy, D. & Caraglio, Y. (2007). Plant Architecture: A Dynamic, Multilevel and Comprehensive Approach to Plant Form, Structure and Ontogeny. *Annals of Botany*, 99, 375–407.
- Baumgarten, F., Gessler, A. & Vitasse, Y. (2023). No risk—no fun: Penalty and recovery from spring frost damage in deciduous temperate trees. *Functional Ecology*, 37, 648–663.
- Baumgarten, F., Zohner, C.M., Gessler, A. & Vitasse, Y. (2021). Chilled to be forced: The best dose to wake up buds from winter dormancy. *New Phytologist*, 230, 1366–1377.
- Böhlenius, H., Huang, T., Charbonnel-Campaa, L., Brunner, A.M., Jansson, S., Strauss, S.H. & Nilsson, O. (2006). CO/FT Regulatory Module Controls Timing of Flowering and Seasonal Growth Cessation in Trees. *Science*, 312, 1040–1043.
- Borchert, R. (1976). The concept of juvenility in woody plants. *Acta Horticulturae*, pp. 21–36.
- Brienen, R.J.W., Caldwell, L., Duchesne, L., Voelker, S., Barichivich, J., Baliva, M., Ceccantini, G., Di Filippo, A., Helama, S., Locosselli, G.M., Lopez, L., Piovesan, G., Schöngart, J., Villalba, R. & Gloor, E. (2020). Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature Communications*, 11, 4241.
- Buttò, V., Shishov, V., Tyckov, I., Popkova, M., He, M., Rossi, S., Deslauriers, A. & Morin, H. (2020). Comparing the Cell Dynamics of Tree-Ring Formation Observed in Microcores and as Predicted by the Vaganov–Shashkin Model. *Frontiers in Plant Science*, 11.
- Caffarra, A. & Donnelly, A. (2011). The ecological significance of phenology in four different tree species: Effects of light and temperature on bud burst. *International Journal of Biometeorology*, 55, 711–721.
- Campioli, M., Marchand, L.J., Zahnd, C., Zuccarini, P., McCormack, M.L., Landuyt, D., Lorer, E., Delpierre, N., Gričar, J. & Vitasse, Y. (2024). Environmental Sensitivity and Impact of Climate

- 361 Change on leaf-, wood- and root Phenology for the Overstory and Understory of Temperate Decid-
362 uous Forests. *Current Forestry Reports*, 11, 1.
- 363 Chapin, F.S., Schulze, E. & Mooney, H.A. (1990). The Ecology and Economics of Storage in Plants.
364 *Annual Review of Ecology and Systematics*, 21, 423–447.
- 365 Cuny, H.E., Rathgeber, C.B., Lebourgeois, F., Fortin, M. & Fournier, M. (2012). Life strategies in
366 intra-annual dynamics of wood formation: Example of three conifer species in a temperate forest in
367 north-east France. *Tree Physiology*, 32, 612–625.
- 368 Damascos, M.A., Prado, C.H.B.A. & Ronquim, C.C. (2005). Bud Composition, Branching Patterns
369 and Leaf Phenology in Cerrado Woody Species. *Annals of Botany*, 96, 1075–1084.
- 370 Delpierre, N., Vitasse, Y., Chuine, I., Guillemot, J., Bazot, S., Rutishauser, T. & Rathgeber, C.B.K.
371 (2016). Temperate and boreal forest tree phenology: From organ-scale processes to terrestrial ecosys-
372 tem models. *Annals of Forest Science*, 73, 5–25.
- 373 Ejsmond, M.J., Czarnoński, M., Kapustka, F. & Kozłowski, J. (2010). How to Time Growth and
374 Reproduction during the Vegetative Season: An Evolutionary Choice for Indeterminate Growers in
375 Seasonal Environments. *The American Naturalist*, 175, 551–563.
- 376 Estiarte, M. & Peñuelas, J. (2015). Alteration of the phenology of leaf senescence and fall in winter
377 deciduous species by climate change: Effects on nutrient proficiency. *Global Change Biology*, 21,
378 1005–1017.
- 379 Etzold, S., Sterck, F., Bose, A.K., Braun, S., Buchmann, N., Eugster, W., Gessler, A., Kahmen, A.,
380 Peters, R.L. & Vitasse, Y. (2021). Number of growth days and not length of the growth period
381 determines radial stem growth of temperate trees. *Ecology Letters*.
- 382 Girard, F., Vennetier, M., Ouarmim, S., Caraglio, Y. & Misson, L. (2011). Polycyclism, a fundamental
383 tree growth process, decline with recent climate change: The example of *Pinus halepensis* Mill. in
384 Mediterranean France. *Trees*, 25, 311–322.
- 385 Guédon, Y., Puntieri, J.G., Sabatier, S. & Barthélémy, D. (2006). Relative Extents of Preformation
386 and Neoformation in Tree Shoots: Analysis by a Deconvolution Method. *Annals of Botany*, 98,
387 835–844.

- 388 Hallé, F., Oldeman, R.A.A. & Tomlinson, P.B. (1978). *Tropical Trees and Forests*. Springer, Berlin,
389 Heidelberg.
- 390 Hariharan, I.K., Wake, D.B. & Wake, M.H. (2016). Indeterminate growth: Could it represent the
391 ancestral condition? *Cold Spring Harbor Perspectives in Biology*, 8, a019174.
- 392 Heide, O.M. (1993). Daylength and thermal time responses of budburst during dormancy release in
393 some northern deciduous trees. *Physiologia Plantarum*, 88, 531–540.
- 394 Heuret, P., Meredieu, C., Coudurier, T., Courdier, F. & Barthélémy, D. (2006). Ontogenetic trends
395 in the morphological features of main stem annual shoots of *Pinus pinaster* (Pinaceae). *American*
396 *Journal of Botany*, 93, 1577–1587.
- 397 Hover, A., Buissart, F., Caraglio, Y., Heinz, C., Pailler, F., Ramel, M., Vennetier, M., Prévosto, B.
398 & Sabatier, S. (2017). Growth phenology in *Pinus halepensis* Mill.: Apical shoot bud content and
399 shoot elongation. *Annals of Forest Science*, 74, 39.
- 400 Hsiao, T.C. (1973). Plant responses to water stress. *Annual review of plant physiology*, 24, 519–570.
- 401 Hu, Z., Wang, H., Dai, J., Ge, Q. & Lin, S. (2022). Stronger Spring Phenological Advance in Future
402 Warming Scenarios for Temperate Species With a Lower Chilling Sensitivity. *Frontiers in Plant*
403 *Science*, 13.
- 404 Huijser, P. & Schmid, M. (2011). The control of developmental phase transitions in plants. *Develop-*
405 *ment*, 138, 4117–4129.
- 406 Jin, S., Zhang, W., Shao, J., Wan, P., Cheng, S., Cai, S., Yan, G. & Li, A. (2022). Estimation of Larch
407 Growth at the Stem, Crown, and Branch Levels Using Ground-Based LiDAR Point Cloud. *Journal*
408 *of Remote Sensing*, 2022.
- 409 Journé, V., Szymkowiak, J., Foest, J., Hacket-Pain, A., Kelly, D. & Bogdziewicz, M. (2024). Summer
410 solstice orchestrates the subcontinental-scale synchrony of mast seeding. *Nature Plants*, 10, 367–373.
- 411 Karkach, A.S. (2006). Trajectories and models of individual growth. *Demographic Research*, 15, 347–
412 400.
- 413 Keenan, T.F. & Richardson, A.D. (2015). The timing of autumn senescence is affected by the timing
414 of spring phenology: Implications for predictive models. *Global Change Biology*, 21, 2634–2641.

- 415 Kikuzawa, K. (1983). Leaf survival of woody plants in deciduous broad-leaved forests. 1. Tall trees.
416 *Canadian Journal of Botany*, 61, 2133–2139.
- 417 Körner, C. (2008). Winter crop growth at low temperature may hold the answer for alpine treeline
418 formation. *Plant Ecology & Diversity*, 1, 3–11.
- 419 Kovalesski, A.P., Londo, J.P. & Finkelstein, K.D. (2019). X-ray phase contrast imaging of *Vitis* spp.
420 buds shows freezing pattern and correlation between volume and cold hardiness. *Scientific Reports*,
421 9, 14949.
- 422 Kozłowski, T.T. (2012). *Seed Germination, Ontogeny, and Shoot Growth*. Elsevier.
- 423 Kozłowski, T.T. & Pallardy, S.G. (1997). *Growth Control in Woody Plants*. Elsevier.
- 424 Kuster, T.M., Dobbertin, M., Günthardt-Goerg, M.S., Schaub, M. & Arend, M. (2014). A Phenological
425 Timetable of Oak Growth under Experimental Drought and Air Warming. *PLoS ONE*, 9, e89724.
- 426 Lang, G.A., Early, J.D., Martin, G.C. & Darnell, R.L. (1987). Endo-, Para-, and Ecodormancy:
427 Physiological Terminology and Classification for Dormancy Research. *HortScience*, 22, 371–377.
- 428 Laube, J., Sparks, T.H., Estrella, N., Hoffer, J., Ankerst, D.P. & Menzel, A. (2014). Chilling outweighs
429 photoperiod in preventing precocious spring development. *Global Change Biology*, 20, 170–82.
- 430 Lechowicz, M.J. (1984). Why Do Temperate Deciduous Trees Leaf Out at Different Times? Adaptation
431 and Ecology of Forest Communities. *The American Naturalist*.
- 432 Leites, L. & Benito Garzón, M. (2023). Forest tree species adaptation to climate across biomes:
433 Building on the legacy of ecological genetics to anticipate responses to climate change. *Global*
434 *Change Biology*, 29, 4711–4730.
- 435 Luo, T., Liu, X., Zhang, L., Li, X., Pan, Y. & Wright, I.J. (2018). Summer solstice marks a seasonal
436 shift in temperature sensitivity of stem growth and nitrogen-use efficiency in cold-limited forests.
437 *Agricultural and Forest Meteorology*, 248, 469–478.
- 438 Lupi, C., Morin, H., Deslauriers, A. & Rossi, S. (2010). Xylem phenology and wood production:
439 Resolving the chicken-or-egg dilemma. *Plant, Cell & Environment*, 33, 1721–1730.
- 440 Lyford, W. & Wilson, B. (1966). Controlled growth of forest tree roots: Technique and application.
441 *Harvard Forest Paper*.

- 442 Makoto, K., Wilson, S.D., Sato, T., Blume-Werry, G. & Cornelissen, J.H.C. (2020). Synchronous and
443 asynchronous root and shoot phenology in temperate woody seedlings. *Oikos*, 129, 643–650.
- 444 Marchand, L.J., Gričar, J., Zuccarini, P., Dox, I., Mariën, B., Verlinden, M., Heinecke, T., Prislan,
445 P., Marie, G. & Lange, H. (2025). No winter halt in below-ground wood growth of four angiosperm
446 deciduous tree species. *Nature Ecology & Evolution*, pp. 1–9.
- 447 Marks, P.L. (1975). On the Relation between Extension Growth and Successional Status of Deciduous
448 Trees of the Northeastern United States. *Bulletin of the Torrey Botanical Club*, 102, 172–177.
- 449 McDaniel, C.N. (1992). Induction and Determination: Developmental Concepts. *Flowering Newsletter*,
450 pp. 3–6.
- 451 Meng, F., Felton, A.J., Mao, J., Cong, N., Smith, W.K., Körner, C., Hu, Z., Hong, S., Knott, J., Yan,
452 Y., Guo, B., Deng, Y., Leisz, S., Dorji, T., Wang, S. & Chen, A. (2024). Consistent time allocation
453 fraction to vegetation green-up versus senescence across northern ecosystems despite recent climate
454 change. *Science Advances*, 10, eadn2487.
- 455 Millet, J., Bouchard, A. & Édelin, C. (1999). Relationship between architecture and successional status
456 of trees in the temperate deciduous forest. *Écoscience*, 6, 187–203.
- 457 Moore, E. (1909). The Study of Winter Buds with Reference to Their Growth and Leaf Content.
458 *Bulletin of the Torrey Botanical Club*, 36, 117–145.
- 459 O’Sullivan, O.S., Heskell, M.A., Reich, P.B., Tjoelker, M.G., Weerasinghe, L.K., Penillard, A., Zhu,
460 L., Egerton, J.J.G., Bloomfield, K.J., Creek, D., Bahar, N.H.A., Griffin, K.L., Hurry, V., Meir, P.,
461 Turnbull, M.H. & Atkin, O.K. (2017). Thermal limits of leaf metabolism across biomes. *Global*
462 *Change Biology*, 23, 209–223.
- 463 Poethig, R.S. (2003). Phase Change and the Regulation of Developmental Timing in Plants. *Science*,
464 301, 334–336.
- 465 Pugnaire, F.I., Serrano, LUIS. & Pardos, JOSE. (1999). Constraints by water stress on plant growth.
466 *Handbook of plant and crop stress*, 2, 271–283.
- 467 Radville, L., McCormack, M.L., Post, E. & Eissenstat, D.M. (2016). Root phenology in a changing
468 climate. *Journal of Experimental Botany*, 67, 3617–3628.

- Richardson, A.D., Hollinger, D.Y., Dail, D.B., Lee, J.T., Munger, J.W. & O'keefe, J. (2009). Influence of spring phenology on seasonal and annual carbon balance in two contrasting New England forests. *Tree Physiology*, 29, 321–331.
- Roskilly, B. & Aitken, S. (2024). Weak Local Adaptation to Climate in Seedlings of a Deciduous Conifer Suggests Limited Benefits and Risks of Assisted Gene Flow. *Evolutionary Applications*, 17, e70001.
- Rossi, S., Deslauriers, A., Anfodillo, T., Morin, H., Saracino, A., Motta, R. & Borghetti, M. (2006). Conifers in cold environments synchronize maximum growth rate of tree-ring formation with day length. *New Phytologist*, 170, 301–310.
- Rossi, S., Deslauriers, A., Gricar, J., Seo, J.W., Rathgeber, C.B.K., Anfodillo, T., Morin, H., Levanic, T., Oven, P. & Jalkanen, R. (2008). Critical temperatures for xylogenesis in conifers of cold climates. *Global Ecology and Biogeography*, 17, 696–707.
- Rudolph, T.D. (1964). Lammas Growth and Prolepsis in Jack Pine in the Lake States. *Forest Science*, 10, a0001–70.
- Sakai, A. & Larcher, W. (1987). *Freezing Injuries in Plants*, Springer Berlin Heidelberg, Berlin, Heidelberg, vol. 62, pp. 39–58.
- Schenker, G., Lenz, A., Körner, C. & Hoch, G. (2014). Physiological minimum temperatures for root growth in seven common European broad-leaved tree species. *Tree Physiology*, 34, 302–313.
- Silvestro, R., Deslauriers, A., Prislan, P., Rademacher, T., Rezaie, N., Richardson, A.D., Vitasse, Y. & Rossi, S. (2025). From Roots to Leaves: Tree Growth Phenology in Forest Ecosystems. *Current Forestry Reports*, 11, 12.
- Silvestro, R., Zeng, Q., Buttò, V., Sylvain, J.D., Drolet, G., Mencuccini, M., Thiffault, N., Yuan, S. & Rossi, S. (2023). A longer wood growing season does not lead to higher carbon sequestration. *Scientific Reports*, 13, 4059.
- Smith, S.A. & Brown, J.W. (2018). Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany*, 105, 302–314.
- Soolanayakanahally, R.Y., Guy, R.D., Street, N.R., Robinson, K.M., Silim, S.N., Albrechtsen, B.R. & Jansson, S. (2015). Comparative physiology of allopatric *Populus* species: Geographic clines in

- 497 photosynthesis, height growth, and carbon isotope discrimination in common gardens. *Frontiers in*
498 *plant science*, 6, 528.
- 499 Stearns, S.C. (1989). Trade-Offs in Life-History Evolution. *Functional Ecology*, 3, 259.
- 500 Stearns, S.C. (1998). *The Evolution Of Life Histories*. Oxford University Press.
- 501 Stevens, M.T. & Lindroth, R.L. (2005). Induced resistance in the indeterminate growth of aspen
502 (*Populus tremuloides*). *Oecologia*, 145, 297–305.
- 503 Valdovinos-Ayala, J., Robles, C., Fickle, J.C., Pérez-de-Lis, G., Pratt, R.B. & Jacobsen, A.L. (2022).
504 Seasonal patterns of increases in stem girth, vessel development, and hydraulic function in deciduous
505 tree species. *Annals of Botany*, 130, 355–365.
- 506 Vitasse, Y., Baumgarten, F., Zohner, C.M., Rutishauser, T., Pietragalla, B., Gehrig, R., Dai, J., Wang,
507 H., Aono, Y. & Sparks, T.H. (2022). The great acceleration of plant phenological shifts. *Nature*
508 *Climate Change*, 12, 300–302.
- 509 Volaire, F., Barkaoui, K., Grémillet, D., Charrier, G., Dangles, O., Lamarque, L.J., Martin-StPaul,
510 N. & Chuine, I. (2023). Is a seasonally reduced growth potential a convergent strategy to survive
511 drought and frost in plants? *Annals of Botany*, 131, 245–254.
- 512 Wang, Q., Liu, W., Leung, C.C., Tarté, D.A. & Gendron, J.M. (2024). Plants distinguish different
513 photoperiods to independently control seasonal flowering and growth. *Science*, 383, eadg9196.
- 514 Wolkovich, E.M., Ettinger, A.K., Chin, A.R., Chamberlain, C.J., Baumgarten, F., Pradhan, K., Man-
515 zanedo, R.D. & Hille Ris Lambers, J. (2025). Why longer seasons with climate change may not
516 increase tree growth. *Nature Climate Change*, 15, 1283–1292.
- 517 Zani, D., Crowther, T.W., Mo, L., Renner, S.S. & Zohner, C.M. (2020). Increased growing-season
518 productivity drives earlier autumn leaf senescence in temperate trees. *Science*, 370, 1066–1071.
- 519 Zhang, Y., Wu, H. & Yang, W. (2019). Forests Growth Monitoring Based on Tree Canopy 3D Recon-
520 struction Using UAV Aerial Photogrammetry. *Forests*, 10, 1052.
- 521 Zohner, C.M., Mirzagholi, L., Renner, S.S., Mo, L., Rebindaine, D., Bucher, R., Palouš, D., Vitasse,
522 Y., Fu, Y.H., Stocker, B.D. & Crowther, T.W. (2023). Effect of climate warming on the timing of
523 autumn leaf senescence reverses after the summer solstice. *Science*, 381, eadf5098.

- 524 Zohner, C.M., Mo, L., Pugh, T.A.M., Bastin, J.F. & Crowther, T.W. (2020). Interactive climate
525 factors restrict future increases in spring productivity of temperate and boreal trees. *Global Change*
526 *Biology*, 26, 4042–4055.
- 527 Zohner, C.M., Renner, S.S., Sebald, V. & Crowther, T.W. (2021). How changes in spring and autumn
528 phenology translate into growth-experimental evidence of asymmetric effects. *Journal of Ecology*,
529 109, 2717–2728.

Box 1: Metrics and proxies related to (in)determinacy

To advance our understanding of growth strategies under climate change, we need not only improved methods for quantifying the degree of (in)determinacy—but also empirical approaches that move beyond a simple binary classification. An organ-level metric at the shoot apical meristem captures the defining features of (in)determinacy and allows it to be explicitly defined and quantified. Temporal dynamics of shoot growth provide a related but indirect proxy that can be applied from individual shoots to crowns and entire forest stands.

Ratio of preformed to neoformed organs: The ratio of the number of leaves (or leaf scars) produced during a growing season to the number of leaf primordia preformed in dormant buds provides a direct, mechanistic, and quantitative measure of shoot-level (in)determinacy. Values close to one indicate growth constrained to preformed organs, whereas values greater than one reflect the production of neoformed organs and thus indeterminate development. Preformation and neoformation have been documented for more than a century (Moore, 1909), and several studies have quantified the number of preformed primordia and compared it with the final number of emerged leaves or phytomers (Damascos *et al.*, 2005; Kikuzawa, 1983; Guédon *et al.*, 2006). Quantification typically relies on bud dissection, but emerging non-destructive approaches such as micro-CT imaging offer promising alternatives for assessing bud structure and preformation (Kovaleski *et al.*, 2019). This metric remains the most conceptually robust approach for defining and quantifying the level of (in)determinacy across species.

Seasonal growth dynamics of the shoot apical meristem: Temporal patterns of shoot elongation provide a complementary perspective on (in)determinacy at the shoot apical meristem. Across species, elongation may occur over a brief period, extend continuously through the growing season, or appear in multiple distinct growth phases, reflecting differences in how apical meristem activity is maintained or reactivated. Such dynamics can be quantified using repeated shoot measurements or phenological observations, and increasingly through high-resolution terrestrial LiDAR and photogrammetric techniques that capture shoot elongation and structural change at the scale of entire crowns or stands (Zhang *et al.*, 2019; Jin *et al.*, 2022). While temporal growth dynamics do not distinguish preformed from neoformed organs directly, they remain closely linked to shoot apical meristem activity and provide a scalable proxy for the seasonal expression of (in)determinacy. Repeated measurements also allow shoot-elongation curves to be reconstructed, whose shape differs systematically among species (single, short sigmoid vs. prolonged or multi-phase extension; Hover *et al.* (2017); Kuster *et al.* (2014)). Specifically, curve-derived parameters such as the duration of shoot expansion relative to the growing season provide a quantitative proxy for the degree of (in)determinacy.




Figure 1: Schematic overview of the possible discrepancy between the potential growing season and the effectively realized vegetative growth (shoot and possibly cambial growth). When environmental factors, such as temperature and soil moisture, exceed growth-promoting thresholds they can act as filters that narrow the window of opportunity available for vegetative growth (potential light and photoperiodic constraints are not shown). The species-specific life history cycle (phenology) can impose another filter by dictating a dormancy cycle and prioritizing developmental processes other than vegetative growth (e.g. flowering, fruit maturation and storage). The degree of (in)determinacy is presented here in two extremes, although we propose to consider this trait as continuous.




Figure 2: Determinate and indeterminate growth within one growing season for species producing terminal buds. Commonly all tree species deploy buds during their first (spring) flush from prebuilt and overwintering leaf primordia (LP) protected by bud scales (BS; A). Determinate growing species set buds (B) that are under hormonal suppression to inhibit any further activity of the shoot apical meristem – SAM (paradormancy). The basic unit of a shoot is the phytomer which is composed of a node, a leaf, the axillary bud and an internode (see bottom right). Indeterminate growing species continue to produce new phytomers directly (C) or through one (D) to several (E) intermediate bud stage(s). Finally, most species set their terminal buds (F) and enter full dormancy (endodormancy).




Figure 3: Hypothesized predictions of growth rates for the three major meristem (apical, cambium and root) classes of trees under current and warmer climates following an extreme determinate (A) and indeterminate (B) growth strategy. The area under the curve is summarized as yearly biomass increment in the respective bar-plot. Arrows indicate the shift of growth phenology under warmer climate conditions. Root meristems appear to be purely temperature opportunistic for both strategies, even growing during warm winter spells. The indicated genera were observed to showcase the illustrated trends. The responses of these two contrasting growth strategies might apply not only to different tree species but also within a population (e.g. along environmental gradients) and even within an individual as it transitions from the juvenile to the adult stage (ontogeny).




Figure 4: Phylogenetic tree of plant species colored by growth determinacy trait values, visually representing the evolutionary distribution of growth strategies across taxa. This tree includes species for which shoot growth determinacy data were compiled (mainly from Hallé *et al.* (1978) and Kikuzawa (1983)), with the phylogeny pruned from the comprehensive seed plant tree of Smith & Brown (2018) to match the trait dataset. Growth determinacy was scored as indeterminate (blue), intermediate (yellow) and determinate (red), with mean values calculated per species when multiple entries existed. These values were then mapped as a continuous trait onto the phylogeny using the `contMap()` function from the `phytools` R package.