



Stockholm  
University

Department of Meteorology

## Ekman Spirals with Variable Eddy Viscosity

June 4, 2026

• Plota  $u(z)$ ,  $v(z)$  och  $\theta(z)$ ,  $|\vec{u}|(z)$  för  
1,5 sticht och den exponentiella profilen

*Author:* Fredrik Bergelv ([fredrik.bergelv@live.se](mailto:fredrik.bergelv@live.se))

*Supervisor:* Jonas Nycander

## Contents

|          |                                                             |           |
|----------|-------------------------------------------------------------|-----------|
| <b>1</b> | <b>Background</b>                                           | <b>2</b>  |
| 1.1      | Introduction . . . . .                                      | 2         |
| <b>2</b> | <b>Theory</b>                                               | <b>2</b>  |
| 2.1      | Problem formulation . . . . .                               | 2         |
| 2.2      | The classical solution . . . . .                            | 3         |
| 2.3      | The two layer model . . . . .                               | 3         |
| 2.4      | The exponential decaying viscosity . . . . .                | 4         |
| 2.5      | The 1.5 layer model . . . . .                               | 4         |
| 2.5.1    | Simple viscosity profiles for the 1.5 layer model . . . . . | 5         |
| 2.6      | The Ekman transport . . . . .                               | 6         |
| <b>3</b> | <b>Methodology</b>                                          | <b>6</b>  |
| 3.1      | Numerical method . . . . .                                  | 6         |
| <b>4</b> | <b>Results</b>                                              | <b>7</b>  |
| 4.1      | The theoretical models . . . . .                            | 7         |
| 4.2      | The numerical 1.5 model . . . . .                           | 8         |
| 4.2.1    | The Ekman transport . . . . .                               | 9         |
| <b>5</b> | <b>Discussion</b>                                           | <b>10</b> |
| <b>6</b> | <b>Conclusion</b>                                           | <b>10</b> |
| <b>A</b> | <b>The Source Code</b>                                      | <b>11</b> |

# 1 Background

(A) Discuss turbulent viscosity

## 1.1 Introduction

The Ekman spiral is a famous consequence of the Coriolis force in meteorology and oceanography. Due to the interaction between shear forces in the atmosphere and ocean, the wind high up in the atmosphere, and current in the deep ocean, will not give rise to a flow in the same direction at the surface. The adjustment of the flow from the surface will thus spiral, and is therefore properly named the Ekman spiral after its discoverer Ekman [1905]. The surface flow is often at an angle from the flow aloft (or down under in the ocean) by an angle of more or less  $45^\circ$  [Ekman, 1905]. Due to this turning of the wind, the net flux of the wind will be perpendicular to the wind aloft (or current below). This is known as the Ekman transport and is crucial for studying both meteorology and oceanography.

However, this derivation with a surface angle of  $45^\circ$  assumes a constant viscosity in the medium. ~~However,~~ this is a simplification of reality, where a turbulent boundary layer is present. Thus, in reality the viscosity should vary with altitude. This would then vary the surface angle from  $45^\circ$ , and thus also the Ekman transport. In this article, a varying viscosity with altitude will be studied under the Ekman equation. This will be done both theoretically and numerically and will also be compared to model data. This will give a deeper insight into the Ekman spiral and provide insight into when a surface angle different from  $45^\circ$  can be observed.

## 2 Theory

### 2.1 Problem formulation

The Ekman equation is derived from the steady-state Navier-Stokes equations on a rotating frame with shear flow. Assuming friction, incompressible flow, negligible vertical motion, gives the horizontal momentum equations as

$$\begin{cases} -fv = -\frac{1}{\rho} \frac{\partial p}{\partial x} + \frac{\partial}{\partial z} \left( \nu \frac{\partial u}{\partial z} \right), \\ fu = -\frac{1}{\rho} \frac{\partial p}{\partial y} + \frac{\partial}{\partial z} \left( \nu \frac{\partial v}{\partial z} \right), \end{cases} \quad (1a)$$

(1b)

where  $u$  and  $v$  are the horizontal velocity components,  $f$  is the Coriolis parameter,  $\rho$  is the fluid density, and  $\nu$  is the viscosity. Far away from the surface, shear flow and friction becomes negligible and the flow will thus approach geostrophic balance,  $(u, v) \rightarrow (u_g, v_g)$ , as were one can neglect the friction term in Equation 1b. Subtracting the geostrophic balance from Equation 1b thus gives

$$\frac{\partial}{\partial z} \rightarrow \frac{\partial}{\partial z} \text{ (d into kurst)} \begin{cases} -f(v - v_g) = \frac{\partial}{\partial z} \left( \nu \frac{\partial u}{\partial z} \right), \\ f(u - u_g) = \frac{\partial}{\partial z} \left( \nu \frac{\partial v}{\partial z} \right). \end{cases} \quad (2a)$$

(2b)

This problem can be formulated with one equation if one introduces a complex velocity defined as

$$U \equiv u + iv, \quad (3)$$

and a complex geostrophic velocity as  $U_g = u_g + iv_g$ . Using Equation 2b with the complex velocities  $U$  and  $U_g$  the coupled equations can be combined into the Ekman equation

$$\frac{\partial}{\partial z} \left( \nu \frac{\partial U}{\partial z} \right) = if(U - U_g). \quad (4)$$

With the boundary conditions

$$U(0) = 0, \quad (5a)$$

$$U(z \rightarrow \infty) = U_g. \quad (5b)$$

*Ge uttrycket för vindprofilen och Ekman-transport.*

## 2.2 The classical solution

The classical solution to the Ekman equation is obtained when the viscosity is assumed to be constant with altitude,  $\nu = \text{constant}$ , and can be found in Ekman [1905]. In this case the classical Ekman spiral solution from Equation 4 can be obtained as

$$U(z) = U_g \left( 1 - e^{-\frac{1+i}{\sqrt{2}} \frac{z}{h_{\text{Ek}}}} \right), \quad (6)$$

where

$$h_{\text{Ek}} \equiv \sqrt{\frac{\nu}{f}} \quad \text{varför inte } \sqrt{\frac{2\nu}{f}}? \quad (7)$$

*skriv om*

is the Ekman depth scale. One can note that by taking the argument of the change by altitude  $\frac{\partial U}{\partial z}|_{z=0}$ , one can measure the angle of Equation 6 as  $45^\circ$ .

## 2.3 The two layer model

The next natural step from a constant viscosity profile is to assume two layers of constant viscosity in each layer. One can label the lower level as having the viscosity  $\nu_1$  up until a height of  $H$ , where the next viscosity of  $\nu_2$  takes over. From this assumption one would obtain a superimposed solution similar to Equation 6 as

$$U_1(z) = A_1 e^{\frac{1+i}{\sqrt{2}} \frac{z}{h_{\text{Ek1}}}} + B_1 e^{-\frac{1+i}{\sqrt{2}} \frac{z}{h_{\text{Ek1}}}} + U_g, \quad (8a)$$

$$U_2(z) = B_2 e^{-\frac{1+i}{\sqrt{2}} \frac{z}{h_{\text{Ek2}}}} + U_g \quad (8b)$$

where the Ekman depth scales are defined for each layer like in Equation 7. To solve for the constants above one has to use the fact that the material flux has to be constant. This gives the jump condition as

*the velocity and*

$$U_1 = U_2 \quad \text{at } z = H, \quad (9a)$$

$$\nu_1 \frac{\partial U_1}{\partial z} = \nu_2 \frac{\partial U_2}{\partial z} \quad \text{at } z = H. \quad (9b)$$

Using this and the boundary conditions at Equation 5 one obtains an expression between the factors as

$$A_1 = -U_g \frac{1 - \gamma}{1 - \gamma + e^{\sqrt{2}(1+i)\varphi} (1 + \gamma)}, \quad \text{where } \gamma = \sqrt{\frac{\nu_2}{\nu_1}} \quad (10a)$$

$$B_1 = -U_g \frac{(1 + \gamma) e^{\sqrt{2}(1+i)\varphi}}{1 - \gamma + e^{\sqrt{2}(1+i)\varphi} (1 + \gamma)}, \quad (10b)$$

$$B_2 = -U_g \frac{2e^{\frac{1+i}{\sqrt{2}} \varphi} \left( 1 + \sqrt{\frac{\nu_1}{\nu_2}} \right)}{1 - \gamma + e^{\sqrt{2}(1+i)\varphi} (1 + \gamma)}, \quad (10c)$$

where  $\varphi$  is the non-dimensional height defined as

$$\varphi = \frac{H}{h_{\text{Ek1}}}. \quad (11)$$

The constant  $R$  is defined as

$$R = \frac{1 - \gamma}{1 + \gamma}. \quad (12)$$

Using this one can <sup>obtain</sup> define the surface angle as

$$\theta = \arg \frac{\partial U_1}{\partial z} \Big|_{z=0} = 45^\circ + \arctan \left[ \frac{2Re^{\sqrt{2}\varphi} \sin(\sqrt{2}\varphi)}{e^{2\sqrt{2}\varphi} - R^2} \right] \quad (13)$$

Observing Equation 13, and assuming that  $\nu_1 > \nu_2$ , one can see that the solution will always provide a result greater than  $45^\circ$ , except between the  $\varphi \in \left(\frac{\pi}{\sqrt{2}}, \sqrt{2}\pi\right)$  where an angle below  $45^\circ$  can be observed.

## 2.4 The exponential decaying viscosity

Another viscosity profile that can provide a fully analytical solution is an exponentially decaying viscosity with altitude. Here the viscosity is set to decay from an initial viscosity of  $\nu_0$ , with a decay rate of  $k$  as  $\nu(z) = \nu_0 \exp(-kz)$ . To solve Equation 4 with this viscosity profile one can introduce two new parameters

$$t = 2\sqrt{i}\varphi e^{\frac{kz}{2}}, \quad U^* = \frac{U - U_g}{t}. \quad (14)$$

The nondimensional parameter  $\varphi$  was introduced above in this problem, just like in the 2-layer model, but here it is defined as

$$\varphi = \frac{1}{k h_{EK0}} = \frac{1}{k} \sqrt{\frac{f}{\nu_0}}. \quad (15)$$

Using these newly defined variables Equation 4 transforms into

$$t^2 \frac{\partial^2 U^*}{\partial t^2} - t \frac{\partial U^*}{\partial t} - t^2 U^* = 0, \quad (16)$$

which is a differential equation with a standard solution. The standard solution is a modified Bessel equation as

$$U^* = C_1 I_1(t) + C_2 K_1(t), \quad (17)$$

where  $C_1$  and  $C_2$  are constants, while  $I_1(t)$  and  $K_1(t)$  are well-defined modified Bessel functions of the first and second kind of order one, respectively. By applying the previously defined boundary conditions from Equation 5 one can find the two constants. Applying the geostrophic boundary condition from Equation 5b one can see that  $C_1 = 0$  since  $I_1(t)$  does not go to zero when  $t \rightarrow \infty$  (which occurs when  $z \rightarrow \infty$ ). Applying the surface boundary condition from Equation 5a one can find an expression for the second constant. Transforming the solution to the normal complex velocity as in Equation 3 one can obtain the solution

$$U(z) = U_g \left[ 1 - \frac{e^{\frac{kz}{2}} K_1(2\sqrt{i}\varphi e^{\frac{kz}{2}})}{K_1(2\sqrt{i}\varphi)} \right]. \quad \sqrt{i} = \frac{1+i}{\sqrt{2}} \quad (18)$$

Using Equation 18 one can derive the surface angle of this model as

$$\theta = \arg \frac{\partial U}{\partial z} \Big|_{z=0} = \arg \left[ U_g \sqrt{i} \frac{K_0(2\sqrt{i}\varphi)}{K_1(2\sqrt{i}\varphi)} \right], \quad (19)$$

where  $K_0$  is the modified Bessel function of the second kind of order zero.

## 2.5 The 1.5 layer model

In the previous sections two completely analytical solutions for two different altitude-varying viscosity profiles were derived. The difficulty in solving Equation 4 comes from the altitude dependence of the viscosity profile  $\nu(z)$ . This introduces variable coefficients into the complex second-order ordinary differential equation, which in general prevents closed-form analytical solutions. Analytical solutions therefore exist only for specific viscosity profiles that allow the equation to be transformed into a solvable form.

Equation 4 is furthermore also difficult to solve in a numerical setting, as the upper boundary condition is a limit. This will put limits on the viscosity profiles one can study, as the viscosity cannot increase

$\varphi$  is analogous, but not the same. Give it a new name.

the flow is laminar and the turbulent viscosity small.

with altitude continuously. In the physical world, this makes sense, since we assume that high up in the atmosphere (or deep down in the ocean), friction and shear stress become negligible. This was what was labelled as the geostrophic limit. However, by limiting ourselves to only upper-level decreasing viscosity profiles, the differential equation becomes more difficult to solve. Furthermore, if one wants to solve for a dimensionless height parameter, too complex viscosity profiles become hard to deal with.

A solution to this problem is to introduce the 1.5 layer model, which is like the 2-layer model, except that we assume that the upper-level viscosity is negligible  $\nu_2 \approx 0$ . This also makes physical sense, since we know that both the atmosphere and the ocean consist of a boundary layer where turbulent mixing takes place. Reducing the problem to a 1.5 model will thus make it possible to test viscosity profiles that do not converge at  $z \rightarrow \infty$ , since the new domain is between  $z \in [0, H]$ . This modifies the boundary conditions to Equation 4 as

$$U(0) = 0, \quad (20a)$$

$$\left. \frac{\partial U}{\partial z} \right|_{z=H} = 0. \quad (20b)$$

Above,  $H$  becomes the height of the boundary layer, and the upper boundary condition was argued from the jump condition in the two-layer model, as seen in Equation 9b.

### 2.5.1 Simple viscosity profiles for the 1.5 layer model

Using the 1.5 layer model one can study some simple viscosity profiles to be evaluated numerically. The easiest, except for the constant viscosity, is a linearly varying viscosity with height as

$$\nu_{\uparrow}(z) = \frac{\nu_{\max}}{H} z, \quad 0 < z < H \quad (21a)$$

$$\nu_{\downarrow}(z) = \frac{\nu_{\max}}{H} (H - z), \quad (21b)$$

where  $\nu_{\max}$  is the maximal viscosity. For both of these profiles one can define a dimensionless height coordinate as

$$\tilde{z} = \frac{z}{H} \Rightarrow \begin{cases} \nu_{\uparrow}(z) = \nu_{\max} \tilde{z} \\ \nu_{\downarrow}(z) = \nu_{\max} (1 - \tilde{z}) \end{cases} \quad 0 < \tilde{z} < 1 \quad (22)$$

One must note that this new coordinate changes the previous  $z$ -derivative, as a factor of  $1/H$  appears when taking the derivative with respect to  $\tilde{z}$ . Using these linear viscosity profiles, together with the new dimensionless height coordinate, one can rewrite Equation 4 for the two profiles as

$$\frac{\partial}{\partial \tilde{z}} \left( \underbrace{\tilde{z}}_{\equiv \tilde{\nu}_{\uparrow}} \frac{\partial U}{\partial \tilde{z}} \right) = i \underbrace{\frac{H f^2}{\nu_{\max}}}_{\equiv \varphi^2} (U - U_g), \quad (23a)$$

$$\frac{\partial}{\partial \tilde{z}} \left( \underbrace{[1 - \tilde{z}]}_{\equiv \tilde{\nu}_{\downarrow}} \frac{\partial U}{\partial \tilde{z}} \right) = i \underbrace{\frac{H f^2}{\nu_{\max}}}_{\equiv \varphi^2} (U - U_g). \quad (23b)$$

Above the dimensionless height  $\varphi$  was again introduced following the logic from before.

Another simple profile to evaluate is a parabolic increasing viscosity as

$$\nu_p = \frac{4\nu_{\max}}{H^2} (H - z) z. \quad (24)$$

Again, using the non-dimensional height coordinate  $\tilde{z}$  and the dimensionless height  $\varphi$ , one can rewrite Equation 4 as

$$\frac{\partial}{\partial \tilde{z}} \left( \underbrace{4\tilde{z}[1 - \tilde{z}]}_{\equiv \tilde{\nu}_p} \frac{\partial U}{\partial \tilde{z}} \right) = i \underbrace{\frac{H f^2}{\nu_{\max}}}_{\equiv \varphi^2} (U - U_g). \quad (25)$$

(A) 1,5 skikt: konstant  $\nu$ , viskositet, och överensstämmande med 2 skikt.

## 2.6 The Ekman transport

For a theoretically or numerically derived solution, one can derive the Ekman transport, which may be a more useful quantity. The Ekman transport is defined as the mass flux per meter for the entire Ekman spiral. In the classical solution the Ekman transport is  $90^\circ$  to the right of the surface flow. However one must note that this angle is obtained by subtracting the Ekman transport relative to the geostrophic flow or the surface flow, i.e.  $135^\circ - 45^\circ = 90^\circ$  in classical theory. However, for an obtained solution  $U(\tilde{z}, \varphi)$  as a function of the dimensionless height and layer thickness, one can obtain the mass flux as

$$\Phi_T(\varphi) = \int_0^1 [U(\tilde{z}, \varphi) - U_g] d\tilde{z}. \quad (26)$$

To obtain the transport angle one can simply compute the argument as

$$\theta_T(\varphi) = \arg[\Phi_T(\varphi)]. \quad (27)$$

Thus the relative angle between the surface transport and the upper geostrophic flow can be calculated as

$$\theta_R = \theta_T - \theta. \quad (28)$$

## 3 Methodology

### 3.1 Numerical method

For arbitrary viscosity profiles, Equation 4 generally does not admit a closed-form analytical solution. The equation was therefore solved numerically as a boundary value problem using the `solve_bvp` routine available in SciPy [Virtanen et al., 2020]. However, it was previously shown that the equation could be rewritten for a dimensionless layer thickness  $\varphi$ . Thus, using an arbitrary height-normalized height-dependent eddy viscosity profile,  $\tilde{\nu}(z)$ , one can write a simpler differential equation. Separating the equation into the zonal and meridional velocity components, one obtains

$$\frac{\partial}{\partial z} \left( \tilde{\nu}(z) \frac{\partial u}{\partial z} \right) = -\varphi^2 v, \quad (29a)$$

$$\frac{\partial}{\partial z} \left( \tilde{\nu}(z) \frac{\partial v}{\partial z} \right) = \varphi^2 (u - u_g), \quad (29b)$$

where the wind aloft was set as a purely zonal wind  $u_g$ . To apply the numerical boundary value solver, the equations were rewritten as a first-order system by introducing the new variables

$$u' = \frac{\partial u}{\partial z}, \quad v' = \frac{\partial v}{\partial z}. \quad \text{och } \tilde{\nu}' = \frac{d\tilde{\nu}}{dz} \quad (30)$$

Thus one can expand Equation 29b into four coupled first-order equations

$$\begin{cases} \frac{\partial u}{\partial z} &= u', \\ \frac{\partial u'}{\partial z} &= \frac{-\varphi^2 v - \tilde{\nu}' u'}{\tilde{\nu}}, \\ \frac{\partial v}{\partial z} &= v', \\ \frac{\partial v'}{\partial z} &= \frac{\varphi^2 (u - u_g) - \tilde{\nu}' v'}{\tilde{\nu}}, \end{cases} \quad (31)$$

with the boundary conditions

$$u(\varphi = 0) = 0, \quad u'(\varphi = 1) = 0, \quad (32a)$$

$$v(\varphi = 0) = 0, \quad v'(\varphi = 1) = 0. \quad (32b)$$

The equations were solved iteratively on a vertical grid. The solver required an initial guess, which was chosen based on the initial setup of the system. The initial guess was therefore set as the classical solution for a constant viscosity. The final solution was found when the iterative solution reached a steady state for the differential equations with the imposed boundary conditions.


(A) Definiera Ekman transporten som komponenten ortogonalt mot  $\tilde{u}_g$ , dvs parallell med  $\nabla p$ .

(B) Vi läser bara fall med förenklat övre randvillkor

## 4 Results


### 4.1 The ~~theoretical~~ <sup>analytical</sup> models

The result from the theoretical two layer model, Equation 13, can be seen in Figure 1. There one can observe solutions for different viscosity ratios, half of which have an higher upper-level viscosity and the other half a lower one. These are plotted together with a 45° reference point to highlight the difference from the classical Ekman solution. One can observe a convergence towards the classical Ekman value when  $\varphi \rightarrow \infty$  for all the viscosity ratios. The zero limit is, however, dependent on the viscosity ratio, where finite ratios approach 45°. However, when  $\frac{\nu_2}{\nu_1} \rightarrow 0$  and  $\varphi \rightarrow 0$  one can observe that  $\theta \rightarrow 90^\circ$ . Respectively, one can observe that when  $\frac{\nu_2}{\nu_1} \rightarrow \infty$  and  $\varphi \rightarrow 0$ ,  $\theta \rightarrow 0^\circ$ .



**Figure 1:** The surface angle as a function of the dimensionless layer thickness  $\varphi = H/h_{EK1}$  for the two-layer viscosity model. Different coloured curves correspond to different layer viscosity ratios. The classical Ekman angle of 45° is shown as a reference.

Figure 2 showcases the result from the exponentially decaying viscosity as in Equation 19. There one can observe a similar limit as in Figure 1 when  $\varphi \rightarrow 0$ , as Equation 19 also approaches 90°. The same limit is also true for  $\varphi \rightarrow \infty$ , which gives  $\theta \rightarrow 45^\circ$ , although this exponential solution converges more slowly than the two-layer solution. However, the shape is much more uniform, with an exponential-like shape. However, all angles are here greater then the reference angle of 45°.



**Figure 2:** Surface angle for the exponentially decaying viscosity model as a function of the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.

Förklara kvalitativt att  $\theta \rightarrow 45^\circ$  både då  $\varphi \rightarrow 0$  (spiralen helt övre skiktet) och  $\varphi \rightarrow \infty$  (spiralen helt i undre skiktet).


slutats:  $\theta > 45^\circ$  när det mesta av spiralen ligger i ett övre skikt med litet  $\nu$ .



## 4.2 The numerical 1.5 model


The result from the numerical 1.5-layer model can be viewed in Figure 3–4. There one can observe similarities between these models, the two-layer model, and the exponentially decaying model. They all have a lower boundary of  $90^\circ$  (except for the decreasing two-layer model, where the lower boundary was  $0^\circ$ ). This strengthens the results from these numerical models, as the theoretical ones behave in a similar manner.

The result from the linear 1.5-layer model is shown in Figure 3. There one can observe slightly different behaviour for the linearly increasing and decreasing viscosities. For the linearly decreasing viscosity, the values go smoothly from  $90^\circ \rightarrow 45^\circ$ . However, for the linearly increasing viscosity, the upper boundary is instead located at  $0^\circ$ , and the curve is approximately  $45^\circ$  at  $\varphi \approx 0.210$ . One must note that both of these upper limits only occur when  $\varphi \rightarrow \infty$ .



**Figure 3:** Numerically computed surface angles for linearly increasing and decreasing viscosity profiles in the 1.5-layer model. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.

The 1.5-layer model with a parabolic viscosity profile gave Figure 4. There one can observe very similar behaviour to the increasing viscosity profile in Figure 3, with the main difference being that it crosses the reference line of  $45^\circ$  at  $\varphi \approx 0.400$ .




**Figure 4:** Numerically computed surface angles for a parabolic viscosity profile in the 1.5-layer model. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.


#### 4.2.1 The Ekman transport

In Figure 5–8 one can observe the angle of the integrated transport for both the theoretical two-layer model, the exponentially decaying model, and the numerical 1.5-layer models. One can see identical behaviour in the plots of Figure 5–8 compared to the surface wind angles in Figure 1–4, with the only difference being that these have a new reference point at  $135^\circ$ . Thus, the plots are merely translated to this new reference point, since their respective shapes remain intact.

Thus one can make the interesting observation that if one computes the relative transport angle, i.e. the angle between the surface angle and the transport angle, one obtains a relative value of  $90^\circ$  for all the different models. Thus, these models give the same Ekman transport angle as in the classical Ekman theory.




**Figure 5:** Numerically computed transport angles for the two layer viscosity profile. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . Different coloured curves correspond to different layer viscosity ratios. The solution assumed a purely zonal geostrophic flow.




**Figure 6:** Numerically computed transport angles for the exponential viscosity profile. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.

(A) Detta kan visas analytiskt. Titta på transporten längs  $\nabla p$ .



**Figure 7:** Numerically computed transport angles for linearly increasing and decreasing viscosity profiles in the 1.5-layer model. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.



**Figure 8:** Numerically computed transport angles for a parabolic viscosity profile in the 1.5-layer model. The angle is dependent on the dimensionless parameter  $\varphi = H/h_{EK}$ . The solution assumed a purely zonal geostrophic flow.

## 5 Discussion

## 6 Conclusion

## A The Source Code

The source code for this article can be found on GitHub under:

<https://github.com/FredrikBergelov/Ekman-Spiral-Project>

## References

- V. Walfrid Ekman. On the Influence of the Earth's Rotation on Ocean-Currents. *Arkiv för matematik, astronomi och fysik*, 2(11):1–52, May 1905.
- Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. Van Der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul Van Mulbregt, SciPy 1.0 Contributors, Aditya Vijaykumar, Alessandro Pietro Bardelli, Alex Rothberg, Andreas Hilboll, Andreas Kloeckner, Anthony Scopatz, Antony Lee, Ariel Rokem, C. Nathan Woods, Chad Fulton, Charles Masson, Christian Häggström, Clark Fitzgerald, David A. Nicholson, David R. Hagen, Dmitrii V. Pasechnik, Emanuele Olivetti, Eric Martin, Eric Wieser, Fabrice Silva, Felix Lenders, Florian Wilhelm, G. Young, Gavin A. Price, Gert-Ludwig Ingold, Gregory E. Allen, Gregory R. Lee, Hervé Audren, Irvin Probst, Jörg P. Dietrich, Jacob Silterra, James T Webber, Janko Slavič, Joel Nothman, Johannes Buchner, Johannes Kulick, Johannes L. Schönberger, José Vinícius De Miranda Cardoso, Joscha Reimer, Joseph Harrington, Juan Luis Cano Rodríguez, Juan Nunez-Iglesias, Justin Kuczynski, Kevin Tritz, Martin Thoma, Matthew Newville, Matthias Kümmerer, Maximilian Bolingbroke, Michael Tartre, Mikhail Pak, Nathaniel J. Smith, Nikolai Nowaczyk, Nikolay Shebanov, Oleksandr Pavlyk, Per A. Brodtkorb, Perry Lee, Robert T. McGibbon, Roman Feldbauer, Sam Lewis, Sam Tygier, Scott Sievert, Sebastiano Vigna, Stefan Peterson, Surhud More, Tadeusz Pudlik, Takuya Osima, Thomas J. Pingel, Thomas P. Robitaille, Thomas Spura, Thouis R. Jones, Tim Cera, Tim Leslie, Tiziano Zito, Tom Krauss, Utkarsh Upadhyay, Yaroslav O. Halchenko, and Yoshiki Vázquez-Baeza. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods*, 17(3):261–272, March 2020. ISSN 1548-7091, 1548-7105. doi: 10.1038/s41592-019-0686-2. URL <https://www.nature.com/articles/s41592-019-0686-2>.