



Stockholm
University

Department of Meteorology

Modelling of Large-Scale Circulation in Atmosphere and Ocean
(MO8004)

Wind Driven Ocean Gyres Sverdrup and Munk Solutions

Fredrik Bergelv

June 3, 2026

Introduction

Ocean circulation gyres are part of the atmospheric circulation.

- 1 Heating air at the equator → Hadley cells.
- 2 Coriolis effect deflects the air movement → Trade winds and westerlies
- 3 Enforce wind stress on the ocean surface → 90° Ekman transport.
- 4 Since f increases with latitude, net southward Ekman transport in the North Atlantic.
- 5 Western boundary current needed to close the circulation.



Figure: Wikimedia Commons, North Atlantic location map, accessed 2026

Introduction, Sverdrup

Introduction

Theory

Sverdrup

Munk

Method

Result

The Sverdrup solution

The Munk solution

Discussion

Conclusion

References

The Munk solution

Harald Ulrik Sverdrup (1888-1957) formulated the net transport in the ocean interior as:

$$\boxed{-\beta \frac{\partial \Psi}{\partial x} = \frac{1}{\rho} \nabla \times \vec{\tau}} \quad (1)$$

which gives the Sverdrup solution:

$$\begin{aligned} \int_x^{x_{\text{East}}} \frac{\partial \Psi}{\partial x} dx &= \int_x^{x_{\text{East}}} vH dx = \cancel{\Psi(x_{\text{East}}, y)} - \Psi(x, y) \\ \Rightarrow \Psi(x, y) &= -\frac{1}{\rho\beta} \int_x^{x_{\text{East}}} (\nabla \times \vec{\tau}) \end{aligned}$$

Introduction, Munk

Walter Heinrich Munk (1917-2019) extended the Sverdrup model by including meridional harmonic Newtonian viscosity (diffusion of momentum):

$$A \nabla^4 \Psi - \beta \frac{\partial \Psi}{\partial x} = \frac{1}{\rho} \nabla \times \vec{\tau} \quad (2)$$

which gives the Munk solution:

$$\Psi(x, y) = \Psi_{\text{Sverdrup}}(x, y) \left(1 - e^{-\frac{x - x_{\text{West}}}{2\delta_m}} \left[\cos\left(\sqrt{3} \frac{x - x_{\text{West}}}{2\delta_m}\right) + \frac{1}{\sqrt{3}} \sin\left(\sqrt{3} \frac{x - x_{\text{West}}}{2\delta_m}\right) \right] \right)$$

$$\text{where } \delta_M \equiv \left(\frac{A}{\beta} \right)^{\frac{1}{3}}$$

Stress set as:

$$\tau = \rho_{\text{air}} C_D U_{10}^2 \sin \left(\frac{\pi y}{L_y} \right). \quad (3)$$

<i>Variable</i>	<i>Sverdrup model</i>	<i>Munk model</i>
Atlantic, L_x, L_y	7000 km, 7000 km	7000 km, 7000 km
Pacific, L_x, L_y	–	14000 km, 7000 km
Grid, $\Delta x, \Delta y$	35 km	35 km
Wind forcing, U_{10}	[4, 5, 6] m/s	5 m/s
Viscosity, A	0	$[3 \cdot 10^4, 10^5, 5 \cdot 10^5] \text{ m}^2/\text{s}$
$\delta_M / \Delta x$	–	[3.3, 4.9, 8.4]

Table: Comparison of parameters used in the Sverdrup and Munk models.

Result, The Sverdrup solution




Figure: Streamfunction for the no-drag f -plane configuration. The circulation remains symmetric with no β -effect and without western intensification.

Result, The Sverdrup solution




Figure: Streamfunction for the no-drag β -plane configuration. The red lines are the theoretical Sverdrup solution.

Result, The Munk solution




Figure: Streamfunction for the Laplace-viscous β -plane configuration. The red lines are the theoretical Munk solution.

Result, The Munk solution




Figure: Streamfunction for the drag β -plane configuration with varying domain size. The red lines are the theoretical Sverdrup solution.

Result, The Munk solution




Figure: Streamfunction for the Laplace-viscous β -plane configuration. The red lines are the theoretical Sverdrup and Munk solutions respectively.

Discussion

Both numerical models show western boundary currents, but only Munk theory does. Why?

- The Sverdrup theory is linear, whereas the Munk theory is non-linear, and frictional force is opposite sign to the motion itself.
- The Munk viscosity creates a boundary layer where the frictional force can take care of mass conservation.
- Friction balances continuous input of energy. *Realistic?*
- Gives western boundary flow in right magnitude to real Gulf Stream ($\sim 2.5 \text{ m s}^{-1}$).

$$A \nabla^4 \Psi - \beta \frac{\partial \Psi}{\partial x} = \frac{1}{\rho} \nabla \times \vec{\tau}$$

Three important conclusions:

- 1 Both numerical models showed western intensification, explained by Rossby wave propagation. But continuous energy input need dissipation to converge with time.
- 2 Only Munk theory showed a western boundary current, explained by frictional layer, although both numerical models showed return flow.
- 3 A larger domain gave same interior meridional velocity, which resulted in a stronger western boundary current.

Result, The Munk solution




Figure: Streamfunction for the Laplace-viscous β -plane configuration. The red lines are the theoretical Munk solution. Twice as high resolution (i.e. $\Delta x = \Delta y = 17.5 \text{ km}$)