



UiO : **Department of Mathematics**  
University of Oslo

# Join of hexagons and Calabi–Yau threefolds

Public defence

**Fredrik Meyer**

**November 8, 2017**

# Outline of the thesis


- Unsuccessful attempt to find new hyper-Kähler varieties.
- The topology of  $C(dP_6)$ .
- New Calabi–Yau varieties and potential mirror partners.

# Calabi–Yau manifolds

## Definition (Calabi–Yau variety)

A Calabi–Yau variety is a smooth projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = k$  and  $h^1(X, \mathcal{O}_X) = h^2(\mathcal{O}_X) = 0$ .
  - The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .
- 
- Easiest invariant are the Euler characteristic and the Hodge numbers.
  - We always have  $\chi = 2(h^{11} - h^{12})$ .



# Calabi–Yau manifolds

## Definition (Calabi–Yau variety)

A Calabi–Yau variety is a smooth projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = k$  and  $h^1(X, \mathcal{O}_X) = h^2(\mathcal{O}_X) = 0$ .
  - The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .
- 
- Easiest invariant are the Euler characteristic and the Hodge numbers.
  - We always have  $\chi = 2(h^{11} - h^{12})$ .

$$\begin{array}{ccccccc}
 & & & & 1 & & \\
 & & & & 0 & & 0 \\
 & & 0 & & h^{11} & & 0 \\
 1 & & h^{12} & & & & h^{12} & 1 \\
 & 0 & & h^{11} & & 0 & & \\
 & & 0 & & 0 & & & \\
 & & & & 1 & & & 
 \end{array}$$

# Hodge numbers

- The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{11} = 1$  and  $h^{12} = 101$ .

## Remark (Heuristic)

The number  $h^{12}$  is the dimension of the “space of parameters” of  $X$ . The following heuristic will give us the correct Hodge number:

- The space of degree 5 polynomials  $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))$  in  $\mathbb{P}^4$  is  $\binom{4+5}{5} = \binom{9}{4} = 126$ -dimensional. Hence  $\mathbb{P}(H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))) = \mathbb{P}^{125}$ .
- This is not unique, but we can act by  $\mathrm{PGL}(5)$  to identify isomorphic quintics. We have  $\dim \mathrm{PGL}(5) = 25 - 1 = 24$ .
- In total:  $125 - 24 = 101$ , which is  $h^{12}(X)$ .

# Mirror symmetry

- Calabi–Yau threefolds seem to “always” have “mirror partners”.
- Mirror partner  $X^\circ$  to  $X$  have “mirrored Hodge diamond”.
- Hence  $\chi(X^\circ) = -\chi(X)$ .

$X$					
<hr/>					
		1			
	0		0		
0		1		0	
1	0	1	0	1	0
	0		1		0
	0		0		
		1			

# Mirror symmetry

- Calabi–Yau threefolds seem to “always” have “mirror partners”.
- Mirror partner  $X^\circ$  to  $X$  have “mirrored Hodge diamond”.
- Hence  $\chi(X^\circ) = -\chi(X)$ .

$X$				
		1		
	0		0	
0		1		0
1	0	101	101	0
	0		1	0
		0		0
			1	

$X^\circ$				
		1		
	0		0	
0		101		0
1	0	1	1	0
	0		101	0
		0		0
			1	

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ .

- 1 Suppose  $X$  has a natural degeneration  $X_0$  with a finite automorphism group  $G$ .
- 2 Find a family  $\pi : \mathcal{X} \rightarrow S$  on which  $G$  act, and such that the general fiber  $X_t$  have only isolated singularities.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .



# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ .

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 Find a family  $\pi : \mathcal{X} \rightarrow S$  on which  $G$  act, and such that the general fiber  $X_t$  have only isolated singularities.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ .

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ .

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There is an action of  $H \triangleq (\mathbb{Z}/5)^5/(\mathbb{Z}/5)$  on  $X_t$ . Crepant resolutions of  $X_t/H$  exists, and is a mirror.

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ .

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There is an action of  $H \triangleq (\mathbb{Z}/5)^5/(\mathbb{Z}/5)$  on  $X_t$ . Crepant resolutions of  $X_t/H$  exists, and is a mirror.

We can use *Roan's formula* to compute the Euler characteristic.

Theorem (Roan's formula)

$$\chi(\widetilde{X_t/H}) = \frac{1}{|H|} \sum_{g,h \in H} \chi(X_t^g \cap X_t^h).$$


# The cone over $dP_6$

- Let  $dP_6 \subset \mathbb{P}^6$  be an anticanonically embedded del Pezzo surface of degree 6. Let  $C(dP_6)$  be its affine cone in  $\mathbb{A}^7$ .
- The equations are

$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y & x_3 \\ x_5 & x_6 & y \end{vmatrix} \leq 1$$

The origin is an isolated singularity.

- There are two smoothing components.
- They come from perturbations of different form of writing the equation.
- Can also write the equations as:



$$\leq 1$$

# The two smoothing components

TODO: trengs denne?

# Construction of a new Calabi–Yau: $X_1$

- Let  $E$  be a vector space with basis  $e_1, e_2, e_3$ . Consider

$$\mathbb{P}^{17} = \mathbb{P}(E \otimes E \oplus E \otimes E).$$

The elements are pairs of  $3 \times 3$  matrices, equal up to scalar multiplication.

- Consider the set of pairs  $M$  of matrices  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $M$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{17}$ . Let  $X_1 \triangleq M \cap \mathbb{P}^{11}$ .

## Theorem

$X_1$  is a smooth Calabi–Yau with Euler-characteristic  $-72$ .

## Construction of a new Calabi–Yau: $X_2$

- Let  $F$  be a vector space with basis  $f_1, f_2$ . Consider

$$\mathbb{P}^{15} = \mathbb{P}(F \otimes F \otimes F \oplus F \otimes F \otimes F),$$

The elements are pairs of  $2 \times 2 \times 2$ -tensors, equal up to scalar multiplication.

- Consider the set of pairs  $N$  of tensors  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $N$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{15}$ . Let  $X_2 \triangleq N \cap \mathbb{P}^{11}$ .

### Theorem

$X_2$  is a smooth Calabi–Yau with Euler-characteristic  $-48$ .



# Construction of a new Calabi–Yau: $X_3$

- Let  $E$  and  $F$  be as before. Consider

$$\mathbb{P}^{16} = \mathbb{P}(E \otimes E \oplus F \otimes F \otimes).$$

- Consider the set of pairs  $W$  of tensors  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $W$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{16}$ . Let  $X_3 \triangleq W \cap \mathbb{P}^{11}$ .

## Theorem

*$X_3$  is a smooth Calabi–Yau with Euler-characteristic  $-60$ .*

# Hodge number heuristics

## Conjecture

$X_1$  have Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$  (= number of parameters).
- 2 The Grassmannian of  $\mathbb{P}^{11}$ 's in  $\mathbb{P}^{17}$  is 72-dimensional.
- 3 We can act by automorphism from  $\prod_{i=1}^4 \mathrm{GL}(E)$ .
- 4 The subgroup  $\{t_1 t_2 = t_3 t_4\} \subset (\mathbb{C}^*)^4 \subset \prod_{i=1}^4 \mathrm{GL}(E)$  acts trivially.
- 5 Hence  $h^{12} = 72 - (\dim \prod_{i=1}^4 \mathrm{GL}(E) - 3) = 72 - 33 = 39$ .

# Mirror candidates

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

■ BB

UiO : **Department of Mathematics**  
University of Oslo



**Fredrik Meyer**



**Join of hexagons and  
Calabi–Yau threefolds**  
Public defence

