



UiO : **Department of Mathematics**  
University of Oslo

# Join of hexagons and Calabi–Yau threefolds

Public defence

**Fredrik Meyer**

**November 8, 2017**

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**
- The topology of smoothings of  $C(dP_6)$ .

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**
- The topology of smoothings of  $C(dP_6)$ .
  - Identify the smoothings as hyperplane complements, and use exact sequences from algebraic topology.

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**
- The topology of smoothings of  $C(dP_6)$ .
  - Identify the smoothings as hyperplane complements, and use exact sequences from algebraic topology.
- New Calabi–Yau varieties and potential mirror partners.

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**
- The topology of smoothings of  $C(dP_6)$ .
  - Identify the smoothings as hyperplane complements, and use exact sequences from algebraic topology.
- New Calabi–Yau varieties and potential mirror partners.
  - Found by smoothing a certain Stanley–Reisner sphere.

# Outline of the thesis

- Attempts to find new hyper-Kähler varieties.
  - Strategy: Naïvely perturb monomial equations of Stanley–Reisner ideals of known triangulations of  $\mathbb{CP}^2$ . **Did not work.**
- The topology of smoothings of  $C(dP_6)$ .
  - Identify the smoothings as hyperplane complements, and use exact sequences from algebraic topology.
- New Calabi–Yau varieties and potential mirror partners.
  - Found by smoothing a certain Stanley–Reisner sphere.

I will now focus on the last point — the construction of the new Calabi–Yau’s.



# Stanley–Reisner schemes

- Given a simplicial complex  $\mathcal{F}$ , we get a Stanley–Reisner scheme  $\mathbb{P}(\mathcal{F})$ .

# Stanley–Reisner schemes



- Given a simplicial complex  $\mathcal{F}$ , we get a Stanley–Reisner scheme  $\mathbb{P}(\mathcal{F})$ .
- Is a union of projective spaces  $\mathbb{P}^{\dim f}$ , where  $f$  is a face of  $\mathcal{F}$ .

# Stanley–Reisner schemes

- Given a simplicial complex  $\mathcal{F}$ , we get a Stanley–Reisner scheme  $\mathbb{P}(\mathcal{F})$ .
- Is a union of projective spaces  $\mathbb{P}^{\dim f}$ , where  $f$  is a face of  $\mathcal{F}$ .

## Example

From a simplicial complex to a union of  $\mathbb{P}^1$ 's.



The ideal is generated by  $x_i x_{i+2} = x_i x_{i+3} = 0$  ( $i = 0, \dots, 5$ ).

# Stanley–Reisner schemes

- **Join** of two subschemes  $X$  and  $Y$ : The (closure of) the union of all lines between  $X$  and  $Y$ .

# Stanley–Reisner schemes

- **Join** of two subschemes  $X$  and  $Y$ : The (closure of) the union of all lines between  $X$  and  $Y$ .
- **Join** of two Stanley–Reisner schemes  $\mathbb{P}(\mathcal{F})$  and  $\mathbb{P}(\mathcal{G})$  is  $\mathbb{P}(\mathcal{F} * \mathcal{G})$ , where the faces of  $\mathcal{F} * \mathcal{G}$  are  $f \sqcup g$  for  $f \in \mathcal{F}$ ,  $g \in \mathcal{G}$ .

# Stanley–Reisner schemes

- **Join** of two subschemes  $X$  and  $Y$ : The (closure of) the union of all lines between  $X$  and  $Y$ .
- **Join** of two Stanley–Reisner schemes  $\mathbb{P}(\mathcal{F})$  and  $\mathbb{P}(\mathcal{G})$  is  $\mathbb{P}(\mathcal{F} * \mathcal{G})$ , where the faces of  $\mathcal{F} * \mathcal{G}$  are  $f \sqcup g$  for  $f \in \mathcal{F}$ ,  $g \in \mathcal{G}$ .

Smoothings of Stanley–Reisner schemes:

- Given a basis for  $T^1(S_{\mathbb{P}(\mathcal{K})}/k, S_{\mathbb{P}(\mathcal{K})}_0)$ , we can try to find a smoothing of  $X_0 = \mathbb{P}(\mathcal{K})$ .

# Stanley–Reisner schemes

- **Join** of two subschemes  $X$  and  $Y$ : The (closure of) the union of all lines between  $X$  and  $Y$ .
- **Join** of two Stanley–Reisner schemes  $\mathbb{P}(\mathcal{F})$  and  $\mathbb{P}(\mathcal{G})$  is  $\mathbb{P}(\mathcal{F} * \mathcal{G})$ , where the faces of  $\mathcal{F} * \mathcal{G}$  are  $f \sqcup g$  for  $f \in \mathcal{F}$ ,  $g \in \mathcal{G}$ .

Smoothings of Stanley–Reisner schemes:

- Given a basis for  $T^1(S_{\mathbb{P}(\mathcal{K})}/k, S_{\mathbb{P}(\mathcal{K})}_0)$ , we can try to find a smoothing of  $X_0 = \mathbb{P}(\mathcal{K})$ .
- A smoothing  $X$  of  $X_0$  will have many of the same properties:
  - The same Hilbert polynomial.
  - By semicontinuity, if  $X_0$  is a sphere,  $X$  will be Calabi–Yau.

# Calabi–Yau varieties

## Definition

A **Calabi–Yau variety** is an irreducible, smooth, projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = \mathbb{C}$  and  $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ .
- The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .



# Calabi–Yau varieties

## Definition

A **Calabi–Yau variety** is an irreducible, smooth, projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = \mathbb{C}$  and  $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ .
- The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .

- Easiest invariants are the **Euler characteristic** and the **Hodge numbers**,  $h^{ij} = h^j(X, \Omega_{X/\mathbb{C}}^i)$ .

$$\begin{array}{ccccccc} & & h^{00} & & & & \\ & h^{01} & & h^{10} & & & \\ h^{02} & & h^{11} & & h^{20} & & \\ h^{03} & h^{12} & & h^{21} & & h^{30} & \\ & h^{13} & h^{22} & & h^{31} & & \\ & & h^{23} & & h^{32} & & \\ & & & h^{33} & & & \end{array}$$

# Calabi–Yau varieties

## Definition

A **Calabi–Yau variety** is an irreducible, smooth, projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = \mathbb{C}$  and  $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ .
- The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .

- Easiest invariants are the **Euler characteristic** and the **Hodge numbers**,  $h^{ij} = h^j(X, \Omega_{X/\mathbb{C}}^i)$ .
- We always have  $\chi = 2(h^{11} - h^{12})$ .

# Calabi–Yau varieties

## Definition

A **Calabi–Yau variety** is an irreducible, smooth, projective scheme  $X/\mathbb{C}$  of dimension 3 satisfying:

- $H^0(X, \mathcal{O}_X) = H^3(X, \mathcal{O}_X) = \mathbb{C}$  and  $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ .
- The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .

- Easiest invariants are the **Euler characteristic** and the **Hodge numbers**,  $h^{ij} = h^j(X, \Omega_{X/\mathbb{C}}^i)$ .

- We always have  $\chi = 2(h^{11} - h^{12})$ .

$$\begin{array}{ccccccc}
 & & & & 1 & & \\
 & & & & 0 & & 0 \\
 & & & & h^{11} & & 0 \\
 & & 0 & & h^{12} & & 0 \\
 1 & & h^{12} & & h^{11} & & 1 \\
 & & 0 & & h^{11} & & 0 \\
 & & 0 & & 0 & & 0 \\
 & & & & 1 & & 
 \end{array}$$

# Hodge numbers heuristic

The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{11} = 1$  and  $h^{12} = 101$ .

# Hodge numbers heuristic

The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{11} = 1$  and  $h^{12} = 101$ .

## Heuristic.

The number  $h^{12}$  is the dimension of the “space of parameters” of  $X$ . The following heuristic will give us the correct Hodge number:

# Hodge numbers heuristic

The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{1,1} = 1$  and  $h^{1,2} = 101$ .

## Heuristic.

The number  $h^{1,2}$  is the dimension of the “space of parameters” of  $X$ . The following heuristic will give us the correct Hodge number:

- The space of degree 5 polynomials  $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))$  in  $\mathbb{P}^4$  is  $\binom{4+5}{4} = \binom{9}{4} = 126$ -dimensional. Hence  $\mathbb{P}(H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))) = \mathbb{P}^{125}$ .

# Hodge numbers heuristic

The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{11} = 1$  and  $h^{12} = 101$ .

## Heuristic.

The number  $h^{12}$  is the dimension of the “space of parameters” of  $X$ . The following heuristic will give us the correct Hodge number:

- The space of degree 5 polynomials  $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))$  in  $\mathbb{P}^4$  is  $\binom{4+5}{4} = \binom{9}{4} = 126$ -dimensional. Hence  $\mathbb{P}(H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))) = \mathbb{P}^{125}$ .
- This is not unique, but we can act by  $\mathrm{PGL}(5)$  to identify isomorphic quintics. We have  $\dim \mathrm{PGL}(5) = 25 - 1 = 24$ .

# Hodge numbers heuristic

The quintic  $X = V(f) \subset \mathbb{P}^4$  is the canonical example of a Calabi–Yau. It has Hodge numbers  $h^{11} = 1$  and  $h^{12} = 101$ .

## Heuristic.

The number  $h^{12}$  is the dimension of the “space of parameters” of  $X$ . The following heuristic will give us the correct Hodge number:

- The space of degree 5 polynomials  $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))$  in  $\mathbb{P}^4$  is  $\binom{4+5}{4} = \binom{9}{4} = 126$ -dimensional. Hence  $\mathbb{P}(H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(5))) = \mathbb{P}^{125}$ .
- This is not unique, but we can act by  $\mathrm{PGL}(5)$  to identify isomorphic quintics. We have  $\dim \mathrm{PGL}(5) = 25 - 1 = 24$ .
- In total:  $125 - 24 = 101$ , which is  $h^{12}(X)$ . ■



# Mirror symmetry

- Calabi–Yau threefolds seem to “always” have “mirror partners”.

$$\begin{array}{ccccccc}
 & & & X & & & \\
 \hline
 & & & 1 & & & \\
 & & 0 & & 0 & & \\
 & 0 & & 1 & & 0 & \\
 1 & & 101 & & 101 & & 1 \\
 & 0 & & 1 & & 0 & \\
 & & 0 & & 0 & & \\
 & & & 1 & & & 
 \end{array}$$

# Mirror symmetry

- Calabi–Yau threefolds seem to “always” have “mirror partners”.
- Mirror partner  $X^\circ$  to  $X$  has “mirrored Hodge diamond”.

$X$					
		1			
	0		0		
0		1		0	
1	101		101		1
	0	1		0	
	0		0		
		1			

$X^\circ$					
		1			
	0		0		
0		101		0	
1	1		1		1
	0	101		0	
	0		0		
		1			

# Mirror symmetry

- Calabi–Yau threefolds seem to “always” have “mirror partners”.
- Mirror partner  $X^\circ$  to  $X$  has “mirrored Hodge diamond”.
- Hence  $\chi(X^\circ) = -\chi(X)$ .

$X$					
		1			
	0		0		
	0	1		0	
1	101		101		1
	0	1		0	
	0		0		
		1			

$X^\circ$					
		1			
	0		0		
	0	101		0	
1	1		1		1
	0	101		0	
	0		0		
		1			

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ :

- 1 Suppose  $X$  has a natural degeneration  $X_0$  with a finite automorphism group  $G$ .
- 2 Find a family  $\pi: \mathcal{X} \rightarrow S$  on which  $G$  act, and such that the general fiber  $X_t$  has only isolated singularities.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ :

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 Find a family  $\pi: \mathcal{X} \rightarrow S$  on which  $G$  act, and such that the general fiber  $X_t$  has only isolated singularities.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ :

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There might be a finite subgroup  $H$  of the big torus acting. A mirror candidate is then a crepant resolution of  $X_t/H$ .

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ :

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There is an action of  $H \triangleq (\mathbb{Z}/5)^5/(\mathbb{Z}/5)$  on  $X_t$ .  
Crepant resolutions of  $X_t/H$  exists, and are mirrors.

# The orbifold heuristic

Sometimes the following method produces a mirror manifold of a Calabi–Yau  $X$ :

- 1 The general quintic degenerates to the singular scheme  $V(x_0x_1x_2x_3x_4)$ .
- 2 The family defined by  $f_t = x_0x_1x_2x_3x_4 + t \sum_{i=1}^5 x_i^5$  is  $S_5$ -invariant.
- 3 There is an action of  $H \triangleq (\mathbb{Z}/5)^5/(\mathbb{Z}/5)$  on  $X_t$ .  
Crepant resolutions of  $X_t/H$  exists, and are mirrors.

We can use **Roan's formula** to compute the Euler characteristic:

Theorem (Roan's formula)

$$\chi(\widetilde{X_t/H}) = \frac{1}{|H|} \sum_{g,h \in H} \chi(X_t^g \cap X_t^h)$$



# The cone over $dP_6$

- Let  $dP_6 \subset \mathbb{P}^6$  be an anticanonically embedded del Pezzo surface of degree 6. Let  $C(dP_6)$  be its affine cone in  $\mathbb{A}^7$ .
- The equations are

$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y & x_3 \\ x_5 & x_6 & y \end{vmatrix} \leq 1.$$

The origin is an isolated singularity.

# The cone over $dP_6$

- Let  $dP_6 \subset \mathbb{P}^6$  be an anticanonically embedded del Pezzo surface of degree 6. Let  $C(dP_6)$  be its affine cone in  $\mathbb{A}^7$ .
- The equations are
- There are two smoothing components.
- They come from perturbations of different forms of writing the equation.

$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y & x_3 \\ x_5 & x_6 & y \end{vmatrix} \leq 1.$$

The origin is an isolated singularity.

# The cone over $dP_6$

- Let  $dP_6 \subset \mathbb{P}^6$  be an anticanonically embedded del Pezzo surface of degree 6. Let  $C(dP_6)$  be its affine cone in  $\mathbb{A}^7$ .
- The equations are

$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y & x_3 \\ x_5 & x_6 & y \end{vmatrix} \leq 1.$$

The origin is an isolated singularity.

- There are two smoothing components.
- They come from perturbations of different forms of writing the equation.
- Can also write the equations as:

$$\leq 1$$

# The two smoothing components of $dP_6$

We can identify one component with  $\mathbb{P}(\mathcal{T}_{\mathbb{P}^2}) \setminus H$ , where  $\mathcal{T}_{\mathbb{P}^2}$  is the tangent bundle of  $\mathbb{P}^2$ .


$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y + t_1 & x_3 \\ x_5 & x_6 & y + t_2 \end{vmatrix} \leq 1$$

# The two smoothing components of $dP_6$

We can identify one component with  $\mathbb{P}(\mathcal{T}_{\mathbb{P}^2}) \setminus H$ , where  $\mathcal{T}_{\mathbb{P}^2}$  is the tangent bundle of  $\mathbb{P}^2$ .

$$\begin{vmatrix} y & x_1 & x_2 \\ x_4 & y + t_1 & x_3 \\ x_5 & x_6 & y + t_2 \end{vmatrix} \leq 1$$

The second component can be identified with  $(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1) \setminus H$ , where  $H$  is a  $(1, 1, 1)$ -divisor.



# Construction of a new Calabi–Yau: $X_1$

# Construction of a new Calabi–Yau: $X_1$

- Let  $E$  be a vector space with basis  $e_1, e_2, e_3$ . Consider

$$\mathbb{P}^{17} = \mathbb{P}((E \otimes E) \oplus (E \otimes E)).$$

The elements are pairs of  $3 \times 3$  matrices, equal up to scalar multiplication.

# Construction of a new Calabi–Yau: $X_1$

- Let  $E$  be a vector space with basis  $e_1, e_2, e_3$ . Consider

$$\mathbb{P}^{17} = \mathbb{P}((E \otimes E) \oplus (E \otimes E)).$$

The elements are pairs of  $3 \times 3$  matrices, equal up to scalar multiplication.

- Consider the set of pairs  $M$  of matrices  $(A, B)$  with rank  $1 + 1$ .



# Construction of a new Calabi–Yau: $X_1$

- Let  $E$  be a vector space with basis  $e_1, e_2, e_3$ . Consider

$$\mathbb{P}^{17} = \mathbb{P}((E \otimes E) \oplus (E \otimes E)).$$

The elements are pairs of  $3 \times 3$  matrices, equal up to scalar multiplication.

- Consider the set of pairs  $M$  of matrices  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $M$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{17}$ . Let  $X_1 \triangleq M \cap \mathbb{P}^{11}$ .

## Construction of a new Calabi–Yau: $X_1$

- Let  $E$  be a vector space with basis  $e_1, e_2, e_3$ . Consider

$$\mathbb{P}^{17} = \mathbb{P}((E \otimes E) \oplus (E \otimes E)).$$

The elements are pairs of  $3 \times 3$  matrices, equal up to scalar multiplication.

- Consider the set of pairs  $M$  of matrices  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $M$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{17}$ . Let  $X_1 \triangleq M \cap \mathbb{P}^{11}$ .

### Theorem

$X_1$  is a smooth Calabi–Yau with Euler characteristic  $-72$ .

## Construction of a new Calabi–Yau: $X_2$

- Let  $F$  be a vector space with basis  $f_1, f_2$ . Consider

$$\mathbb{P}^{15} = \mathbb{P}((F \otimes F \otimes F) \oplus (F \otimes F \otimes F)).$$

The elements are pairs of  $2 \times 2 \times 2$ -tensors, equal up to scalar multiplication.

- Consider the set of pairs  $N$  of tensors  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $N$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{15}$ . Let  $X_2 \triangleq N \cap \mathbb{P}^{11}$ .

### Theorem

$X_2$  is a smooth Calabi–Yau with Euler characteristic  $-48$ .

# Construction of a new Calabi–Yau: $X_3$

- Let  $E$  and  $F$  be as before. Consider

$$\mathbb{P}^{16} = \mathbb{P}((E \otimes E) \oplus (F \otimes F \otimes F)).$$

- Consider the set of pairs  $W$  of tensors  $(A, B)$  with rank  $1 + 1$ .
- Intersect  $W$  with a generic  $\mathbb{P}^{11} \subset \mathbb{P}^{16}$ . Let  $X_3 \triangleq W \cap \mathbb{P}^{11}$ .

## Theorem

$X_3$  is a smooth Calabi–Yau with Euler characteristic  $-60$ .

# Hodge number heuristics

## Conjecture

$X_1$  has Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$ .

# Hodge number heuristics

## Conjecture

$X_1$  has Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$ .
- 2 The Grassmannian of  $\mathbb{P}^{11}$ 's in  $\mathbb{P}^{17}$  is 72-dimensional.

# Hodge number heuristics

## Conjecture

$X_1$  has Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$ .
- 2 The Grassmannian of  $\mathbb{P}^{11}$ 's in  $\mathbb{P}^{17}$  is 72-dimensional.
- 3 We can act with automorphisms from  $\prod_{i=1}^4 \mathrm{GL}(E)$ .

# Hodge number heuristics

## Conjecture

$X_1$  has Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$ .
- 2 The Grassmannian of  $\mathbb{P}^{11}$ 's in  $\mathbb{P}^{17}$  is 72-dimensional.
- 3 We can act with automorphisms from  $\prod_{i=1}^4 \mathrm{GL}(E)$ .
- 4 The subgroup  $\{t_1 t_2 = t_3 t_4\} \subset (\mathbb{C}^*)^4 \subset \prod_{i=1}^4 \mathrm{GL}(E)$  acts trivially.



# Hodge number heuristics

## Conjecture

$X_1$  has Hodge numbers  $h^{11} = 3$  and  $h^{12} = 39$ .

## “Reason”.

- 1 We know the Euler characteristic, so it is enough to find  $h^{12}$ .
- 2 The Grassmannian of  $\mathbb{P}^{11}$ 's in  $\mathbb{P}^{17}$  is 72-dimensional.
- 3 We can act with automorphisms from  $\prod_{i=1}^4 \mathrm{GL}(E)$ .
- 4 The subgroup  $\{t_1 t_2 = t_3 t_4\} \subset (\mathbb{C}^*)^4 \subset \prod_{i=1}^4 \mathrm{GL}(E)$  acts trivially.
- 5 Hence  $h^{12} = 72 - \left( \dim \prod_{i=1}^4 \mathrm{GL}(E) - 3 \right) = 72 - 33 = 39$ . ■

## Mirror candidate for $X_1$

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

- There is an  $H = \mathbb{Z}/3$ -action on  $E$  defined by  $e_i \mapsto \omega^i e_i$ .

# Mirror candidate for $X_1$

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

- There is an  $H = \mathbb{Z}/3$ -action on  $E$  defined by  $e_i \mapsto \omega^i e_i$ .
- Another  $\mathbb{Z}/3$ -action  $e_i \mapsto e_{i+1}$ .

# Mirror candidate for $X_1$

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

- There is an  $H = \mathbb{Z}/3$ -action on  $E$  defined by  $e_i \mapsto \omega^i e_i$ .
- Another  $\mathbb{Z}/3$ -action  $e_i \mapsto e_{i+1}$ .
- Extends to actions on  $\mathbb{P}((E \otimes E) \oplus (E \otimes E)) = \mathbb{P}^{17}$ .

# Mirror candidate for $X_1$

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

- There is an  $H = \mathbb{Z}/3$ -action on  $E$  defined by  $e_i \mapsto \omega^i e_i$ .
- Another  $\mathbb{Z}/3$ -action  $e_i \mapsto e_{i+1}$ .
- Extends to actions on  $\mathbb{P}((E \otimes E) \oplus (E \otimes E)) = \mathbb{P}^{17}$ .
- Choose invariant  $\mathbb{P}^{11}$ : Defined by

$$f_{ij}^\alpha = e_{ij}^\alpha + t_{-i-j}^\alpha e_{-i-j, -i-j}^{\alpha+1}$$

for  $i, j \in \mathbb{Z}/3 \times \mathbb{Z}/3$  ( $i \neq j$ ) and  $\alpha = 0, 1$ .

# Mirror candidate for $X_1$

Using the mirror Ansatz, we propose mirror candidates for  $X_1$  and  $X_2$ .

- There is an  $H = \mathbb{Z}/3$ -action on  $E$  defined by  $e_i \mapsto \omega^i e_i$ .
- Another  $\mathbb{Z}/3$ -action  $e_i \mapsto e_{i+1}$ .
- Extends to actions on  $\mathbb{P}((E \otimes E) \oplus (E \otimes E)) = \mathbb{P}^{17}$ .
- Choose invariant  $\mathbb{P}^{11}$ : Defined by

$$f_{ij}^\alpha = e_{ij}^\alpha + t_{-i-j}^\alpha e_{-i-j, -i-j}^{\alpha+1}$$

for  $i, j \in \mathbb{Z}/3 \times \mathbb{Z}/3$  ( $i \neq j$ ) and  $\alpha = 0, 1$ .

- The resulting  $X_{H_t} \stackrel{\Delta}{=} \mathbb{P}^{11} \cap M$  is singular with 48 isolated double point singularities.

# Mirror candidate for $X_1$

- We divide out by the  $H$ -action and resolve:  $X_1^\circ \triangleq \widetilde{X_{H_t}}/H$ .

## Mirror candidate for $X_1$

- We divide out by the  $H$ -action and resolve:  $X_1^\circ \triangleq \widetilde{X_{H_t}}/H$ .
- Roan's formula gives:

$$\chi(X_1^\circ) = \frac{1}{3} (24 + 8 \cdot 24) = 72.$$

Based on this calculation and the mirror heuristic, we conjecture:

Conjecture

$X_1^\circ$  is a mirror of  $X_1$ .



## Mirror candidate for $X_1$

- We divide out by the  $H$ -action and resolve:  $X_1^\circ \triangleq \widetilde{X_{H_t}}/H$ .
- Roan's formula gives:

$$\chi(X_1^\circ) = \frac{1}{3} (24 + 8 \cdot 24) = 72.$$

Based on this calculation and the mirror heuristic, we conjecture:

### Conjecture

$X_1^\circ$  is a mirror of  $X_1$ .

### Remark

A very similar construction gives a mirror candidate for  $X_2$ .

UiO : **Department of Mathematics**  
University of Oslo



**Fredrik Meyer**



**Join of hexagons and  
Calabi–Yau threefolds**  
Public defence

