

Calculations

Fredrik Meyer

August 25, 2016

Contents

1	Computations on dP6	1
1.1	Finding equations of deformations	1
1.2	Intersecting with two special hyperplanes	3

1 Computations on dP6

1.1 Finding equations of deformations

Consider the del Pezzo surface dP_6 of degree 6 embedded in \mathbb{P}^6 . Its ideal is defined as follows:

```
1 restart
S = QQ[x_1..x_6,y_0]
3 I = ideal(x_1*x_3-x_2*y_0,
           x_2*x_4-x_3*y_0,
5    x_3*x_5-x_4*y_0,
  x_4*x_6-x_5*y_0,
7    x_5*x_1-x_6*y_0,
  x_6*x_2-x_1*y_0,
9    x_1*x_4-y_0^2,
  x_2*x_5-y_0^2,
11   x_3*x_6-y_0^2)
```

We compute the two deformations of its affine cone using the package **VersalDeformations**.

```
1 (F,R,G,C) = versalDeformation(gens I);
  decompose ideal transpose mingens ideal G
```

The output are four lists of matrices entries in $\mathbb{Q}[\mathbf{x}] \otimes \mathbb{Q}[t_1, t_2, t_3]$. The list F consists of the equations of the family, and the list R of the relations. The list G gives equations for the base space. We have that F_0 is the matrix of generators of I , and that $F_i R_i \equiv 0 \pmod{t^{i+1}}$.

The decomposition of `ideal G` is the following:

```
i9 : decompose ideal transpose mingens ideal G
o9 = {ideal (t1 - t3), ideal (t2 - t3, t1)}
```

Thus the base space splits into two components meeting transversely at the origin, of dimension 2 and 1, respectively. By doing a change of variables we can get rid of the linear terms:

```
1 T = QQ[x_1..x_6,y_0,t_1,t_2,t_3,s_1,s_2,s_3];
   gsub = sub(sub(ideal mingens ideal G,T), {t_2 => s_2+s_3-s_1,
   t_1 => s_3, t_3 => s_3-s_1})
3 fsub = transpose sub(sub(sum F,T), {t_2 => s_2+s_3-s_1, t_1 =>
   s_3, t_3 => s_3-s_1})
```

Now the equations are easier:

```
1 i13 : decompose gsub
o13 = {ideal (s1), ideal (s3, s2)}
```

We can get equations for each of these families by setting $s_1 = 0$ and $s_3 = s_2 = 0$, respectively:

```
1 i25 : fsub1 = sub(fsub, s_1 => 0)
3 o25 = {-2} | x_1x_3-x_2y_0 |
   {-2} | x_2x_4-x_3y_0+x_3s_2+x_3s_3 |
   {-2} | x_3x_5-x_4y_0+x_4s_3 |
   {-2} | x_4x_6-x_5y_0 |
   {-2} | x_1x_5-x_6y_0+x_6s_2+x_6s_3 |
   {-2} | x_2x_6-x_1y_0+x_1s_3 |
   {-2} | x_1x_4-y_0^2+y_0s_2+y_0s_3 |
   {-2} | x_2x_5-y_0^2+y_0s_2+2y_0s_3-s_2s_3-s_3^2 |
11  {-2} | x_3x_6-y_0^2+y_0s_3 |
```

```

13 |
    |          9          1
o25 : Matrix T <----- T
    |

```

And:

```

i26 : fsub2 = sub(fsub, {s_3 => 0, s_2 => 0})
2
o26 = {-2} | x_1x_3-x_2y_0 |
4         {-2} | x_2x_4-x_3y_0-x_3s_1 |
         {-2} | x_3x_5-x_4y_0 |
6         {-2} | x_4x_6-x_5y_0-x_5s_1 |
         {-2} | x_1x_5-x_6y_0 |
8         {-2} | x_2x_6-x_1y_0-x_1s_1 |
         {-2} | x_1x_4-y_0^2-y_0s_1 |
10        {-2} | x_2x_5-y_0^2-y_0s_1 |
         {-2} | x_3x_6-y_0^2-y_0s_1 |
12
    |          9          1
14 o26 : Matrix T <----- T
    |

```

1.2 Intersecting with two special hyperplanes

Consider dP_6 defined as above. Then consider the two hyperplanes

$$h_1 = x_1 + x_2 + x_3 + x_4 + x_5 + x_6$$

and

$$h_2 = x_1 - x_2 + x_3 - x_4 + x_5 - x_6.$$

We can compute the intersection with dP_6 in Macaulay2 as follows:

```

h1 = x_1+x_2+x_3+x_4+x_5+x_6
2 h2 = x_1-x_2+x_3-x_4+x_5-x_6
4 SS = S[r]/(r^2+3)
  apply(decompose(sub(I + h1 + h2, SS)), j -> ideal mingens sub(j,
    y_0 => 1))

```

The reason we create a new ring is that we need $\sqrt{-3}$ in order for the ideals to decompose. The results is the following list of ideals:

1. $(x_5 - 1, x_4 + x_6 + 1, x_3 + x_6 + 1, x_2 - 1, x_1 - x_6, r - 2x_6 - 1).$

2. $(x_5 - 1, x_4 + x_6 + 1, x_3 + x_6 + 1, x_2 - 1, x_1 - x_6, r + 2x_6 + 1).$
3. $(x_6 - 1, x_4 - x_5, x_3 - 1, x_2 + x_5 + 1, x_1 + x_5 + 1, r - 2x_5 - 1).$
4. $(x_6 - 1, x_4 - x_5, x_3 - 1, x_2 + x_5 + 1, x_1 + x_5 + 1, r + 2x_5 + 1).$
5. $(x_5 - x_6, x_4 - 1, x_3 + x_6 + 1, x_2 + x_6 + 1, x_1 - 1, r - 2x_6 - 1).$
6. $(x_5 - x_6, x_4 - 1, x_3 + x_6 + 1, x_2 + x_6 + 1, x_1 - 1, r + 2x_6 + 1).$

From this we can read off the coordinates in \mathbb{P}^6 :

Table 1: Table of singular points.

x_1	x_2	x_3	x_4	x_5	x_6	y_0
$\frac{-1+\sqrt{-3}}{2}$	1	$\frac{-1-\sqrt{-3}}{2}$	$\frac{-1-\sqrt{-3}}{2}$	1	$\frac{-1+\sqrt{-3}}{2}$	1
$\frac{-1+\sqrt{-3}}{2}$	1	$\frac{-1+\sqrt{-3}}{2}$	$\frac{-1+\sqrt{-3}}{2}$	1	$\frac{-1-\sqrt{-3}}{2}$	1
$\frac{-1-\sqrt{-3}}{2}$	$\frac{-1-\sqrt{-3}}{2}$	1	$\frac{-1+\sqrt{-3}}{2}$	$\frac{-1+\sqrt{-3}}{2}$	1	1
$\frac{-1+\sqrt{-3}}{2}$	$\frac{-1+\sqrt{-3}}{2}$	1	$\frac{-1-\sqrt{-3}}{2}$	$\frac{-1-\sqrt{-3}}{2}$	1	1
1	$\frac{-1-\sqrt{-3}}{2}$	$\frac{-1-\sqrt{-3}}{2}$	1	$\frac{-1+\sqrt{-3}}{2}$	$\frac{-1+\sqrt{-3}}{2}$	1
1	$\frac{-1+\sqrt{-3}}{2}$	$\frac{-1+\sqrt{-3}}{2}$	1	$\frac{-1-\sqrt{-3}}{2}$	$\frac{-1-\sqrt{-3}}{2}$	1

Note that the hexagonal group D_6 act transitively on the set of singular points.