

Notes toric

FM

1 The torus

Toric geometry is about the *algebraic torus*, and not the *topological torus*. The latter is a product of copies of S^1 , and the latter is a product of copies of \mathbb{C}^* . They are not completely dissimilar: the algebraic torus deformation retracts onto the topological torus, so they do have the same topological cohomology groups.

From now on, the word “torus” will always mean the algebraic torus.

What are the functions on the torus? Since we have the inclusion $(K^*)^n \subseteq K^n$, we have an inclusion $K[\mathbb{A}^n] \subseteq K[(K^*)^n]$. A short moment’s thought tells that the remaining functions are precisely the monomials. Thus we can say that $K[(K^*)^n] = K[\mathbb{Z}^n]$.

The invertible functions are the monomials. Those that map $1_T \mapsto 1_{K^*}$ are the monomials with coefficient 1, the *characters*: $\{\chi^u \mid u \in \mathbb{Z}^n\}$.

What are maps between tori? That is, we want to describe $\text{Hom}(T, T')$. We want the maps to preserve units. We can compose in any coordinate direction $T \rightarrow T' \rightarrow K^*$. A moment’s thought tells that the invertibility preserving maps are precisely the maps between the character lattices of the tori (in the opposite direction).

This is contravariant, and we would want to have a covariant correspondence. Introduce the *dual lattice*, namely $N = \text{Hom}(M, \mathbb{Z})$. Thus we see that maps between tori $T \rightarrow T'$ correspond bijectively naturally to maps between the dual lattices $N \rightarrow N'$.

The lattice N is often called the lattice of 1-parameter subgroups $K^* \rightarrow T$.

Definition 1.1. A *toric variety* is a normal equivariant partial compactification of the torus T . ■

Normality means that the coordinate ring of an affine patch is integrally closed in its fraction field. This is equivalent to Serre’s famous “R1+S2” condition.