

# Results so far

Fredrik Meyer

December 4, 2014

## 1 Introduction

In the article [Gai09], it is proven that there exists a triangulation of  $\mathbb{CP}^2$  with 15 vertices and other nice properties [[expand]]. The triangulation is constructed by “glueing” cones of 3-spheres along a triangle. Call this triangulation for  $\mathcal{S}$ .

A smoothing of the associated Stanley-Reisner scheme  $\mathbb{P}(\mathcal{S})$  would have interesting properties. In particular, it would be a hyper-Kähler variety [[EXPLAIN]]

The Hilbert polynomial of  $\mathbb{P}(\mathcal{S})$  is  $(9/2)t^4 + (15/2)t^2 + 3$ . The simplicial complex  $\mathcal{S}$  have  $f$ -vector  $(15, 90, 240, 270, 108)$ . In particular, the degree of  $\mathbb{P}(\mathcal{S})$  is 108.

Here are a few computations:

**Lemma 1.1.** *The module of first-order deformations of  $\mathbb{P}(\mathcal{S})$  is 90-dimensional, i.e.  $\dim_k T^1(\mathbb{P}(\mathcal{S})) = 90$ .*

**Lemma 1.2.** *The obstruction module have  $\dim_k T^2(\mathbb{P}(\mathcal{K})) = 306$ .*

## 2 Deformations of the 5-dimensional toric variety

Recall that the 5-dimensional toric variety  $T$  had 2-dimensional singularities (actually two disjoint copies of  $dP_6$ ).

**Theorem 2.1.** *There exists a flat deformation of  $\mathbb{P}(\mathcal{K} * \Delta^1)$ ,  $\mathfrak{X} \rightarrow S$ , such that  $\mathfrak{X}_{t_1} = T$  for some  $t_1 \in S$  and such that the general fiber  $\tilde{T}$  have one-dimensional singularities.*

*Proof.* As per now, the proof is purely by computer. The technique is this: First, consider the monomial degeneration of  $T$  to the Stanley-Reisner ring

$A(\mathcal{K} * \Delta^1)$  (recall that  $\mathcal{K} = D_6 * D_6$ ). Choose deformation parameters  $t_i$  perturbing the equations “in the direction of  $T$ ”, meaning that we only choose parameters introducing terms already occurring in the equations of  $T$ .

Now, using the package `VersalDeformations`, it is possible to produce a flat family  $\mathfrak{X} \rightarrow S'$  with  $\mathfrak{X}_0 = \mathbb{P}(\mathcal{K} * \Delta^1)$ .

The base space is a union of toric varieties of dimensions 14, 13, 13, 12, respectively. Call the largest one for  $S$ . The equations are

$$\begin{vmatrix} t_1 & t_2 & t_2 & \cdots & t_6 \\ t_7 & t_8 & t_9 & \cdots & t_{12} \end{vmatrix} \leq 1 \quad \begin{vmatrix} t_{13} & t_{14} & t_{15} & \cdots & t_{18} \\ t_{19} & t_{20} & t_{21} & \cdots & t_{24} \end{vmatrix} \leq 1$$

By restriction, we get the claimed family  $\mathfrak{X} \rightarrow T$ . It is checked by setting  $t_i = i$  for  $i = 1, \dots, 12$  and  $t_i = 2i$  for  $i = 13, \dots, 24$  that the resulting fiber have one-dimensional singularities. The reason we don't set all  $t_i = 1$ , is that this point lies in the intersection of the components of  $S$ .  $\square$

**Corollary 2.2.** *The Stanley-Reisner scheme  $\mathbb{P}(\mathcal{K})$  smooths to a smooth Calabi-Yau variety  $X$ .*

*Proof.* The scheme  $\mathbb{P}(\mathcal{K})$  sits as a complete intersection in  $\mathbb{P}(\mathcal{K} * \Delta^1)$ . Complete intersections deform together with the ambient variety, so  $\mathbb{P}(\mathcal{K} * \Delta^1)$  deforms to a general complete intersection in  $\tilde{T}$ . Since  $\tilde{T}$  have curve singularities, it follows by two applications of Bertini's theorem [Har77, Theorem II.8.18], that  $X$  is smooth.  $\square$

Now what we would really like to do is to compute the Hodge numbers  $h^{ij} = \dim_k H^j(X, \Omega_X^i)$  of  $X$ .

We can however compute the Hodge numbers of  $\tilde{T}$ . The hope is that there is some sort of Lefschetz theorem giving us the Hodge numbers of  $X$ .

**Theorem 2.3.** *We have  $h^{11}(\tilde{T}) = 1$  and  $h^{12}(\tilde{T}) = 12$ .*

*Proof.* Again, this is purely computational. We use long exact sequences together with sheaf cohomology computations in `Macaulay2`.

Since the ideal of  $\tilde{T}$  is rather complicated, doing this naively does not work. The trick is to choose the right term order. Since we know that  $\tilde{T}$  has a nice degeneration, we would like to find a term order such that its initial ideal is precisely the Stanley-Reisner ideal.

The `Macaulay2` package `gfanInterface` provides an interface with `gfan`, which is a program that can compute weight vectors given polynomials with prescribed initial terms. The weight vector is

$$\omega = (1, 1, 4, 7, 7, 4, 1, 1, 4, 7, 7, 4, 1, 1).$$

With this term order, giving a very small Gröbner basis (18 elements), the computations are much faster than with the standard term order. We are able to compute resolutions of all the relevant modules within a few minutes in total.

We have an exact sequence of sheaves on  $\tilde{T}$ :

$$0 \rightarrow \mathcal{I}_1 \hookrightarrow \mathcal{I}/\mathcal{I}^2 \xrightarrow{d} \Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}} \rightarrow \Omega_{\tilde{T}}^1 \rightarrow 0.$$

This sequence can be broken into two short exact sequences. The relevant one is this:

$$0 \rightarrow \text{im } d \rightarrow \Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}} \rightarrow \Omega_{\tilde{T}}^1 \rightarrow 0. \quad (1)$$

We also have the restricted Euler sequence:

$$0 \rightarrow \Omega_{\mathbb{P}^{13}} \otimes \mathcal{O}_{\tilde{T}} \rightarrow \mathcal{O}_{\tilde{T}}(-1)^{14} \rightarrow \mathcal{O}_{\tilde{T}} \rightarrow 0. \quad (2)$$

We first compute  $h^{11}$ . From (1) we get a long exact sequence

$$\dots \rightarrow H^1(\text{im } d) \rightarrow H^1(\Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}}) \rightarrow H^1(\Omega_{\tilde{T}}^1) \rightarrow H^2(\text{im } d) \rightarrow \dots$$

The cohomology of  $H^1(\text{im } d)$  and  $H^2(\text{im } d)$  was computed with `Macaulay2` to be both zero. Thus  $H^1(\Omega_{\tilde{T}}^1) \simeq H^1(\Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}})$ . From the Euler sequence we get

$$\dots \rightarrow H^0(\mathcal{O}_{\tilde{T}}(-1)^{14}) \rightarrow H^0(\mathcal{O}_{\tilde{T}}) \rightarrow H^1(\Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}}) \rightarrow H^1(\mathcal{O}_{\tilde{T}}(-1)^{14}) \rightarrow \dots$$

But the left and right terms are both zero. Hence  $h^{11} = 1$ . We now compute  $h^{12}$ .

From (1) we again get

$$0 \rightarrow H^2(\Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}}) \rightarrow H^2(\Omega_{\tilde{T}}^1) \rightarrow H^3(\text{im } d) \rightarrow H^3(\Omega_{\mathbb{P}^{13}}^1 \otimes \mathcal{O}_{\tilde{T}}) \rightarrow \dots,$$

where we have used that  $H^2(\text{im } d) = 0$ . But from the Euler sequence we get that the right term is also zero. Thus  $H^2(\Omega_{\tilde{T}}^1) \simeq H^3(\text{im } d)$ . This last group can be computed in `Macaulay2` to be 12-dimensional.  $\square$

**Remark.** *The reader may wonder we just didn't ask `Macaulay2` to compute the cohomology sheaf  $\Omega_{\tilde{T}}^1$  directly, e.g. by the command `HH^i(cotangentSheaf Proj A)`. The reason is that `Macaulay2`'s algorithms actually compute the sheaf, and not just the dimension, and this is too computationally intensive.*

**Remark (Question).** *A computation reveals that  $H^i(\mathcal{I}/\mathcal{I}^2) \simeq H^i(\text{im } d)$  for  $i \geq 2$ . This could be because the singularities are of dimension 1. Is there a theoretical result to this effect?*

We can compute it `Macaulay2` that:

**Lemma 2.4.** *The first cotangent module of  $\tilde{T}$  has  $\dim_k T^1(\tilde{T}/k) = 26$ .*

We have that  $h^0(\mathbb{P}(*\Delta^1), \mathcal{T}) = 14$  by Theorem 5.2 in [AC10]. Thus these numbers fit in the narrative that we should have  $T_{X_0}^1 = h^1(\mathcal{T}_{X_t}) + h^0(\mathcal{T}_{X_0})$ . (IS THERE ANY HEURISTIC FOR THIS??)

## References

- [AC10] Klaus Altmann and Jan Arthur Christoffersen. Deforming Stanley-Reisner schemes. *Math. Ann.*, 348(3):513–537, 2010.
- [Gaĭ09] A. A. Gaĭfullin. A minimal triangulation of the complex projective plane that admits a chessboard coloring of four-dimensional simplices. *Tr. Mat. Inst. Steklova*, 266(Geometriya, Topologiya i Matematicheskaya Fizika. II):33–53, 2009.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.