

Representation theory

Fredrik Meyer

Abstract

These are notes from the course MAT4270 on representation theory, autumn 2015. The lectures were held by Sergey Neshyevy. The notes are mine. The first half is about representations of finite groups, the second half about representations of compact Lie groups.

1 Lecture 1

From now on G is always a *finite group* and V a finite-dimensional vector space over the complex number. Most results will hold over any algebraically closed field of characteristic zero.

1.1 Motivating example

In 1896 Dedekind made the following observation. Let $\mathbb{C}[x_g \mid g \in G]$ be the free \mathbb{C} -algebra with basis the elements of G . Then one can form the following polynomial:

$$P_G(x_{\rho_1}, \dots, x_{\rho_n}) = \det \left((x_{\rho_i g_j})_{i,j=1}^n \right).$$

Note that the matrix is just the multiplication table for the group G . The problem is to decompose P_G into irreducible polynomials.

Example 1.1. Let $G = S_3$. Then G is generated by two elements r and s with the relation $srs = r^2$. If we write the elements of G as $\{e, r, r^2, s, sr, sr^2\}$, then the multiplication table looks like

$$\begin{pmatrix} e & r & r^2 & s & sr & sr^2 \\ r & r^2 & e & sr^2 & s & sr \\ r^2 & e & r & sr & sr^2 & s \\ s & sr & sr^2 & e & r & r^2 \\ sr & sr^2 & s & r^2 & e & r \\ sr^2 & s & sr & r & r^2 & e \end{pmatrix}.$$

The determinant is quite long, so I won't write it out. Suffice it to say it is a degree 6 polynomial with 146 terms. Using a computer algebra system such as `Macaulay2`, one can decompose it. It decomposes into three factors: one quadratic and two linear. They are the following (we change notation to avoid confusion about exponents):

$$x_e + x_r + x_{r^2} + x_s + x_{sr} + x_{sr^2}$$

and

$$x_e + x_r + x_{r^2} - x_s - x_{sr} - x_{sr^2}$$

and

$$x_e^2 - x_e x_r + x_{r^2} - x_e x_{r^2} - x_r x_{r^2} + x_{r^2}^2 - x_s^2 + x_s x_{sr} - x_{sr^2}^2 + x_s x_{sr^2} - x_{sr^2}^2 - x_{sr^2}^2.$$

We will see later that the first factor corresponds to the trivial representation, the second to the alternating representation, and the third is the so-called standard representation of G . ★

Frobenius proved the following theorem:

Theorem 1.2. *Let $P_G = \prod_{i=1}^r P_i^{m_i}$ be the decomposition of P_G into irreducibles. Then*

1. $m_i = \deg P_i$ for every i .
2. r is the number of conjugacy classes in G .

In particular, since $r = |G|$ if and only if G is abelian, P_G decomposes into linear factors if and only if G is abelian.

In trying to prove the above theorem, Frobenius had to develop representation theory.

1.2 Representations

A **representation of G on V** is a homomorphism $\pi : G \rightarrow \text{GL}(V)$. We will denote a representation interchangeably by π , (π, V) , V_π or V , depending upon the situation.

This means that a representation is a *linear* action of G on V .

Example 1.3. The trivial representation ϵ of any group is given by letting $V = \mathbb{C}$ and $\epsilon(g) = e$ for all $g \in G$. ★

Example 1.4 (Left regular representation). Let $\mathbb{C}[G]$ be the space of functions on G . It is a finite-dimensional vector space under pointwise addition. Define $\lambda : G \rightarrow \text{GL}(\mathbb{C}[G])$ by

$$g \mapsto (f \mapsto (h \mapsto f(g^{-1}h)))$$

This vector space have a basis consisting of the characteristic functions e_g for $g \in G$. On these elements one sees that the action is given by $g \cdot e_h = e_{gh}$. ★

Example 1.5. Similarly, one has the **permutation representation**. Let X be a G -set, i.e. a set on which G acts. Then one forms the space $\mathbb{C}[X]$ of X -valued functions. It has a basis e_x of characteristic functions, and the action of G on $\mathbb{C}[X]$ is given by $ge_x = e_{gx}$. ★

The notion of isomorphism is not surprising. An **equivalence** of representations (π, V) and (θ, W) is given by a vector space isomorphism $T : V \rightarrow W$ such that $T(g \cdot v) = g \cdot T(v)$ for all $g \in G$.

In case $T : V \rightarrow W$ is not an isomorphism, we say that T is an **intertwiner** of V and W . The set of intertwiners is denoted by $\text{Mor}(\pi, \theta)$ or $\text{Hom}_G(V, W)$.

A subspace $W \subset V$ is **invariant** if $g \cdot W \subset W$ for all $g \in G$. Letting $\theta(g) = \pi(g)|_W$, we get another representation of G , called a **subrepresentation** of G . We write $\pi|_W$ for θ .

If we have two representation (π, V) and (π', V') , we can form the **direct sum representation** by letting $g \in G$ act on $V \oplus V'$ componentwise. Note that π is a subrepresentation of $V \oplus V'$.

Proposition 1.6 (Mascke's theorem). *Let (π, V) be a representation of G and $W \subset V$ an invariant subspace. Then there exists a complementary invariant subspace W^\perp . That is, W is also invariant and such that $V = W \oplus W^\perp$.*

Proof. We prove this by “averaging”. Let $P : V \rightarrow W$ be any projection from V to W . Then define

$$\Pi(v) = \frac{1}{|G|} \sum_{g \in G} gP(g^{-1}v).$$

This is a G -linear morphism, because

$$\begin{aligned} \Pi(h \cdot v) &= \frac{1}{|G|} \sum_{g \in G} gP(g^{-1}h \cdot v) \\ &= \frac{1}{|G|} \sum_{g \in G} gP((h^{-1}g) \cdot v) \\ &= \frac{h}{|G|} \sum_{g \in G} h^{-1}gP((h^{-1}g) \cdot v). \end{aligned}$$

It is also surjective, since if $v \in W$, then clearly $\Pi(v) = v$. Hence Π is a G -linear surjection onto W . Then it is clear that $\ker \Pi$ is a G -invariant subspace of V complementary to W . □

We say that a representation is **irreducible** if it has no proper invariant subspaces. A representation is **completely reducible** if it decomposes into a direct sum of irreducible representations.

Example 1.7. Let $G = (\mathbb{R}, +)$ act on \mathbb{R}^2 by the matrices

$$\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}.$$

This leaves the x -axis fixed, so the representation is not irreducible. But it does not split into irreducible representations, since there is no complementary subspaces. ★

However, if G is finite (or **compact**, as we shall see later), things are nicer:

Theorem 1.8. *Any finitely-dimensional representation of a finite group G is completely reducible.*

The proof follows directly from Maschke's theorem (1.6).

We next prove Schur's lemma, which although has a very simple proof, has great consequences.

Proposition 1.9 (Schur's lemma). *Let (V, π) and (W, θ) be irreducible representations. Then*

1. *If $\pi \not\sim \theta$, then $\text{Hom}_G(V, W) = 0$.*
2. *If $\pi \sim \theta$, then $\text{Hom}_G(V, W)$ is 1-dimensional.*

Proof. Both the kernel and the image of a morphism of representations is a representation. But since V, W are irreducible, it follows that $\ker \varphi = 0$ or $\ker \varphi = V$ for any $\varphi : V \rightarrow W$. This proves i).

Now suppose $\Pi : V \rightarrow W$ is an equivalence of representations. Let $\varphi \in \text{Hom}_G(V, W)$. Then φ must be an equivalence as well, since the neither the kernel nor image can be non-zero. Consider $T = \varphi^{-1} \circ \Pi : V \rightarrow V$. This is a linear map, and since we work over \mathbb{C} , it has an eigenvalue λ . Now consider $T - \lambda \text{id}_V$. This has non-zero invariant kernel, hence $T - \lambda \text{id}_V = 0$. □

Proposition 1.10. *If G is abelian, then any irreducible representation of G is 1-dimensional.*

Proof. Each $g \in G$ gives a map $\pi(g) : V \rightarrow V \in \text{End}(V)$. This is even an intertwining map, because $\pi(g)\pi(h) = \pi(gh) = \pi(hg) = \pi(h)\pi(g)$.

If π is not trivial, by Schur's lemma, this map is just multiplication by some constant. But any 1-dimensional subspace of V is invariant under this action, so V must be 1-dimensional in order to be irreducible. □

Thus, up to equivalence, every irreducible representation of a finite abelian group is just a homomorphism $\chi : G \rightarrow \mathbb{C}$. But since G is finite, this actually maps into $\mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$. Such maps are called **characters** of G . Denote by \hat{G} the set of irreducible representations of G .

We call the \hat{G} the **Pontryagin dual** of G . It is an abelian group under pointwise multiplication. See Exercise 2 for more on this.

Remark. If G is non-abelian, there will always exist irreducible representations of dimension ≥ 2 . This follows because the regular representation $G \rightarrow \text{GL}(\mathbb{C}[G])$ is injective.

2 Lecture 2 - Density theorems

2.1 Density theorems

Let (V, π) be an irreducible representation. The set of operators $\pi(g)$ for $g \in G$ is quite large. By definition, if $x \neq 0$, then

$$V = \text{Span}\{\pi(g)x \mid g \in G\}.$$

By Schur's lemma, we also know that

$$\text{End}(\pi) = \{T \in \text{End}(V) \mid T\pi(g) = \pi(g)T \forall g \in G\} = \mathbb{C} \cdot \text{id}_V.$$

Theorem 2.1. Assume π_1, \dots, π_n are pairwise inequivalent irreducible representations, with $V_i = V_{\pi_i}$. Consider the direct sum representation $V_1 \oplus \dots \oplus V_n$.

Then

$$\text{Span}\{\pi(G)\} = \text{End}(V_1) \oplus \dots \oplus \text{End}(V_n).$$

Proof/exercise: We deduce the first theorem from the second: First break (V, π) into irreducible representations V_i . Then it follows directly from Schur's lemma that $\pi(G)' = \mathbb{C}\text{id}_{V_1} \oplus \dots \oplus \mathbb{C}\text{id}_{V_n}$.

Now, the elements of $\mathbb{C}\text{id}_{V_1} \oplus \dots \oplus \mathbb{C}\text{id}_{V_n}$ are block diagonal matrices. It is an easy exercise with matrices to see that the commutator of this set is just

$$\text{End}(V_1) \oplus \dots \oplus \text{End}(V_n).$$

□

First we need some notation. Let V be some vector space and $X \subset \text{End}(V)$ a subset of the endomorphisms of V . Let

$$X' \triangleq \{T \in \text{End}(V) \mid TS = ST \text{ for all } S \in X\}.$$

We call X' the **commutant** of X . Thus if (V, π) is a representation, then by definition:

$$\text{End}(\pi) = \pi(G)'.$$

Theorem 2.2. For any finite-dimensional representation (V, π) we have

$$\pi(G)'' = \text{Span}\{\pi(G)\}.$$

Proof. Let $T \in \pi(G)''$. Let x_1, \dots, x_n be a basis of V . Consider the representation $\theta = \pi \oplus \dots \oplus \pi$ (n times) on $W = \oplus_{i=1}^n V$. Consider $x = (x_1, \dots, x_n) \in W$ and the operator $S = T \oplus \dots \oplus T$ on W .

It is an exercise to show that $S \in \Theta(G)''$.

Now $\text{Span}\{\theta(G)x\}$ is an invariant subspace of W . Then there exists an invariant complementary subspace. Let $P : W \rightarrow \text{Span}\{\theta(G)x\}$ be the projection with kernel that invariant subspace, so that $P \in \Theta(G)'$ (that is, P is a G -morphism). Then

$$PSx = SPx = Sx$$

since $Px \in \text{Span}\{\theta(G)x\}$. Hence $Sx \in \text{Span}\{\theta(G)x\}$ as well. This means that there exist $\alpha_g \in \mathbb{C}$ such that

$$Sx = \sum_g \alpha_g \theta(g)x$$

This is the same as saying that

$$Tx_i = \sum_g \alpha_g \pi(g)x_i,$$

hence $T = \sum \alpha \pi(g)$. □

2.2 Orthogonality relations

Let (V, π) be a representation and let $\rho \in V^* = \text{Hom}(V, \mathbb{C})$. The functions $\alpha_{\rho, x}^\pi$ on G defined by $\alpha_{\rho, x}^\pi(g) = \rho(\pi(g)x)$ is called a **matrix coefficient** of π . If we fix a basis e_1, \dots, e_n of V and consider the dual basis e_1^*, \dots, e_n^* of V^* , then we write α_{ij}^π instead of $\alpha_{e_i^*, e_j}^\pi$.

In the basis e_1, \dots, e_n we have

$$\pi(g) = (\alpha_{ij}^\pi(g))_{ij}$$

Theorem 2.3. *We have:*

1. *If π and θ are inequivalent irreducible representations, then*

$$\frac{1}{|G|} \sum_g \alpha_{f, x}^\pi(g) \alpha_{\rho, y}^\theta(g^{-1}) = 0$$

for all $f \in V_\pi^$, $\rho \in V_\theta^*$, $x \in V_\pi$ and $y \in V_\theta$.*

2. *If π is irreducible, then*

$$\frac{1}{|G|} \sum_g \alpha_{f, x}^\pi(g) \alpha_{\rho, y}^\pi(g^{-1}) = \frac{f(x)\rho(y)}{\dim \pi}.$$

Proof. It is the averaging trick again. First observe that if $T : V_\theta \rightarrow V_\pi$ is any linear operator, then

$$\frac{1}{|G|} \sum_{g \in G} \pi(g) T \theta(g^{-1})$$

is in $\text{Hom}_G(\theta, \pi)$. □

A Exercises

A.1 Lecture 1

Exercise 1. Suppose $V_1 \oplus \dots \oplus V_r \sim W_1 \oplus \dots \oplus W_s$ as representations and that the V_i, W_i are irreducible. Show that $n = m$ and there exists $\rho \in S_r$ such that $V_i \sim W_{\rho(i)}$ for all i . ♠

Solution 1. First note a lemma: If $V \oplus W \sim V' \oplus W'$ and $V \sim V'$, then $W \sim W'$. This follows from the five-lemma.

Now consider the composition $V_1 \hookrightarrow V_1 \oplus \dots \oplus V_r \rightarrow W_1 \oplus \dots \oplus W_s \rightarrow W_i$. This has to be non-zero for at least one i . Hence $V_1 \sim W_i$ for some i . By rearranging, we are in the situation of the lemma above. Hence, inductively, if $r \leq s$, we find $0 \sim W_{r+1} \oplus \dots \oplus W_s$, which is impossible. Similarly for $s \leq r$. Hence $r = s$ and we conclude. ♥

Exercise 2. Let \hat{G} denote the Pontryagin dual of G . Then, for any finite abelian group G , we have $G \simeq \hat{\hat{G}}$. ♠

Solution 2. We first do this when $G = \mathbb{Z}/n$ for some $n \geq 0$. Let $\varphi : \mathbb{Z}/n \rightarrow \mathbb{C}^*$ be a character. Then $\phi(1) = z$ for some z , but $\phi(n) = \phi(0) = z^n = 1$, so z must be a n 'th root of unity.

Realize \mathbb{Z}/n as the n th roots of unity. Then we can define a homomorphism $\widehat{\mathbb{Z}/n} \rightarrow \mathbb{Z}/n$ by $\varphi \mapsto \varphi(1)$. Similarly, we can define an inverse map by sending $e^{2\pi i/n}$ to the character $m \mapsto e^{2\pi i m/n}$. These are inverses.

Now, every finite abelian group is a product of these groups. So it remains to show (by induction) that if G, H are two finite abelian groups, then $\widehat{G \times H} \simeq \hat{G} \times \hat{H}$. The inclusion maps $G \rightarrow G \times H$ and $H \rightarrow G \times H$ induces a map $\widehat{G \times H} \rightarrow \hat{G} \times \hat{H}$ by $(\varphi_1, \varphi_2) \mapsto ((g, h) \mapsto \varphi_1(g)\varphi_2(h))$.

It is easy to see that this map is injective. To see that it is surjective, let $\varphi : G \times H \rightarrow \mathbb{C}$ be a character. Write φ as $gh \mapsto \varphi(gh)$. Then we can define characters on G, H by $g \mapsto \varphi(g \cdot 1)$ which maps to φ . Ok. ♥

Exercise 3. Show that the S defined in the proof of Theorem (2.2) lies in $\Theta(G)''$. ♠

Solution 3. This is, I think, definition-hunting. DETAILS COME LATER ♡

B Some worked examples

B.1 The representation ring of S_3

We want to explicitly compute the representation ring of S_3 .

First we find all irreducible representations. The first one is the trivial representation $\epsilon : S_3 \rightarrow \mathbb{C}$. We also have the *sign* representation given by $g \mapsto \text{sgn}(g) \in \mathbb{C}$. Both of these are one-dimensional representations.

S_3 have a natural action on \mathbb{R}^3 given by permuting the basis vectors. But it acts trivially on the subspace spanned by $e_1 + e_2 + e_3$, hence \mathbb{R}^3 decomposes as $\epsilon \oplus V$, where V is some 2-dimensional representation. It is irreducible: if not, any $v \in V$ would be sent to a scalar multiple of itself, but this is not the case. This representation is called the **standard representation** of S_3 . We have also now found all representations of S_3 , since $1^2 + 1^2 + 2^2 = 6$.

Thus the representation ring $R(S_3)$ is $\mathbb{Z}[A, S]$ modulo some relations to be found. Here A is the alternating representation and S is the standard representation. The trivial representation is 1. It is easy to see that $A \otimes A \sim \epsilon$, so $A^2 = 1 \in R(S_3)$. The representation $S \otimes S$ is 4-dimensional. To compute how it decomposes, we use character theory.

Recall that characters are class functions on G , that is, they only depend on the conjugacy classes of G . So we write a character table:

	e	τ	σ
ϵ	1	1	1
A	1	1	-1
S	2	-1	0

To compute the character of the standard representation, we first note that it is, as a vector space given by $\mathbb{R}^3/(1,1,1)$. A basis is then given by the images of e_1, e_2 . Let τ be the transposition (123), sending e_i to e_{i+1} . Let σ be reflection fixing e_1 and exchanging e_2 and e_3 . In this basis that means e_2 is sent to $-e_1 - e_2$. Writing up the corresponding matrices lets us find the value of the character.

Now one can compute by hand (or use a result on characters), that the character of $S \otimes S$ is given by $\chi_S \cdot \chi_S$, so that its entry in the character table is $(4, 1, 0)$. If $S \otimes S = V_1 \oplus V_2$, then $\chi_S = \chi_{V_1} + \chi_{V_2}$.

Using also that the characters are linearly independent, we see that the only option is $S \otimes S \sim \epsilon \oplus A \oplus S$.

Hence $S^2 = 1 + A + S$ in the representation ring. Similarly, we find that $AS = S$ in the representation ring. All in all

$$R(S_3) = \mathbb{Z}[A, S]/(A^2 - 1, S^2 - 1 - A - S, AS - S).$$