

All matematikk

Fredrik Meyer

24. september 2014

Kapittel 1

Sets

1.1 Axioms

Kapittel 2

Relations

Definition 2.0.1. A *binary relation* is a map $S \times S \rightarrow \{\text{true}, \text{false}\}$ ■

Kapittel 3

Groups

Definition 3.0.2. A *group* is a triple (G, μ, ι) where G is a set and μ is a binary operation $G \times G \rightarrow G$ and ι is a function $G \rightarrow G$, satisfying the following axioms:

- Associativity: The following diagram commutes:

$$\begin{array}{ccc} G \times G \times G & \xrightarrow{\text{id} \times \mu} & G \times G \\ \mu \times \text{id} \downarrow & & \downarrow \mu \\ G \times G & \xrightarrow{\mu} & G \end{array}$$

- Identity element. There exist an element $e \in G$ such that the following two diagrams commute:

$$\begin{array}{ccc} \{e\} \times G & \xrightarrow{e \times \text{id}} & G \times G \\ & \searrow \pi_2 & \downarrow \mu \\ & & G \end{array}$$

and

$$\begin{array}{ccc} G \times \{e\} & \xrightarrow{\text{id} \times e} & G \times G \\ & \searrow \pi_2 & \downarrow \mu \\ & & G \end{array}$$

where the top maps are the natural inclusions.

- Inverse element. The following diagram is commutative:

$$\begin{array}{ccccc}
 G & \xrightarrow{\text{id} \times \iota} & G \times G & \xrightarrow{\mu} & G \\
 & \searrow & & \nearrow & \\
 & & \{e\} & &
 \end{array}$$

The same should also hold with $\text{id} \times \iota$ replaced with $\iota \times \text{id}$.

■

We will never write a group G as a triple, but only refer to the group (G, μ, ι) as just G , the maps being implicit. We will write $\iota(g)$ as g^{-1} , and $\mu(g, h)$ as gh .

Lemma 3.0.3. *The identity element $e \in G$ is unique.*

Bevis. Suppose e' is another identity element. Then $ee' = e'$ by the first diagram. But also $ee' = e$ by the second diagram. Then $e = e'$. Alternatively, the following is a commutative diagram:

$$\begin{array}{ccccc}
 & & \{e\} \times \{e'\} & & \\
 & \swarrow & \downarrow \Delta & \searrow & \\
 \{e\} \times G & \xrightarrow{e \times \text{id}} & G \times G & \xleftarrow{e' \times \text{id}} & \{e'\} \times G \\
 \searrow \pi_2 & & \downarrow \mu & & \swarrow \pi_2 \\
 & & G & &
 \end{array}$$

□

Kapittel 4

Rings

Kapittel 5

Fields

Definition 5.0.4. A *field* is a commutative ring k with only one ideal. ■

5.1 Ordered fields

Kapittel 6

The real numbers