

All of it

Fredrik Meyer

May 2, 2016



# Chapter 1

## Preliminary definitions

We work over  $\mathbb{C}$ .

### 1.1 Stanley-Reisner basics

Given a simplicial complex  $\mathcal{K}$ , one can associate to it a projective scheme  $\mathbb{P}(\mathcal{K})$  defined as follows. Let  $P$  be the polynomial ring with one variable for each vertex of  $\mathcal{K}$ . Then the *Stanley-Reisner ideal*  $I_{\mathcal{K}}$  corresponding to  $\mathcal{K}$  is generated by the monomials corresponding to *non-faces* of  $\mathcal{K}$ . Then we define the *Stanley-Reisner scheme* to be  $\text{Proj } P/I_{\mathcal{K}}$ .

**Example 1.1.1.** Let  $\mathcal{K}$  be the square, with vertices  $v_0, v_1, v_2, v_3$ . Then the Stanley-Reisner ideal is generated by  $v_0v_2$  and  $v_1v_3$ . ★

Some of the topology of the simplicial complex is encoded in the scheme structure of  $\mathbb{P}(\mathcal{K})$ . In particular, the simplicial (co)homology groups of  $\mathcal{K}$  can be computed as the sheaf cohomology of  $\mathbb{P}(\mathcal{K})$

**Lemma 1.1.2.** Let  $H^i(K; \mathbb{C})$  denote the singular cohomology groups of  $\mathcal{K}$ . Then there are isomorphisms  $H^i(K; \mathbb{C}) = H^i(\mathbb{P}(\mathcal{K}), \mathcal{O}_{\mathbb{P}(\mathcal{K})})$  for all  $i$ .

*Proof.* ———to come———

□

### 1.2 Calabi-Yau basics

**Definition 1.2.1.** A *Calabi-Yau variety* is a smooth projective variety satisfying the following two conditions:

1.  $H^i(X, \mathcal{O}_X) = 0$  for  $0 < i < \dim X$ .

2. The canonical sheaf is trivial:  $\omega_X \simeq \mathcal{O}_X$ .

■

The classical example of a Calabi-Yau manifold is the quintic threefold in  $\mathbb{P}^5$ . Another example is the following:

**Example 1.2.2.** Let  $X$  be the double cover of  $\mathbb{P}^3$  ramified along a smooth octic. The projection map is affine, so the conditions on  $H^i(X, \mathcal{O}_X)$  are fulfilled. To see that the canonical sheaf is trivial, we use the adjunction formula, which says that  $K_X = 2K_{\mathbb{P}^3}|_X + \deg R$ , where  $R$  is the ramification divisor. Then, since  $\omega_{\mathbb{P}^3} = \mathcal{O}_{\mathbb{P}^3}(-4)$ , it follows that  $K_X = 0$ . ★

If  $\mathcal{K}$  is a simplicial sphere, then a smoothing of  $\mathbb{P}(\mathcal{K})$  will give a Calabi-Yau manifold.

— ref: bayer-eisenbud graph curves.

### 1.3 Deformation theory

## Chapter 2

# Two topologically distinct smoothings

Denote by  $dP_6$  the del Pezzo surface of degree 6 embedded in  $\mathbb{P}^6$ . This can be realized as the blow-up of  $\mathbb{P}^2$  in three points not lying on a line. Let  $X$  denote the affine cone over  $dP_6$ . Then it has long been known that  $X$  has two smoothing components, and we show here that they are topologically distinct.

Recall that a *del Pezzo* surface is a surface such that the anti-canonical bundle is ample. The degree is the degree given by the anticanonical embedding. It is a classical result that every del Pezzo surface is obtained either by blowing up  $\mathbb{P}^2$  in  $r = 0, \dots, 6$  points in suitable positions, or as the 2-uple embedding of a quadric surface in  $\mathbb{P}^3$ .

### 2.1 Different embeddings of $dP_6$

We first obtain the equations of  $dP_6$  directly from the description of it as blow-up. Let  $x_0, x_1, x_2$  be coordinates of  $\mathbb{P}^2$ . Recall that the blowup of  $\mathbb{P}^2$  in the point  $(1 : 0 : 0)$  can be realized as the closed subscheme of  $\mathbb{P}^2 \times \mathbb{P}^1$  given by the equation  $r_0x_1 - r_1x_2 = 0$ , where  $r_0, r_1$  are coordinates on  $\mathbb{P}^1$ . We can repeat this process on the points  $(0 : 1 : 0)$  and  $(0 : 0 : 1)$  to obtain similar equations. Collecting these, we see that  $dP_6$  is given by the matrix equation

$$M\vec{x} = \begin{pmatrix} 0 & r_0 & -r_1 \\ s_1 & 0 & -s_0 \\ -t_0 & t_1 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \\ z_0 \end{pmatrix} = 0.$$

in  $\mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ . Here  $r_i, s_i$  and  $t_i$  ( $i = 0, 1$ ) are of course coordinates on  $\mathbb{P}^1$ .

We can do more than this however.

**Lemma 2.1.1.** *We can also realize  $dP_6$  embedded in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$  with equation  $r_0 s_0 t_0 = r_1 s_1 t_1$ .*

*Proof.* Note that the matrix cannot have rank 1 or lower.

Now consider the projection onto the last three factors:

$$\pi : \mathbb{P}^2 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1.$$

Each point  $P$  in the product on the right-hand side gives a matrix  $M_P$  of rank 2. Thus there is a line of solutions, which correspond exactly to a point in  $\mathbb{P}^2$ .

Hence the restriction of  $\pi$  to  $dP_6$  is an isomorphism onto the hypersurface given by  $\det M = 0$  in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ .  $\square$

Another way to realize blow-ups is this: let  $\mathfrak{d}$  be the linear system of quadrics with assigned basepoints  $(1 : 0 : 0)$ ,  $(0 : 1 : 0)$  and  $(0 : 0 : 1)$  in  $\mathbb{P}^2$ . We can choose a basis given by  $x_0 x_1, x_0 x_2$  and  $x_1 x_2$ . This gives a rational map  $\mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ . The closure of the graph of this map is a subvariety of  $\mathbb{P}^2 \times \mathbb{P}^2$  defined by two bilinear equations. Each of the projections correspond to the blowup.


Explicitly, if we let  $y_0, y_1, y_2$  be coordinates on the other  $\mathbb{P}^2$ , then the equations are  $x_1 y_0 - x_1 y_1 = x_1 y_1 - x_2 y_2 = 0$ .

We also have a natural embedding in  $\mathbb{P}^6$  as follows. Denote by  $E_1, E_2, E_3$  the exceptional divisors on the blowup. Let  $L$  be a line in  $\mathbb{P}^2$ . Then the divisor  $\pi^* 3L - E_1 - E_2 - E_3$  is ample, and gives an embedding in  $\mathbb{P}^6$  (see [1, Chapter V, Theorem 4.6]). A basis for the corresponding linear system is given by all monomials in  $\Gamma(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(3))$  except  $x^3, y^3$  and  $z^3$ .


The equations can be arranged in a particularly symmetric form: let  $y, x_1, \dots, x_5$  be coordinates on  $\mathbb{P}^6$ . Then the equations of  $dP_6$  are the  $2 \times 2$  minors of the matrix

$$\begin{pmatrix} x_1 & y & x_6 \\ x_2 & x_3 & y \\ y & x_4 & x_5 \end{pmatrix}.$$

This gives 9 equations, which can be compactly written as  $x_i x_{i+2} - y x_i = 0$  and  $x_i x_{i+3} - y^2 = 0$ , for  $i = 1, \dots, 6$  (where  $i$  is taken modulo 6). Note that the equations have a visible  $D_6$ -symmetry, where  $D_6$  denotes the dihedral group.



(a) The hexagon.

(b) The fan of  $dP_6$ .

### 2.1.1 As a toric variety

There is a nice combinatorial description of  $dP_6$  as a toric variety associated to a polytope. Namely, let  $P$  denote the hexagon in Figure 2.1a. Then the normal fan of this polytope defines a fan in  $N_{\mathbb{R}}$ , defining a toric variety.

The polytope is reflexive, implying that the normal fan of  $P$  is the face fan over the same polytope. See Figure 2.1b. From standard toric geometry, it is clear that  $dP_6$  is the blowup of  $\mathbb{P}^2$  in the three torus-fixed points.

## 2.2 The affine cone

## 2.3 The two smoothings





## Chapter 3

# A smooth Calabi-Yau

Consider the hexagon  $E_6$ . The join  $E_6 * E_6$  is a 3-dimensional sphere, and so a smoothing of the corresponding Stanley-Reisner scheme would correspond to a smooth Calabi-Yau manifold. In this chapter I prove that there does indeed exist a smoothing, and I describe some of its properties.



# Bibliography

- [1] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977.  
Graduate Texts in Mathematics, No. 52.