

# IMPLICITLY CONSTITUTED FLUID FLOW MODELS: ANALYSIS AND APPROXIMATION

Endre Süli

Mathematical Institute  
University of Oxford

Acknowledgements:

Miroslav Bulíček (Prague), Josef Málek (Prague),  
Lars Diering (Bielefeld), Christian Kreuzer (Dortmund),  
Tabea Tscherpel (Bielefeld), Alexei Gazca Orozco (Oxford), Patrick Farrell (Oxford)

Paris  
December 2020

# 1. Implicitly constituted fluid flow models

# 1. Implicitly constituted fluid flow models

Continuum mechanics of viscous fluids (at uniform temperature):

$$\begin{aligned}(\varrho \mathbf{u})_t + \operatorname{div}(\varrho \mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \varrho \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \varrho_t + \operatorname{div}(\varrho \mathbf{u}) &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) - \varrho \frac{d}{dt} \psi(\varrho) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

# 1. Implicitly constituted fluid flow models

Continuum mechanics of viscous fluids (at uniform temperature):

$$(\varrho \mathbf{u})_t + \operatorname{div}(\varrho \mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} = \varrho \mathbf{f} \quad \text{with } \mathbf{T} = \mathbf{T}^T,$$

$$\varrho_t + \operatorname{div}(\varrho \mathbf{u}) = 0,$$

$$\mathbf{T} : \mathbf{D}(\mathbf{u}) - \varrho \frac{d}{dt} \psi(\varrho) =: \xi \quad \text{with } \xi \geq 0.$$

$\varrho$  density

$\mathbf{u}$  velocity

$\mathbf{T}$  Cauchy stress tensor

$\mathbf{f}$  density of the external force

$\mathbf{D}(\mathbf{u}) := \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$  rate of strain tensor

$\psi = \psi(\varrho)$  Helmholtz free energy

$\xi$  entropy production

Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\boldsymbol{u} \otimes \boldsymbol{u}) - \operatorname{div} \boldsymbol{T} &= \boldsymbol{f} && \text{with } \boldsymbol{T} = \boldsymbol{T}^T, \\ \operatorname{div} \boldsymbol{u} &= 0, \\ \boldsymbol{T} : \boldsymbol{D}(\boldsymbol{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \operatorname{div} \mathbf{u} &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

To close the system, a constitutive relation, relating  $\mathbf{T}$  to  $\mathbf{D}(\mathbf{u})$ , is required.

Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \operatorname{div} \mathbf{u} &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

To close the system, a constitutive relation, relating  $\mathbf{T}$  to  $\mathbf{D}(\mathbf{u})$ , is required.

$\mathbf{T} = \mathbf{S} - p\mathbf{I}$ , with  $\mathbf{S}$  the deviatoric (shear) stress:  $\operatorname{tr}(\mathbf{S}) = 0$ ; and  $p$  the pressure.

Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \operatorname{div} \mathbf{u} &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

To close the system, a constitutive relation, relating  $\mathbf{T}$  to  $\mathbf{D}(\mathbf{u})$ , is required.

$\mathbf{T} = \mathbf{S} - p\mathbf{I}$ , with  $\mathbf{S}$  the deviatoric (shear) stress:  $\operatorname{tr}(\mathbf{S}) = 0$ ; and  $p$  the pressure.

Example: Newtonian (Navier–Stokes) fluid

$$\mathbf{S} = 2\mu \mathbf{D}(\mathbf{u}), \quad \mu > 0 \text{ viscosity coefficient.}$$



Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \operatorname{div} \mathbf{u} &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

To close the system, a constitutive relation, relating  $\mathbf{T}$  to  $\mathbf{D}(\mathbf{u})$ , is required.

$\mathbf{T} = \mathbf{S} - p\mathbf{I}$ , with  $\mathbf{S}$  the deviatoric (shear) stress:  $\operatorname{tr}(\mathbf{S}) = 0$ ; and  $p$  the pressure.

Example: Newtonian (Navier–Stokes) fluid

$$\mathbf{S} = 2\mu \mathbf{D}(\mathbf{u}), \quad \mu > 0 \text{ viscosity coefficient.}$$

Then:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \mu \Delta \mathbf{u} + \nabla p &= \mathbf{f} && \text{in } \Omega, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } \Omega,\end{aligned}$$

Special case: isothermal, homogeneous fluid at steady state:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{T} &= \mathbf{f} && \text{with } \mathbf{T} = \mathbf{T}^T, \\ \operatorname{div} \mathbf{u} &= 0, \\ \mathbf{T} : \mathbf{D}(\mathbf{u}) &=: \xi && \text{with } \xi \geq 0.\end{aligned}$$

To close the system, a constitutive relation, relating  $\mathbf{T}$  to  $\mathbf{D}(\mathbf{u})$ , is required.

$\mathbf{T} = \mathbf{S} - p\mathbf{I}$ , with  $\mathbf{S}$  the deviatoric (shear) stress:  $\operatorname{tr}(\mathbf{S}) = 0$ ; and  $p$  the pressure.

Example: Newtonian (Navier–Stokes) fluid

$$\mathbf{S} = 2\mu \mathbf{D}(\mathbf{u}), \quad \mu > 0 \text{ viscosity coefficient.}$$

Then:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \mu \Delta \mathbf{u} + \nabla p &= \mathbf{f} && \text{in } \Omega, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } \Omega,\end{aligned}$$

$$\begin{aligned}\xi := \mathbf{T} : \mathbf{D}(\mathbf{u}) &= \mathbf{S} : \mathbf{D}(\mathbf{u}) = \mu |\mathbf{D}(\mathbf{u})|^2 + \frac{1}{4\mu} |\mathbf{S}|^2 \geq \min\left(\mu, \frac{1}{4\mu}\right) (|\mathbf{D}(\mathbf{u})|^2 + |\mathbf{S}|^2) \\ &\geq 0. \quad \checkmark\end{aligned}$$

## Examples of constitutive relations

- Power-law (Ostwald–de Waele) fluids:

$$\boldsymbol{S} = 2\mu |\boldsymbol{D}(\boldsymbol{u})|^{r-2} \boldsymbol{D}(\boldsymbol{u}), \quad \text{with } 1 < r < \infty;$$

## Examples of constitutive relations

- Power-law (Ostwald–de Waele) fluids:

$$\boldsymbol{S} = 2\mu |\boldsymbol{D}(\boldsymbol{u})|^{r-2} \boldsymbol{D}(\boldsymbol{u}), \quad \text{with } 1 < r < \infty;$$

- Generalized power-law fluids:  $\boldsymbol{S} = 2\mu(|\boldsymbol{D}(\boldsymbol{u})|^2) \boldsymbol{D}(\boldsymbol{u});$

# Examples of constitutive relations

- Power-law (Ostwald–de Waele) fluids:

$$\mathbf{S} = 2\mu |\mathbf{D}(\mathbf{u})|^{r-2} \mathbf{D}(\mathbf{u}), \quad \text{with } 1 < r < \infty;$$

- Generalized power-law fluids:  $\mathbf{S} = 2\mu(|\mathbf{D}(\mathbf{u})|^2) \mathbf{D}(\mathbf{u})$ ;
- Stress power-law fluids and generalizations:  $\mathbf{D}(\mathbf{u}) = \alpha(|\mathbf{S}|^2) \mathbf{S}$ ;

## Examples of constitutive relations

- Power-law (Ostwald–de Waele) fluids:

$$\mathbf{S} = 2\mu|\mathbf{D}(\mathbf{u})|^{r-2} \mathbf{D}(\mathbf{u}), \quad \text{with } 1 < r < \infty;$$

- Generalized power-law fluids:  $\mathbf{S} = 2\mu(|\mathbf{D}(\mathbf{u})|^2) \mathbf{D}(\mathbf{u})$ ;
- Stress power-law fluids and generalizations:  $\mathbf{D}(\mathbf{u}) = \alpha(|\mathbf{S}|^2) \mathbf{S}$ ;
- Activated fluids, such as Bingham and Herschel–Bulkley fluids ( $\mathbf{D} := \mathbf{D}(\mathbf{u})$ ):

$$|\mathbf{S}| \leq \tau_* \Leftrightarrow \mathbf{D} = \mathbf{0} \quad \text{and} \quad |\mathbf{S}| > \tau_* \Leftrightarrow \mathbf{S} = \frac{\tau_* \mathbf{D}}{|\mathbf{D}|} + 2\nu(|\mathbf{D}|^2) \mathbf{D}.$$

# Examples of constitutive relations

- Power-law (Ostwald–de Waele) fluids:

$$\mathbf{S} = 2\mu |\mathbf{D}(\mathbf{u})|^{r-2} \mathbf{D}(\mathbf{u}), \quad \text{with } 1 < r < \infty;$$

- Generalized power-law fluids:  $\mathbf{S} = 2\mu(|\mathbf{D}(\mathbf{u})|^2) \mathbf{D}(\mathbf{u})$ ;
- Stress power-law fluids and generalizations:  $\mathbf{D}(\mathbf{u}) = \alpha(|\mathbf{S}|^2) \mathbf{S}$ ;
- Activated fluids, such as Bingham and Herschel–Bulkley fluids ( $\mathbf{D} := \mathbf{D}(\mathbf{u})$ ):

$$|\mathbf{S}| \leq \tau_* \Leftrightarrow \mathbf{D} = \mathbf{0} \quad \text{and} \quad |\mathbf{S}| > \tau_* \Leftrightarrow \mathbf{S} = \frac{\tau_* \mathbf{D}}{|\mathbf{D}|} + 2\nu(|\mathbf{D}|^2) \mathbf{D}.$$

Equivalently:

$$2\nu(|\mathbf{D}|^2) (\tau_* + (|\mathbf{S}| - \tau_*)_+) \mathbf{D} = (|\mathbf{S}| - \tau_*)_+ \mathbf{S}, \quad \tau_* > 0.$$

# Examples of constitutive relations between $S$ and $D$




Image source: S. K. Kawatra & A. K. Bakshi, Int. J. Mineral Process., Vol. 47(3–4), pp. 275–283, 1996.



# Examples of constitutive relations between $S$ and $D$




Image source: S. K. Kawatra & A. K. Bakshi, *Int. J. Mineral Process.*, Vol. 47(3–4), pp. 275–283, 1996.



K. R. Rajagopal. *On implicit constitutive theories for fluids*, *J. Fluid Mech.* **550** (2006), pp. 243–249.



K. R. Rajagopal & A. R. Srinivasa. *On the thermodynamics of fluids defined by implicit constitutive relations*, *Z. Angew. Math. Phys.* **59**(4) (2008), pp. 715–729.

# Examples of constitutive relations between $S$ and $D$




Image source: S. K. Kawatra & A. K. Bakshi, Int. J. Mineral Process., Vol. 47(3–4), pp. 275–283, 1996.



M. Bulíček, P. Gwiazda, J. Málek, and A. Świerczewska-Gwiazda. *On steady flows of incompressible fluids with implicit power-law-like rheology*. Advances in Calculus of Variations, 2009.



M. Bulíček, P. Gwiazda, J. Málek, and A. Świerczewska-Gwiazda. *On unsteady flows of implicitly constituted incompressible fluids*. SIAM Journal on Mathematical Analysis, 2012.

## Problem

In a bounded open Lipschitz domain  $\Omega \subset \mathbb{R}^d$ ,  $d \in \{2, 3\}$ , consider:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{S} + \nabla p &= \mathbf{f} && \text{in } \Omega, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } \Omega, \\ \mathbf{u} &= \mathbf{0} && \text{on } \partial\Omega.\end{aligned}\tag{Eq}$$

# Problem

In a bounded open Lipschitz domain  $\Omega \subset \mathbb{R}^d$ ,  $d \in \{2, 3\}$ , consider:

$$\begin{aligned}\operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div} \mathbf{S} + \nabla p &= \mathbf{f} && \text{in } \Omega, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } \Omega, \\ \mathbf{u} &= \mathbf{0} && \text{on } \partial\Omega.\end{aligned}\tag{Eq}$$

Here:

- $\mathbf{u} : \Omega \rightarrow \mathbb{R}^d$ : velocity of the fluid;
- $p : \Omega \rightarrow \mathbb{R}$ : pressure;
- $\mathbf{S} = \mathbf{S}^T : \Omega \rightarrow \mathbb{R}_{\text{sym},0}^{d \times d}$ : shear stress and  $\mathbf{D}(\mathbf{u}) := \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$  are assumed to be related through the **implicit constitutive relation**

$$\mathbf{G}(\mathbf{S}, \mathbf{D}(\mathbf{u})) = \mathbf{0},$$

where  $\mathbf{G} : \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d} \rightarrow \mathbb{R}_{\text{sym},0}^{d \times d}$ .

We identify the implicit relation with a graph  $\mathcal{A} \subset \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ :

$$\mathbf{G}(\mathbf{S}, \mathbf{D}) = \mathbf{0} \iff (\mathbf{D}, \mathbf{S}) \in \mathcal{A}.$$

We assume that  $\mathcal{A}$  is a *maximal monotone  $r$ -graph*:

We identify the implicit relation with a graph  $\mathcal{A} \subset \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ :

$$\mathbf{G}(S, D) = \mathbf{0} \iff (D, S) \in \mathcal{A}.$$

We assume that  $\mathcal{A}$  is a *maximal monotone  $r$ -graph*:

**(A1)**  $\mathcal{A}$  includes the origin; i.e.,  $(\mathbf{0}, \mathbf{0}) \in \mathcal{A}$ ;

We identify the implicit relation with a graph  $\mathcal{A} \subset \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ :

$$\mathbf{G}(\mathbf{S}, \mathbf{D}) = \mathbf{0} \iff (\mathbf{D}, \mathbf{S}) \in \mathcal{A}.$$

We assume that  $\mathcal{A}$  is a *maximal monotone r-graph*:

**(A1)**  $\mathcal{A}$  includes the origin; i.e.,  $(\mathbf{0}, \mathbf{0}) \in \mathcal{A}$ ;

**(A2)**  $\mathcal{A}$  is a monotone graph; i.e.,

$$(\mathbf{S}_1 - \mathbf{S}_2) : (\mathbf{D}_1 - \mathbf{D}_2) \geq 0 \text{ for all } (\mathbf{D}_1, \mathbf{S}_1), (\mathbf{D}_2, \mathbf{S}_2) \in \mathcal{A};$$

We identify the implicit relation with a graph  $\mathcal{A} \subset \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ :

$$\mathbf{G}(\mathbf{S}, \mathbf{D}) = \mathbf{0} \iff (\mathbf{D}, \mathbf{S}) \in \mathcal{A}.$$

We assume that  $\mathcal{A}$  is a *maximal monotone r-graph*:

**(A1)**  $\mathcal{A}$  includes the origin; i.e.,  $(\mathbf{0}, \mathbf{0}) \in \mathcal{A}$ ;

**(A2)**  $\mathcal{A}$  is a monotone graph; i.e.,

$$(\mathbf{S}_1 - \mathbf{S}_2) : (\mathbf{D}_1 - \mathbf{D}_2) \geq 0 \text{ for all } (\mathbf{D}_1, \mathbf{S}_1), (\mathbf{D}_2, \mathbf{S}_2) \in \mathcal{A};$$

**(A3)**  $\mathcal{A}$  is a maximal monotone graph; i.e., for any  $(\mathbf{D}, \mathbf{S}) \in \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ ,

$$\text{if } (\mathbf{S} - \tilde{\mathbf{S}}) : (\mathbf{D} - \tilde{\mathbf{D}}) \geq 0 \text{ for all } (\tilde{\mathbf{D}}, \tilde{\mathbf{S}}) \in \mathcal{A}, \text{ then } (\mathbf{D}, \mathbf{S}) \in \mathcal{A};$$



We identify the implicit relation with a graph  $\mathcal{A} \subset \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ :

$$\mathbf{G}(\mathbf{S}, \mathbf{D}) = \mathbf{0} \iff (\mathbf{D}, \mathbf{S}) \in \mathcal{A}.$$

We assume that  $\mathcal{A}$  is a *maximal monotone  $r$ -graph*:

**(A1)**  $\mathcal{A}$  includes the origin; i.e.,  $(\mathbf{0}, \mathbf{0}) \in \mathcal{A}$ ;

**(A2)**  $\mathcal{A}$  is a monotone graph; i.e.,

$$(\mathbf{S}_1 - \mathbf{S}_2) : (\mathbf{D}_1 - \mathbf{D}_2) \geq 0 \text{ for all } (\mathbf{D}_1, \mathbf{S}_1), (\mathbf{D}_2, \mathbf{S}_2) \in \mathcal{A};$$

**(A3)**  $\mathcal{A}$  is a maximal monotone graph; i.e., for any  $(\mathbf{D}, \mathbf{S}) \in \mathbb{R}_{\text{sym},0}^{d \times d} \times \mathbb{R}_{\text{sym},0}^{d \times d}$ ,

$$\text{if } (\mathbf{S} - \tilde{\mathbf{S}}) : (\mathbf{D} - \tilde{\mathbf{D}}) \geq 0 \text{ for all } (\tilde{\mathbf{D}}, \tilde{\mathbf{S}}) \in \mathcal{A}, \text{ then } (\mathbf{D}, \mathbf{S}) \in \mathcal{A};$$

**(A4)**  $\mathcal{A}$  is an  $r$ -graph; i.e., there exists a constant  $C_1 > 0$  s.t.:

$$\mathbf{S} : \mathbf{D} \geq C_1(|\mathbf{D}|^r + |\mathbf{S}|^{r'}) \text{ for all } (\mathbf{D}, \mathbf{S}) \in \mathcal{A},$$


where  $\frac{1}{r} + \frac{1}{r'} = 1$  and  $1 < r < \infty$ .

## 2. Numerical approximation

- Approximate the implicit constitutive law  $\mathcal{S}$  by a sequence of explicit laws  $\mathcal{S}^n$  by convolving a selection  $\mathcal{S}^*$  with compactly supported  $\theta_n \in L^1(\mathbb{R}; \mathbb{R}_{\geq 0})$ .


## 2. Numerical approximation

- Approximate the implicit constitutive law  $\mathcal{S}$  by a sequence of explicit laws  $\mathcal{S}^n$  by convolving a selection  $\mathcal{S}^*$  with compactly supported  $\theta_n \in L^1(\mathbb{R}; \mathbb{R}_{\geq 0})$ .



## 2. Numerical approximation


- Approximate the implicit constitutive law  $\mathcal{S}$  by a sequence of explicit laws  $\mathcal{S}^n$  by convolving a selection  $\mathcal{S}^*$  with compactly supported  $\theta_n \in L^1(\mathbb{R}; \mathbb{R}_{\geq 0})$ .



Alternative: Yosida regularization of the graph  $\mathcal{A} \subset \mathbb{R}^{d \times d}_{\text{sym},0} \times \mathbb{R}^{d \times d}_{\text{sym},0}$ :


$$\mathcal{A}^n := \left\{ \left( D + \frac{1}{n} |\mathcal{S}|^{\frac{2-q}{q-1}} \mathcal{S}, \mathcal{S} \right) \in \mathbb{R}^{d \times d}_{\text{sym},0} \times \mathbb{R}^{d \times d}_{\text{sym},0} : (D, \mathcal{S}) \in \mathcal{A} \right\}, \quad q \in (1, \infty).$$

## Computational grid: grid-size $h \ll 1$



Discretize the resulting PDE by a family of finite element spaces  $\mathbb{V}_h \times \mathbb{Q}_h$  contained in  $W_0^{1,r}(\Omega)^d \times L_0^{\tilde{r}}(\Omega)$ , on conforming and shape-regular grids  $\mathcal{T}_h$ , where the union of  $\{\mathbb{V}_h\}_{h>0}$  and  $\{\mathbb{Q}_h\}_{h>0}$  is dense in the respective space.

## Computational grid: grid-size $h \ll 1$



Discretize the resulting PDE by a family of finite element spaces  $\mathbb{V}_h \times \mathbb{Q}_h$  contained in  $W_0^{1,r}(\Omega)^d \times L_0^{\tilde{r}}(\Omega)$ , on conforming and shape-regular grids  $\mathcal{T}_h$ , where the union of  $\{\mathbb{V}_h\}_{h>0}$  and  $\{\mathbb{Q}_h\}_{h>0}$  is dense in the respective space.

Notation:

$$\tilde{r} = \begin{cases} \min \left\{ r', \frac{dr}{2(d-r)} \right\}, & \text{if } r < d, \\ r', & \text{if } r \geq d \end{cases} \leq r'.$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) \cdot \mathbf{u} \, dx + \int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx - \int_{\Omega} p(\operatorname{div} \mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) \cdot \mathbf{u} \, dx + \int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$



## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\boldsymbol{v} = \boldsymbol{u}$ ,  $\operatorname{div} \boldsymbol{u} = 0$ :

$$\int_{\Omega} \boldsymbol{S} : \boldsymbol{D}(\boldsymbol{u}) \, dx = \int_{\Omega} \boldsymbol{f} \cdot \boldsymbol{u} \, dx$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \|\mathbf{f}\|_{W^{-1,r'}(\Omega)} \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \|\mathbf{f}\|_{W^{-1,r'}(\Omega)} \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}$$

$$\|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \leq C_2 \|\mathbf{f}\|_{W^{-1,r'}(\Omega)}^{r'}.$$

## Formal energy (in)equality

Take scalar product of momentum eqn. with  $\mathbf{v} = \mathbf{u}$ ,  $\operatorname{div} \mathbf{u} = 0$ :

$$\int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{u}) \, dx = \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \int_{\Omega} \mathbf{f} \cdot \mathbf{u} \, dx$$

$$C_1 \left( \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \right) \leq \|\mathbf{f}\|_{W^{-1,r'}(\Omega)} \|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}$$

$$\|\mathbf{D}(\mathbf{u})\|_{L^r(\Omega)}^r + \|\mathbf{S}\|_{L^{r'}(\Omega)}^{r'} \leq C_2 \|\mathbf{f}\|_{W^{-1,r'}(\Omega)}^{r'}.$$

→ This argument must now be replicated at the discrete level.

## Choice of the finite element spaces $\mathbb{V}_h$ and $\mathbb{Q}_h$

Must have:

$$\int_{\Omega} q_h (\operatorname{div} \mathbf{u}_h) \, dx = 0 \quad \forall q_h \in \mathbb{Q}_h.$$

## Choice of the finite element spaces $\mathbb{V}_h$ and $\mathbb{Q}_h$

Must have:

$$\int_{\Omega} q_h (\operatorname{div} \mathbf{u}_h) \, dx = 0 \quad \forall q_h \in \mathbb{Q}_h.$$

**Option (A):** Taylor–Hood (P2/P1), Conforming Crouzeix–Raviart Stokes element (P2+bubble/P1); Mini element (P1+bubble/P1); P2/P0 ( $d = 2$ ); P3/P0 ( $d = 3$ );

## Choice of the finite element spaces $\mathbb{V}_h$ and $\mathbb{Q}_h$

Must have:

$$\int_{\Omega} q_h (\operatorname{div} \mathbf{u}_h) \, dx = 0 \quad \forall q_h \in \mathbb{Q}_h.$$

**Option (A):** Taylor–Hood (P2/P1), Conforming Crouzeix–Raviart Stokes element (P2+bubble/P1); Mini element (P1+bubble/P1); P2/P0 ( $d = 2$ ); P3/P0 ( $d = 3$ );

**Option (B):** Pointwise divergence-free velocity space (Scott–Vogelius; Guzmán–Neilan).



## Choice of the finite element spaces $\mathbb{V}_h$ and $\mathbb{Q}_h$

Must have:

$$\int_{\Omega} q_h (\operatorname{div} \mathbf{u}_h) \, dx = 0 \quad \forall q_h \in \mathbb{Q}_h.$$

**Option (A):** Taylor–Hood (P2/P1), Conforming Crouzeix–Raviart Stokes element (P2+bubble/P1); Mini element (P1+bubble/P1); P2/P0 ( $d = 2$ ); P3/P0 ( $d = 3$ );

**Option (B):** Pointwise divergence-free velocity space (Scott–Vogelius; Guzmán–Neilan).

Problem (in the case of Option (A)):

In general  $\operatorname{div} \mathbf{u}_h \not\equiv 0$  pointwise on  $\Omega$ , and thus

$$\int_{\Omega} \operatorname{div}(\mathbf{u}_h \otimes \mathbf{u}_h) \cdot \mathbf{u}_h \, dx \neq 0.$$

## Choice of the finite element spaces $\mathbb{V}_h$ and $\mathbb{Q}_h$

Must have:

$$\int_{\Omega} q_h (\operatorname{div} \mathbf{u}_h) \, dx = 0 \quad \forall q_h \in \mathbb{Q}_h.$$

**Option (A):** Taylor–Hood (P2/P1), Conforming Crouzeix–Raviart Stokes element (P2+bubble/P1); Mini element (P1+bubble/P1); P2/P0 ( $d = 2$ ); P3/P0 ( $d = 3$ );

**Option (B):** Pointwise divergence-free velocity space (Scott–Vogelius; Guzmán–Neilan).

Problem (in the case of Option (A)):

In general  $\operatorname{div} \mathbf{u}_h \not\equiv 0$  pointwise on  $\Omega$ , and thus

$$\int_{\Omega} \operatorname{div}(\mathbf{u}_h \otimes \mathbf{u}_h) \cdot \mathbf{u}_h \, dx \neq 0.$$

Fix (in the case of option (A)): replace  $\int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{v}) \cdot \mathbf{w} \, dx$  by

$$\mathcal{B}[\mathbf{u}, \mathbf{v}, \mathbf{w}] := \frac{1}{2} \int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{v}) \cdot \mathbf{w} - \operatorname{div}(\mathbf{u} \otimes \mathbf{w}) \cdot \mathbf{v} \, dx$$



$$\mathcal{B}[\mathbf{v}, \mathbf{v}, \mathbf{v}] = 0 \quad \forall \mathbf{v}, \quad \mathcal{B}[\mathbf{u}, \mathbf{v}, \mathbf{w}] = \int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{v}) \cdot \mathbf{w} \, dx \quad \text{if } \operatorname{div} \mathbf{u} \equiv 0.$$

### 3. Convergence theorem

For  $n \in \mathbb{N}$ , compute  $(\mathbf{u}_{n,h}, p_{n,h}) \in \mathbb{V}_h \times \mathbb{Q}_h \subset W_0^{1,r}(\Omega)^d \times L_0^{\tilde{r}}(\Omega)$ , s.t.:

$$\begin{aligned} \int_{\Omega} \mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h})) : \mathbf{D}\mathbf{v}_h \, dx + \mathcal{B}[\mathbf{u}_{n,h}, \mathbf{u}_{n,h}, \mathbf{v}_h] &= \int_{\Omega} p_{n,h} \operatorname{div} \mathbf{v}_h + \mathbf{f} \cdot \mathbf{v}_h \, dx \quad \forall \mathbf{v}_h \in \mathbb{V}_h, \\ \int_{\Omega} q_h \operatorname{div} \mathbf{u}_{n,h} \, dx &= 0 \quad \forall q_h \in \mathbb{Q}_h. \end{aligned}$$

### 3. Convergence theorem

For  $n \in \mathbb{N}$ , compute  $(\mathbf{u}_{n,h}, p_{n,h}) \in \mathbb{V}_h \times \mathbb{Q}_h \subset W_0^{1,r}(\Omega)^d \times L_0^{\tilde{r}}(\Omega)$ , s.t.:

$$\begin{aligned} \int_{\Omega} \mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h})) : \mathbf{D}\mathbf{v}_h \, dx + \mathcal{B}[\mathbf{u}_{n,h}, \mathbf{u}_{n,h}, \mathbf{v}_h] &= \int_{\Omega} p_{n,h} \operatorname{div} \mathbf{v}_h + \mathbf{f} \cdot \mathbf{v}_h \, dx \quad \forall \mathbf{v}_h \in \mathbb{V}_h, \\ \int_{\Omega} q_h \operatorname{div} \mathbf{u}_{n,h} \, dx &= 0 \quad \forall q_h \in \mathbb{Q}_h. \end{aligned}$$

#### Theorem

Suppose that **Case (A):**  $r > \frac{2d}{d+1}$ ; or **Case (B):**  $r > \frac{2d}{d+2}$ ,  $\frac{1}{r} + \frac{1}{r'} = 1$  and  $\mathbf{f} \in W^{-1,\tilde{r}}(\Omega)^d$ . Then, as  $n \rightarrow \infty$ ,  $h \rightarrow 0$ :

$$\begin{aligned} \mathbf{u}_{n,h} &\rightharpoonup \mathbf{u} \quad \text{in } W_0^{1,r}(\Omega)^d, \\ \mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h})) &\rightharpoonup \mathbf{S} \quad \text{in } L^{r'}(\Omega)^{d \times d}, \quad \text{and} \\ p_{n,h} &\rightharpoonup p \quad \text{in } L_0^{\tilde{r}}(\Omega). \end{aligned}$$

Also,  $(\mathbf{u}, p, \mathbf{S}) \in W_0^{1,r}(\Omega)^d \times L_0^{\tilde{r}}(\Omega) \times L^{r'}(\Omega)^{d \times d}$  is such that  $(\mathbf{D}(\mathbf{u}), \mathbf{S}) \in \mathcal{A}$  and

$$\begin{aligned} \int_{\Omega} \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) \cdot \mathbf{v} \, dx + \int_{\Omega} \mathbf{S} : \mathbf{D}(\mathbf{v}) \, dx - \int_{\Omega} p (\operatorname{div} \mathbf{v}) \, dx &= \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \, dx \quad \forall \mathbf{v} \in W_0^{1,\tilde{r}'}(\Omega)^d, \\ \int_{\Omega} (\operatorname{div} \mathbf{u}) q \, dx &= 0 \quad \forall q \in L_0^{r'}(\Omega). \end{aligned}$$

## 4. Convergence proof

**STEP 1.**  $[n \rightarrow \infty, h > 0 \text{ fixed}]$

## 4. Convergence proof

**STEP 1.**  $[n \rightarrow \infty, h > 0 \text{ fixed}]$

The existence of  $\mathbf{u}_{n,h} \in \mathbb{V}_h$  follows by Brouwer's fixed point theorem.

## 4. Convergence proof

**STEP 1.**  $[n \rightarrow \infty, h > 0 \text{ fixed}]$

The existence of  $\mathbf{u}_{n,h} \in \mathbb{V}_h$  follows by Brouwer's fixed point theorem.

Next, note that, taking  $\mathbf{v}_h = \mathbf{u}_{n,h}$ :

$$\|\mathbf{D}(\mathbf{u}_{n,h})\|_{L^r(\Omega)}^r + \|\mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h}))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall n \in \mathbb{N}, \forall h > 0.$$

## 4. Convergence proof

**STEP 1.**  $[n \rightarrow \infty, h > 0 \text{ fixed}]$

The existence of  $\mathbf{u}_{n,h} \in \mathbb{V}_h$  follows by Brouwer's fixed point theorem.

Next, note that, taking  $\mathbf{v}_h = \mathbf{u}_{n,h}$ :

$$\|\mathbf{D}(\mathbf{u}_{n,h})\|_{L^r(\Omega)}^r + \|\mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h}))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall n \in \mathbb{N}, \forall h > 0.$$

Hence, by the Bolzano–Weierstrass thm. and Alaoglu's thm. + Minty's method, as  $n \rightarrow \infty$ ,  $h > 0$  fixed,

$$\begin{aligned} \mathbf{D}(\mathbf{u}_{n,h}) &\rightharpoonup \mathbf{D}(\mathbf{u}_h) && \text{in } L^r(\Omega)^{d \times d}, \\ \mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h})) &\rightharpoonup \mathbf{S}(\mathbf{D}(\mathbf{u}_h)) && \text{in } L^{r'}(\Omega)^{d \times d}, \quad \frac{1}{r} + \frac{1}{r'} = 1. \end{aligned}$$



## 4. Convergence proof

**STEP 1.**  $[n \rightarrow \infty, h > 0 \text{ fixed}]$

The existence of  $\mathbf{u}_{n,h} \in \mathbb{V}_h$  follows by Brouwer's fixed point theorem.

Next, note that, taking  $\mathbf{v}_h = \mathbf{u}_{n,h}$ :

$$\|\mathbf{D}(\mathbf{u}_{n,h})\|_{L^r(\Omega)}^r + \|\mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h}))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall n \in \mathbb{N}, \forall h > 0.$$

Hence, by the Bolzano–Weierstrass thm. and Alaoglu's thm. + Minty's method, as  $n \rightarrow \infty$ ,  $h > 0$  fixed,

$$\begin{aligned} \mathbf{D}(\mathbf{u}_{n,h}) &\rightharpoonup \mathbf{D}(\mathbf{u}_h) && \text{in } L^r(\Omega)^{d \times d}, \\ \mathbf{S}^n(\mathbf{D}(\mathbf{u}_{n,h})) &\rightharpoonup \mathbf{S}(\mathbf{D}(\mathbf{u}_h)) && \text{in } L^{r'}(\Omega)^{d \times d}, \quad \frac{1}{r} + \frac{1}{r'} = 1. \end{aligned}$$

We also pass to the limit  $n \rightarrow \infty$  in the above inequality.

**STEP 2.**  $[h \rightarrow 0]$

It follows from STEP 1 by (weak) lower semicontinuity that

$$\|\mathbf{D}(\mathbf{u}_h)\|_{L^r(\Omega)}^r + \|\mathbf{S}(\mathbf{D}(\mathbf{u}_h))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall h > 0.$$

**STEP 2.**  $[h \rightarrow 0]$

It follows from STEP 1 by (weak) lower semicontinuity that

$$\|D(u_h)\|_{L^r(\Omega)}^r + \|S(D(u_h))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall h > 0.$$

Hence, by Alaoglu's theorem, as  $h \rightarrow 0$ ,

$$\begin{aligned} D(u_h) &\rightharpoonup D(u) && \text{in } L^r(\Omega)^{d \times d}, \\ S(D(u_h)) &\rightharpoonup \overline{S} && \text{in } L^{r'}(\Omega)^{d \times d}, \quad \frac{1}{r} + \frac{1}{r'} = 1. \end{aligned}$$

## STEP 2. $[h \rightarrow 0]$

It follows from STEP 1 by (weak) lower semicontinuity that

$$\|D(u_h)\|_{L^r(\Omega)}^r + \|S(D(u_h))\|_{L^{r'}(\Omega)}^{r'} \leq C \quad \forall h > 0.$$

Hence, by Alaoglu's theorem, as  $h \rightarrow 0$ ,

$$\begin{aligned} D(u_h) &\rightharpoonup D(u) && \text{in } L^r(\Omega)^{d \times d}, \\ S(D(u_h)) &\rightharpoonup \bar{S} && \text{in } L^{r'}(\Omega)^{d \times d}, \quad \frac{1}{r} + \frac{1}{r'} = 1. \end{aligned}$$

**Question:**  $(D(u), \bar{S}) \in \mathcal{A}$  a.e. in  $\Omega$ ?

In other words: is  $\bar{S} = S(D(u))$ ?

**STEP 3.** To show that  $(\boldsymbol{D}(\boldsymbol{u}), \overline{\boldsymbol{S}}) \in \mathcal{A}$  a.e. in  $\Omega$ , we consider

$$a_h := (\boldsymbol{S}(\boldsymbol{D}(\boldsymbol{u}_h)) - \boldsymbol{S}(\boldsymbol{D}(\boldsymbol{u}))) : \boldsymbol{D}(\boldsymbol{u}_h - \boldsymbol{u}).$$

**STEP 3.** To show that  $(\mathbf{D}(\mathbf{u}), \overline{\mathbf{S}}) \in \mathcal{A}$  a.e. in  $\Omega$ , we consider

$$a_h := (\mathbf{S}(\mathbf{D}(\mathbf{u}_h)) - \mathbf{S}(\mathbf{D}(\mathbf{u}))) : \mathbf{D}(\mathbf{u}_h - \mathbf{u}).$$

Using the inequality from STEP 2, we see that

$$\int_{\Omega} |(\mathbf{S}(\mathbf{D}(\mathbf{u}_h)) - \mathbf{S}(\mathbf{D}(\mathbf{u}))) : \mathbf{D}(\mathbf{u}_h - \mathbf{u})| \, dx \leq C.$$

**STEP 3.** To show that  $(D(u), \overline{S}) \in \mathcal{A}$  a.e. in  $\Omega$ , we consider

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u).$$

Using the inequality from STEP 2, we see that

$$\int_{\Omega} |(S(D(u_h)) - S(D(u))) : D(u_h - u)| \, dx \leq C.$$

We apply **Chacon's biting lemma**<sup>1</sup>

---

<sup>1</sup> J.M. Ball & F. Murat, Proc. AMS, Vol. 107, No. 3 (Nov., 1989), pp. 655–663.

**STEP 3.** To show that  $(D(u), \overline{S}) \in \mathcal{A}$  a.e. in  $\Omega$ , we consider

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u).$$

Using the inequality from STEP 2, we see that

$$\int_{\Omega} |(S(D(u_h)) - S(D(u))) : D(u_h - u)| \, dx \leq C.$$

We apply **Chacon's biting lemma**<sup>1</sup> to find a nondecreasing countable sequence of measurable sets  $\Omega_1 \subset \cdots \subset \Omega_k \subset \Omega_{k+1} \subset \cdots \subset \Omega$  such that

$$\lim_{k \rightarrow \infty} |\Omega \setminus \Omega_k| \rightarrow 0$$

and such that for any  $k$  there is a subsequence such that

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u) \quad \text{converges weakly in } L^1(\Omega_k).$$

---

<sup>1</sup> J.M. Ball & F. Murat, Proc. AMS, Vol. 107, No. 3 (Nov., 1989), pp. 655–663.



## Interlude: Lipschitz truncation

## Interlude: Lipschitz truncation

Why is Lipschitz truncation needed?

If  $r < \frac{3d}{d+2}$  then  $\mathbf{u}_h - \mathbf{u} \notin W_0^{1, \tilde{r}'}(\Omega)^d$ , so it is **not** a valid test function.

## Interlude: Lipschitz truncation

Why is Lipschitz truncation needed?

If  $r < \frac{3d}{d+2}$  then  $\mathbf{u}_h - \mathbf{u} \notin W_0^{1,\tilde{r}'}(\Omega)^d$ , so it is **not** a valid test function.

Objective:

Approximate a Sobolev function by a Lipschitz function.

## Interlude: Lipschitz truncation

Why is Lipschitz truncation needed?

If  $r < \frac{3d}{d+2}$  then  $\mathbf{u}_h - \mathbf{u} \notin W_0^{1,\tilde{r}'}(\Omega)^d$ , so it is **not** a valid test function.

Objective:

Approximate a Sobolev function by a Lipschitz function.

Requirement:

Change the Sobolev function only on a set of small measure.

Cannot use convolution as it changes the function on a large set!

## Interlude: Lipschitz truncation

Why is Lipschitz truncation needed?

If  $r < \frac{3d}{d+2}$  then  $\mathbf{u}_h - \mathbf{u} \notin W_0^{1,\tilde{r}'}(\Omega)^d$ , so it is **not** a valid test function.

Objective:

Approximate a Sobolev function by a Lipschitz function.

Requirement:

Change the Sobolev function only on a set of small measure.

Cannot use convolution as it changes the function on a large set!



E. Acerbi & N. Fusco. An approximation lemma for  $W^{1,p}$  functions. In: Material Instabilities in Continuum Mechanics (Edinburgh, 1985-1986), pp. 15, OUP, 1988.

For  $\boldsymbol{v} \in W_0^{1,1}(\Omega)^d$ , consider the Hardy–Littlewood maximal fn.

$$M(\nabla \boldsymbol{v})(x) := \sup_{R>0} \int_{B_R(x)} |\nabla \boldsymbol{v}| \, dy.$$

For  $\boldsymbol{v} \in W_0^{1,1}(\Omega)^d$ , consider the Hardy–Littlewood maximal fn.

$$M(\nabla \boldsymbol{v})(x) := \sup_{R>0} \int_{B_R(x)} |\nabla \boldsymbol{v}| \, dy.$$

Then  $\boldsymbol{v}$  is Lipschitz continuous on

$$\mathcal{G}_\lambda = \mathcal{G}_\lambda(\boldsymbol{v}) = \{x \in \Omega : M(\nabla \boldsymbol{v}) \leq \lambda\}.$$

For  $\mathbf{v} \in W_0^{1,1}(\Omega)^d$ , consider the Hardy–Littlewood maximal fn.

$$M(\nabla \mathbf{v})(x) := \sup_{R>0} \int_{B_R(x)} |\nabla \mathbf{v}| \, dy.$$

Then  $\mathbf{v}$  is Lipschitz continuous on

$$\mathcal{G}_\lambda = \mathcal{G}_\lambda(\mathbf{v}) = \{x \in \Omega : M(\nabla \mathbf{v}) \leq \lambda\}.$$

Indeed,

$$\begin{aligned} |\mathbf{v}(x) - \mathbf{v}(y)| &\leq c (M(\nabla \mathbf{v})(x) + M(\nabla \mathbf{v})(y)) |x - y| \\ &\leq c \lambda |x - y| \quad \forall x, y \in \mathcal{G}_\lambda. \end{aligned}$$



For  $\mathbf{v} \in W_0^{1,1}(\Omega)^d$ , consider the Hardy–Littlewood maximal fn.

$$M(\nabla \mathbf{v})(x) := \sup_{R>0} \int_{B_R(x)} |\nabla \mathbf{v}| \, dy.$$

Then  $\mathbf{v}$  is Lipschitz continuous on

$$\mathcal{G}_\lambda = \mathcal{G}_\lambda(\mathbf{v}) = \{x \in \Omega : M(\nabla \mathbf{v}) \leq \lambda\}.$$

Indeed,

$$\begin{aligned} |\mathbf{v}(x) - \mathbf{v}(y)| &\leq c (M(\nabla \mathbf{v})(x) + M(\nabla \mathbf{v})(y)) |x - y| \\ &\leq c \lambda |x - y| \quad \forall x, y \in \mathcal{G}_\lambda. \end{aligned}$$

Extend  $\mathbf{v}|_{\mathcal{G}_\lambda}$  to  $\mathbf{v}^\lambda \in W_0^{1,\infty}(\Omega)^d$  using Kirszbraun's Extension Theorem.

Good news:  $v^\lambda = v$  on  $\mathcal{G}_\lambda$ ; and  $|\Omega \setminus \mathcal{G}_\lambda| = |v^\lambda \neq v|$  is “small”.

Good news:  $v^\lambda = v$  on  $\mathcal{G}_\lambda$ ; and  $|\Omega \setminus \mathcal{G}_\lambda| = |v^\lambda \neq v|$  is “small”.

Bad news:

Lipschitz truncation of a FE function does **not** belong to the same FE space.

Good news:  $\mathbf{v}^\lambda = \mathbf{v}$  on  $\mathcal{G}_\lambda$ ; and  $|\Omega \setminus \mathcal{G}_\lambda| = |\mathbf{v}^\lambda \neq \mathbf{v}|$  is “small”.

Bad news:

Lipschitz truncation of a FE function does **not** belong to the same FE space.

Consider a locally supported, discretely divergence-preserving, and stable projector  $\Pi_h : W_0^{1,1}(\Omega)^d \rightarrow \mathbb{V}_h \subset W_0^{1,1}(\Omega)^d$ :

$$\int_{\Omega} (\operatorname{div} \mathbf{v}) q_h \, dx = \int_{\Omega} (\operatorname{div} \Pi_h \mathbf{v}) q_h \, dx \quad \forall q_h \in \mathbb{Q}_h.$$

Good news:  $\mathbf{v}^\lambda = \mathbf{v}$  on  $\mathcal{G}_\lambda$ ; and  $|\Omega \setminus \mathcal{G}_\lambda| = |\mathbf{v}^\lambda \neq \mathbf{v}|$  is “small”.

Bad news:

Lipschitz truncation of a FE function does **not** belong to the same FE space.

Consider a locally supported, discretely divergence-preserving, and stable projector  $\Pi_h : W_0^{1,1}(\Omega)^d \rightarrow \mathbb{V}_h \subset W_0^{1,1}(\Omega)^d$ :

$$\int_{\Omega} (\operatorname{div} \mathbf{v}) q_h \, dx = \int_{\Omega} (\operatorname{div} \Pi_h \mathbf{v}) q_h \, dx \quad \forall q_h \in \mathbb{Q}_h.$$

Consider

$$\mathbf{e}_h := \Pi_h(\mathbf{u}_h - \mathbf{u}).$$

# Discrete Lipschitz truncation

# Discrete Lipschitz truncation

## Theorem (Diening, Kreuzer, Süli)

There exists  $\{\lambda_{h,j}\}_j \subset \mathbb{R}$  with  $2^{2^j} \leq \lambda_{h,j} \leq 2^{2^{j+1}-1}$ , such that:

$$e_h^j := \Pi_h e_h^{\lambda_{h,j}} \quad (+ \text{ terms to ensure that } \operatorname{div} e_h^j = 0 \text{ discretely})$$

satisfies:

- ①  $e_h^j \in W_{0}^{1,\infty}(\Omega)^d$  with  $\|\nabla e_h^j\|_{L^\infty(\Omega)} \leq c\lambda_{h,j}$ ;
- ②  $e_h^j \rightarrow 0$  in  $L^\infty(\Omega)^d$  as  $h \rightarrow 0$ ;
- ③  $\nabla e_h^j \rightharpoonup^* 0$  in  $L^\infty(\Omega)^{d \times d}$  as  $h \rightarrow 0$ ;
- ④  $\lambda_{h,j} |e_h^j - e_h|^{\frac{1}{r}} \lesssim 2^{-\frac{j}{r}} \|\nabla e_h\|_{L^r(\Omega)}$ , with  $r > 1$ .



L. Diening, Ch. Kreuzer & E. Süli. *Finite element approximation of steady flows of incompressible fluids with implicit power-law-like rheology*. SIAM J. Numer. Anal., 51(2), 984–1015, 2013.

## STEP 4.

Recall:

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u).$$



#### STEP 4.

Recall:

$$a_h := (S(D(\mathbf{u}_h)) - S(D(\mathbf{u}))) : D(\mathbf{u}_h - \mathbf{u}).$$

For

$$\mathbf{e}_h := \Pi_h(\mathbf{u}_h - \mathbf{u}) \in \mathbb{V}_h$$

let  $\mathbf{e}_h^j$  be the **discrete Lipschitz truncation** of  $\mathbf{e}_h$  at level  $2^{2^j}$ .

## STEP 4.

Recall:

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u).$$

For

$$e_h := \Pi_h(u_h - u) \in \mathbb{V}_h$$

let  $e_h^j$  be the **discrete Lipschitz truncation** of  $e_h$  at level  $2^{2^j}$ .

For a fixed  $j \in \mathbb{N}$ , we have by Hölder's inequality that

$$\begin{aligned} \int_{\Omega} |a_h|^{1/2} \, dx &= \int_{\{e_h^j = e_h\}} |a_h|^{1/2} \, dx + \int_{\{e_h^j \neq e_h\}} |a_h|^{1/2} \, dx \\ &\leq |\Omega|^{1/2} \underbrace{\left( \int_{\{e_h^j = e_h\}} |a_h| \, dx \right)^{1/2}}_{\lesssim 2^{-j/r}} + \underbrace{|\{e_h^j \neq e_h\}|^{1/2}}_{\lesssim 2^{-2^j r}} \left( \int_{\Omega} |a_h| \, dx \right)^{1/2}. \end{aligned}$$

Hence,

$$\limsup_{h \rightarrow 0} \int_{\Omega} |a_h|^{1/2} \, dx \leq c |\Omega|^{1/2} 2^{-j/(2r)} + c 2^{-2^j r/2} \quad \forall j \in \mathbb{N}.$$

Therefore, since  $j$  was arbitrary, we deduce that for a subsequence:

$$a_h := (S(D(u_h)) - S(D(u))) : D(u_h - u) \rightarrow 0 \quad \text{a.e. in } \Omega.$$

## STEP 5.

By Vitali's theorem we then deduce that

$$a_h := (\mathcal{S}(\mathcal{D}(\mathbf{u}_h)) - \mathcal{S}(\mathcal{D}(\mathbf{u}))) : \mathcal{D}(\mathbf{u}_h - \mathbf{u}) \rightarrow 0 \quad \text{strongly in } L^1(\Omega_k).$$

## STEP 5.

By Vitali's theorem we then deduce that

$$a_h := (\mathbf{S}(\mathbf{D}(\mathbf{u}_h)) - \mathbf{S}(\mathbf{D}(\mathbf{u}))) : \mathbf{D}(\mathbf{u}_h - \mathbf{u}) \rightarrow 0 \quad \text{strongly in } L^1(\Omega_k).$$

The weak convergence of  $\mathbf{S}(\mathbf{D}(\mathbf{u}_h))$  to  $\overline{\mathbf{S}}$  and  $\mathbf{D}(\mathbf{u}_h)$  to  $\mathbf{D}(\mathbf{u})$  gives:

$$\lim_{h \rightarrow 0} \int_{\Omega_k} \mathbf{S}(\mathbf{D}(\mathbf{u}_h)) : \mathbf{D}(\mathbf{u}_h) \, dx = \int_{\Omega_k} \overline{\mathbf{S}} : \mathbf{D}(\mathbf{u}) \, dx.$$

## STEP 5.

By Vitali's theorem we then deduce that

$$a_h := (\mathcal{S}(D(\mathbf{u}_h)) - \mathcal{S}(D(\mathbf{u}))) : D(\mathbf{u}_h - \mathbf{u}) \rightarrow 0 \quad \text{strongly in } L^1(\Omega_k).$$

The weak convergence of  $\mathcal{S}(D(\mathbf{u}_h))$  to  $\overline{\mathcal{S}}$  and  $D(\mathbf{u}_h)$  to  $D(\mathbf{u})$  gives:

$$\lim_{h \rightarrow 0} \int_{\Omega_k} \mathcal{S}(D(\mathbf{u}_h)) : D(\mathbf{u}_h) \, dx = \int_{\Omega_k} \overline{\mathcal{S}} : D(\mathbf{u}) \, dx.$$

The assumption that  $\mathcal{A}$  is a maximal monotone  $r$ -graph then implies that

$$(D(\mathbf{u}), \overline{\mathcal{S}}) \in \mathcal{A} \quad \text{a.e. in } \Omega_k, \, k = 1, 2, \dots$$

## STEP 5.

By Vitali's theorem we then deduce that

$$a_h := (\mathcal{S}(D(\mathbf{u}_h)) - \mathcal{S}(D(\mathbf{u}))) : D(\mathbf{u}_h - \mathbf{u}) \rightarrow 0 \quad \text{strongly in } L^1(\Omega_k).$$

The weak convergence of  $\mathcal{S}(D(\mathbf{u}_h))$  to  $\overline{\mathcal{S}}$  and  $D(\mathbf{u}_h)$  to  $D(\mathbf{u})$  gives:

$$\lim_{h \rightarrow 0} \int_{\Omega_k} \mathcal{S}(D(\mathbf{u}_h)) : D(\mathbf{u}_h) \, dx = \int_{\Omega_k} \overline{\mathcal{S}} : D(\mathbf{u}) \, dx.$$

The assumption that  $\mathcal{A}$  is a maximal monotone  $r$ -graph then implies that

$$(D(\mathbf{u}), \overline{\mathcal{S}}) \in \mathcal{A} \quad \text{a.e. in } \Omega_k, \, k = 1, 2, \dots$$

Finally, by a diagonal procedure and  $\lim_{k \rightarrow \infty} |\Omega \setminus \Omega_k| \rightarrow 0$  we deduce that

$$(D(\mathbf{u}), \overline{\mathcal{S}}) \in \mathcal{A} \quad \text{a.e. in } \Omega. \quad \square$$

# Extensions

- T. Tscherpel & E. Süli. *Finite element approximation of unsteady flows of incompressible implicitly constituted fluids*. IMA J. Numer. Anal., 30(2), 2020.
- P. Farrell, A. Gazca-Orozco & E. Süli. *Numerical analysis of unsteady implicitly constituted incompressible fluids: three-field formulation*. SIAM J. Numer. Anal. (58)1, 757–787, 2020.
- P. Farrell, A. Gazca-Orozco & E. Süli. *Finite element approximation and augmented Lagrangian preconditioning for anisothermal implicitly-constituted non-Newtonian flow*. (Submitted). Available from: [arXiv:2011.03024](https://arxiv.org/abs/2011.03024), November 2020.



# Extensions

- T. Tschempel & E. Süli. *Finite element approximation of unsteady flows of incompressible implicitly constituted fluids*. IMA J. Numer. Anal., 30(2), 2020.
- P. Farrell, A. Gazca-Orozco & E. Süli. *Numerical analysis of unsteady implicitly constituted incompressible fluids: three-field formulation*. SIAM J. Numer. Anal. (58)1, 757–787, 2020.
- P. Farrell, A. Gazca-Orozco & E. Süli. *Finite element approximation and augmented Lagrangian preconditioning for anisothermal implicitly-constituted non-Newtonian flow*. (Submitted). Available from: [arXiv:2011.03024](https://arxiv.org/abs/2011.03024), November 2020.


## Note

All numerical simulations shown on the next slides were performed by Alexei Gazca-Orozco (Oxford (now at Erlangen–Nürnberg)).


Finite element software: `firedrake`

# Unsteady Navier–Stokes/Euler activated fluid

$$\mathbf{D} = \begin{cases} \begin{cases} \delta_s \frac{\mathbf{S}}{|\mathbf{S}|} + \frac{1}{2\mu} \mathbf{S} & \text{if } |\mathbf{D}| > \tau_y \\ \mathbf{S} = \mathbf{0} & \text{if } |\mathbf{D}| \leq \tau_y \\ \frac{1}{2\mu} \mathbf{S} & \text{otherwise} \end{cases} & \text{if } (x - \frac{1}{2})^2 + (y - \frac{1}{2})^2 \leq (\frac{9}{64})^2, \\ & \text{otherwise} \end{cases}$$



(a) Navier–Stokes fluid



(b) Navier–Stokes/Euler activated fluid

$$\Omega = (0, 1)^2, \quad \delta_s = 2.5, \quad \mu = 0.5, \quad \tau_m = 5 \times 10^{-6}, \quad \mathbb{P}_2 \times \mathbb{P}_1 \times \mathbb{P}_{-1}.$$

## 5. Nonmonotone constitutive laws

The problem under consideration in  $Q := (0, T) \times \Omega$  is:

$$\begin{aligned} \mathbf{u}_t - \operatorname{div}(\mathbf{S} - \mathbf{u} \otimes \mathbf{u}) + \nabla p &= \mathbf{f} && \text{in } Q, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } Q, \\ \mathbf{D}(\mathbf{u}) &= \mathcal{D}(\mathbf{S}) && \text{in } Q, \\ \mathbf{u} &= \mathbf{0} && \text{in } (0, T) \times \partial\Omega, \\ \mathbf{u}(0, \cdot) &= \mathbf{u}_0(\cdot) && \text{in } \Omega. \end{aligned}$$

Here  $\mathcal{D} : \mathbb{R}_{\operatorname{sym},0}^{d \times d} \rightarrow \mathbb{R}_{\operatorname{sym},0}^{d \times d}$  is a continuous function that satisfies:

- $\mathcal{D}(\mathbf{0}) = \mathbf{0}$ ;
- There exists an  $r \in (1, \infty)$  and constants  $c, C > 0$  such that

$$\begin{aligned} c(|\mathbf{D}|^r + |\mathbf{S}|^{r'}) &\leq \mathbf{S} : \mathbf{D} \quad \text{whenever } \mathbf{D} = \mathcal{D}(\mathbf{S}), \\ |\mathcal{D}(\mathbf{S})| &\leq C(1 + |\mathbf{S}|^{r'-1}) \quad \forall \mathbf{S} \in \mathbb{R}_{\operatorname{sym},0}^{d \times d}. \end{aligned}$$

## 5. Nonmonotone constitutive laws

The problem under consideration in  $Q := (0, T) \times \Omega$  is:

$$\begin{aligned} \mathbf{u}_t - \operatorname{div}(\mathbf{S} - \mathbf{u} \otimes \mathbf{u}) + \nabla p &= \mathbf{f} && \text{in } Q, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } Q, \\ \mathbf{D}(\mathbf{u}) &= \mathcal{D}(\mathbf{S}) && \text{in } Q, \\ \mathbf{u} &= \mathbf{0} && \text{in } (0, T) \times \partial\Omega, \\ \mathbf{u}(0, \cdot) &= \mathbf{u}_0(\cdot) && \text{in } \Omega. \end{aligned}$$

Here  $\mathcal{D} : \mathbb{R}_{\operatorname{sym},0}^{d \times d} \rightarrow \mathbb{R}_{\operatorname{sym},0}^{d \times d}$  is a continuous function that satisfies:

- $\mathcal{D}(\mathbf{0}) = \mathbf{0}$ ;
- There exists an  $r \in (1, \infty)$  and constants  $c, C > 0$  such that

$$\begin{aligned} c(|\mathbf{D}|^r + |\mathbf{S}|^{r'}) &\leq \mathbf{S} : \mathbf{D} \quad \text{whenever } \mathbf{D} = \mathcal{D}(\mathbf{S}), \\ |\mathcal{D}(\mathbf{S})| &\leq C(1 + |\mathbf{S}|^{r'-1}) \quad \forall \mathbf{S} \in \mathbb{R}_{\operatorname{sym},0}^{d \times d}. \end{aligned}$$

David & Filip (2004), Galindo-Rosales et al. (2011), Fardin et al. (2012), Divoux et al. (2016).

The nonmonotone response seems crucial in modelling complex non-Newtonian phenomena such as shear banding.

## 5. Nonmonotone constitutive laws

The problem under consideration in  $Q := (0, T) \times \Omega$  is:

$$\begin{aligned} \mathbf{u}_t - \operatorname{div}(\mathbf{S} - \mathbf{u} \otimes \mathbf{u}) + \nabla p &= \mathbf{f} && \text{in } Q, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } Q, \\ \mathbf{D}(\mathbf{u}) &= \mathcal{D}(\mathbf{S}) && \text{in } Q, \\ \mathbf{u} &= \mathbf{0} && \text{in } (0, T) \times \partial\Omega, \\ \mathbf{u}(0, \cdot) &= \mathbf{u}_0(\cdot) && \text{in } \Omega. \end{aligned}$$

Here  $\mathcal{D} : \mathbb{R}_{\operatorname{sym},0}^{d \times d} \rightarrow \mathbb{R}_{\operatorname{sym},0}^{d \times d}$  is a continuous function that satisfies:

- $\mathcal{D}(\mathbf{0}) = \mathbf{0}$ ;
- There exists an  $r \in (1, \infty)$  and constants  $c, C > 0$  such that

$$\begin{aligned} c(|\mathbf{D}|^r + |\mathbf{S}|^{r'}) &\leq \mathbf{S} : \mathbf{D} \quad \text{whenever } \mathbf{D} = \mathcal{D}(\mathbf{S}), \\ |\mathcal{D}(\mathbf{S})| &\leq C(1 + |\mathbf{S}|^{r'-1}) \quad \forall \mathbf{S} \in \mathbb{R}_{\operatorname{sym},0}^{d \times d}. \end{aligned}$$

**Notation:** For a probability measure  $\nu \in \mathcal{P}(\mathbb{R}_{\operatorname{sym},0}^{d \times d})$  we write

$$\langle \nu, f(\tilde{\mathbf{S}}) \rangle := \int_{\mathbb{R}_{\operatorname{sym},0}^{d \times d}} f(\mathbf{S}) \, d\nu(\mathbf{S}) \quad \forall f \in C_0(\mathbb{R}_{\operatorname{sym},0}^{d \times d}).$$

## Definition

A couple  $(\mathbf{S}, \mathbf{u})$  is called a **Young-measure solution** of the problem if

$$(a) \quad \mathbf{S} \in L^{r'}_{\text{sym},0}(Q)^{d \times d}, \quad \mathbf{u} \in L^r(0, T; W^{1,r}_{0,\text{div}}(\Omega)^d) \cap C_w([0, T]; L^2_{\text{div}}(\Omega)^d), \\ \mathbf{u}_t \in L^{\tilde{r}'}(0, T; (W^{1,\tilde{q}}_{0,\text{div}}(\Omega)^d)') \quad \text{satisfy the eqn.}$$

$$-\int_0^T \int_{\Omega} \mathbf{u} \cdot \mathbf{v}_t + \int_0^T \int_{\Omega} (\mathbf{S} - \mathbf{u} \otimes \mathbf{u}) : \mathbf{D}(\mathbf{v}) = \int_0^T \langle \mathbf{f}, \mathbf{v} \rangle + \int_{\Omega} \mathbf{u}_0 \cdot \mathbf{v}(0, \cdot) \\ \forall \mathbf{v} \in C^\infty_{0,\text{div}}((-T, T) \times \Omega)^d;$$

(b) There exists a Young-measure  $\{\nu_{\mathbf{z}}\}_{\mathbf{z} \in Q} \subset \mathcal{P}(\mathbb{R}^{d \times d}_{\text{sym},0})$  such that

$$\mathbf{S} = \langle \nu_{\bullet}, \tilde{\mathbf{S}} \rangle, \quad \mathbf{D}(\mathbf{u}) = \langle \nu_{\bullet}, \mathcal{D}(\tilde{\mathbf{S}}) \rangle \quad \text{a.e. on } Q.$$


## Theorem

Suppose  $r > \frac{2d}{d+2}$ ,  $\mathbf{u}_0 \in L^2_{\text{div}}(\Omega)^d$ , and  $\mathbf{f} \in L^{r'}(0, T; W^{-1, r'}(\Omega)^d)$ . Then, the numerical method admits a solution  $(\mathbf{S}_j^{m,n}, \mathbf{u}_j^{m,n}) \in \Sigma^n \times V^n_{\text{div}}$ ,  $j \in \{1, \dots, T/\tau_m\}$ ,  $m \in \mathbb{N}$ . Furthermore, there is a subsequence (not indicated) such that the piecewise constant in time interpolants satisfy

$$\begin{aligned} \overline{\mathbf{S}}^{m,n} &\rightharpoonup \mathbf{S} && \text{weakly in } L^{r'}_{\text{sym},0}(Q)^{d \times d}, \\ \overline{\mathbf{u}}^{m,n} &\rightharpoonup \mathbf{u} && \text{weakly}^* \text{ in } L^\infty(0, T; L^2_{\text{div}}(\Omega)^d), \\ \overline{\mathbf{u}}^{m,n} &\rightharpoonup \mathbf{u} && \text{weakly in } L^r(0, T; W^{1,r}_{0,\text{div}}(\Omega)^d), \\ \overline{\mathbf{u}}^{m,n} &\rightarrow \mathbf{u} && \text{strongly in } L^2(Q)^d, \end{aligned}$$

where  $(\mathbf{S}, \mathbf{u})$  is a Young-measure solution of the problem.

## An example



$$\mathcal{D}(S) = \alpha_1(1 + \beta_1|S|^2)^{\frac{2-q}{2(q-1)}} S + \alpha_2(1 + \beta_2|S|^2)^{\frac{2-r}{2(r-1)}} S$$



where  $q \in (-\infty, -1)$ ,  $r \in (1, \infty)$ , and  $\alpha_1, \alpha_2, \beta_1, \beta_2 > 0$ .


We shall plot the *apparent viscosity*  $|S|/|D|$ .




A. Janečka, J. Málek, V. Pruša & G. Tierra, Acta Mech., 230, 729–747 (2019).










(e) Monotone  $\uparrow$  part 1 of the graph



(f) Monotone  $\uparrow$  part 2 of the graph

Numerical simulations by Alexei Gazca (Oxford/Erlangen–Nürnberg)  
finite element software: `firedrake`

## 6. Conclusions and open problems

- We have shown the convergence of a large class of finite element methods for implicitly constituted incompressible fluid flow models in the steady and unsteady cases, with both monotone and nonmonotone constitutive laws.

## 6. Conclusions and open problems

- We have shown the convergence of a large class of finite element methods for implicitly constituted incompressible fluid flow models in the steady and unsteady cases, with both monotone and nonmonotone constitutive laws.
- If the solution to the PDE is unique, then the sequence of approximations converges to the unique solution.

## 6. Conclusions and open problems

- We have shown the convergence of a large class of finite element methods for implicitly constituted incompressible fluid flow models in the steady and unsteady cases, with both monotone and nonmonotone constitutive laws.
- If the solution to the PDE is unique, then the sequence of approximations converges to the unique solution.
- There are significant analytical and computational challenges:

## 6. Conclusions and open problems

- We have shown the convergence of a large class of finite element methods for implicitly constituted incompressible fluid flow models in the steady and unsteady cases, with both monotone and nonmonotone constitutive laws.
- If the solution to the PDE is unique, then the sequence of approximations converges to the unique solution.
- There are significant analytical and computational challenges:
  - ▶ strong nonlinearities;
  - ▶ potential nonuniqueness of solutions;
  - ▶ the approximation of measure-valued solutions.

## 6. Conclusions and open problems

- We have shown the convergence of a large class of finite element methods for implicitly constituted incompressible fluid flow models in the steady and unsteady cases, with both monotone and nonmonotone constitutive laws.
- If the solution to the PDE is unique, then the sequence of approximations converges to the unique solution.
- There are significant analytical and computational challenges:
  - ▶ strong nonlinearities;
  - ▶ potential nonuniqueness of solutions;
  - ▶ the approximation of measure-valued solutions.