

# On Stability of Newton Schulz Iterations in an Approximate Algebra

Matt Challacombe\* and Nicolas Bock†

Theoretical Division, Los Alamos National Laboratory

## I. INTRODUCTION

In many areas of application, finite correlations lead to matrices with decay properties. Matrix decay involves an approximate (perhaps bounded  $\| \cdot \|$ ) inverse relationship between matrix elements and a related distance; this may be a simple inverse exponential relationship between elements and the Cartesian distance between support functions, or it may involve a generalized distance, *e.g.* a statistical measure between strings. In electronic structure, correlations manifest in decay properties of the gap shifted matrix sign function, as projector of the effective Hamiltonian (Fig. 1). More broadly, matrix decay properties may corespond to statistical matrices [1–5], including learned correlations in a generalized, non-orthogonal metric  $\| \cdot \|$ . More broadly still, problems with local, non-orthogonal support are often solved with congruential transformations of the matrix inverse square root [6, 7] or a related factorization [5]; these transformations correlate support with a representation independent linear form, *eg.* of the generalized eigenproblem. Interestingly, the matrix sign function and the matrix inverse square root function are related by Higham’s identity:

$$\text{sign} \left( \begin{bmatrix} 0 & s \\ I & 0 \end{bmatrix} \right) = \begin{bmatrix} 0 & s^{1/2} \\ s^{-1/2} & 0 \end{bmatrix}. \quad (1)$$

A complete overview of matrix function theory and computation is given in Higham’s enjoyable reference [8].

A well conditioned matrix  $s$  may often correspond to matrix sign and inverse square root functions with rapid exponential decay, and be amenable to the sparse matrix approximation  $\bar{s} = s + \epsilon_\tau^s$ , where  $\epsilon_\tau^s$  is the error introduced according to some criteria  $\tau$ . This criteria might be a drop-tolerance,  $\epsilon_\tau^s = \{-s_{ij} * \hat{e}_i \mid |s_{ij}| < \tau\}$ , a radial cutoff,  $\epsilon_\tau^s = \{-s_{ij} * \hat{e}_i \mid \|r_i - r_j\| > \tau\}$ , or some other approach to truncation, perhaps involving a sparsity pattern chosen *a priori*. Then, conventional computational kernels may be employed, such as the sparse general matrix-matrix multiply (SpGEMM) [9–12], yielding fast solutions for multiplication rich iterations and a modulated fill in. These and related incomplete/inexact approaches to the computation of sparse approximate matrix functions often lead to  $\mathcal{O}(n)$  algorithms, finding wide use in technologically important preconditioning schemes, the

information sciences, electronic structure and many other disciplines. Comprehensive surveys of these methods in the numerical linear algebra is given by Benzi [13? ]. See also Bowler [14] and Benzi [15] for a complete development of these methods in electronic structure.




FIG. 1: Examples from electronic structure of decay for the spectral projector (gap shifted sign function) with respect to local (atomic) support. Shown is decay for systems with correlations that are short (insulating water), medium (semi-conducting 4,3 nanotube), and long (metallic 3,3 nanotube) ranged, from exponential (insulating) to algebraic (metallic).

Incompleteness -; sparse approximations dense problems, uses conventional sparse infrastructure, second order errors in matrix multiplication. Often adhoc.

$$\tilde{a} \cdot \tilde{b} = a \cdot b + \delta a \cdot b + a \cdot \delta b + \delta a \cdot \delta b$$

The variations do not express in the overall context of the product. Because the error in the incomplete case is additive, the

For example,  $a$  may be small, but  $\delta a \cdot b$  large, leading extra work. Also, once a truncation error is committed, it is encountered in all subsequent steps; it becomes difficult to impossible to manage error flows of differing magnitude in complex maps.

For extended quasi-degenerate correlations, these matrix functions may encounter ill-conditioning, and associated slow rates of decay. For extremely slow decay, maybe even oscillatory, low order algebraic decay, methods that compression.... For fast decay,

correlation and the support

Also, matrices with decay arise from the application of . Generally, ill-conditioning is associated with slower decay,

\*Electronic address: matt.challacombe@freeon.org; URL: <http://www.freeon.org>

†Electronic address: nicolasbock@freeon.org; URL: <http://www.freeon.org>

Decay principles, often very sparse but very ill-conditioned problems.

### A. Retaining the Eigenspace

Gradients lack convergence properties Iteration without orig drives away from basis NS has both. Difference between scalar iteration, Higham page 92.

### B. Approximate Algebra as $N$ -Body Problem

SpAMM is the recursive Cauchy-Schwarz occlusion product  $\otimes_\tau$  on matrix quadrees

$$\mathbf{a}^i = \begin{bmatrix} \mathbf{a}_{00}^{i+1} & \mathbf{a}_{01}^{i+1} \\ \mathbf{a}_{10}^{i+1} & \mathbf{a}_{11}^{i+1} \end{bmatrix} \quad (2)$$

$$\mathbf{a}^i \otimes_\tau \mathbf{b}^i = \begin{cases} \emptyset & \text{if } \|\mathbf{a}^i\| \|\mathbf{b}^i\| < \tau \\ \mathbf{a}^i \cdot \mathbf{b}^i & \text{if (i = leaf)} \\ \begin{bmatrix} \mathbf{a}_{00}^{i+1} \otimes_\tau \mathbf{b}_{00}^{i+1} + \mathbf{a}_{01}^{i+1} \otimes_\tau \mathbf{b}_{10}^{i+1} & \mathbf{a}_{00}^{i+1} \otimes_\tau \mathbf{b}_{01}^{i+1} + \mathbf{a}_{01}^{i+1} \otimes_\tau \mathbf{b}_{11}^{i+1} \\ \mathbf{a}_{10}^{i+1} \otimes_\tau \mathbf{b}_{00}^{i+1} + \mathbf{a}_{11}^{i+1} \otimes_\tau \mathbf{b}_{10}^{i+1} & \mathbf{a}_{10}^{i+1} \otimes_\tau \mathbf{b}_{01}^{i+1} + \mathbf{a}_{11}^{i+1} \otimes_\tau \mathbf{b}_{11}^{i+1} \end{bmatrix} & \text{else} \end{cases} \quad (3)$$

database orientation, Cauchy sch Approximate Algebra, SpAMM Cauchy Schwarz occlusion, n-body approach to numerical linear algebra, first order errors in matrix multiplication. Based on Cauchy Schwarz inequality.

$$\mathbf{a} \otimes_\tau \mathbf{b} = \mathbf{a} \cdot \mathbf{b} + \Delta_\tau^{a \cdot b} \quad (4)$$

where  $\Delta_\tau^{a \cdot b}$  is a deterministic (assymetric) first order variation cooresponding to the branch pattern set by Cauchy-Schwarz occlusion, with length  $\|\Delta_\tau^{a \cdot b}\| \leq \tau \|\mathbf{a}\| \|\mathbf{b}\|$ . The operator  $\otimes_\tau$  leads to a non-associative algebra with Lie bracket

$$[\mathbf{a}, \mathbf{b}]_\tau = \mathbf{a} \otimes_\tau \mathbf{b} - \mathbf{b} \otimes_\tau \mathbf{a} = [\mathbf{a}, \mathbf{b}] + \Delta_\tau^{a \cdot b} - \Delta_\tau^{b \cdot a}. \quad (5)$$

determined by the occlusion field. Our challenge is to master the error flows of these occlusion fields under iteration, for ill-conditioned problems and with permissive values of  $\tau$ .

## II. NEWTON SHULZ ITERATION

### A. Idempotence

### B. The Scaled Map

### C. Alternative Formulations

dual, stabilized and naive

## III. OCCLUSION FLOWS

$\delta \mathbf{x}_k$  and  $\delta \mathbf{z}_k$  arrive from iteration with  $\otimes_\tau$ , and are deterministic flows away from the manifold of  $\mathbf{s}$  determined by sensitivity of the NS iteration to these numerical inputs.

$$\delta \mathbf{x}_k^{\text{naiv}} = \delta \tilde{\mathbf{z}}_k \cdot \mathbf{s} \cdot \tilde{\mathbf{z}}_k + \tilde{\mathbf{z}}_k \cdot \mathbf{s} \cdot \delta \tilde{\mathbf{z}}_k \quad (6)$$

$$\delta \mathbf{x}_k^{\text{dual}} = \delta \tilde{\mathbf{y}}_k \cdot \tilde{\mathbf{z}}_k + \tilde{\mathbf{y}}_k \cdot \delta \tilde{\mathbf{z}}_k \quad (7)$$

$$\tilde{\mathbf{x}}_k = f[\tilde{\mathbf{z}}_{k-1}, \tilde{\mathbf{x}}_{k-1}] \quad (8)$$

$$= \mathbf{m}[\tilde{\mathbf{x}}_{k-1}] \cdot \tilde{\mathbf{z}}_{k-1}^\dagger \cdot \mathbf{s} \cdot \tilde{\mathbf{z}}_{k-1} \cdot \mathbf{m}[\tilde{\mathbf{x}}_{k-1}]$$

$$\delta \mathbf{x}_k = f_{\delta \mathbf{z}_{k-1}} \|\delta \mathbf{z}_{k-1}\| + f_{\delta \mathbf{x}_{k-1}} \|\delta \mathbf{x}_{k-1}\| + \mathcal{O}(\tau^2) \quad (9)$$

generalized Gateaux differential

$$\begin{aligned} f_{\delta \mathbf{z}_{k-1}} &= \lim_{\tau \rightarrow 0} \frac{f[\mathbf{z}_{k-1} + \tau \delta \tilde{\mathbf{z}}_{k-1}, \tilde{\mathbf{x}}_{k-1}] - f[\mathbf{z}_{k-1}, \tilde{\mathbf{x}}_{k-1}]}{\tau} \\ &= L_{\tilde{\mathbf{x}}_k}(\tilde{\mathbf{z}}_k, \delta \tilde{\mathbf{z}}_{k-1}) \end{aligned} \quad (10)$$

$$\begin{aligned} f_{\delta \mathbf{x}_{k-1}} &= \lim_{\tau \rightarrow 0} \frac{f[\tilde{\mathbf{z}}_{k-1}, \mathbf{x}_{k-1} + \tau \delta \tilde{\mathbf{x}}_{k-1}] - f[\tilde{\mathbf{z}}_{k-1}, \mathbf{x}_{k-1}]}{\tau} \\ &= L_{\tilde{\mathbf{x}}_k}(\tilde{\mathbf{z}}_k, \delta \tilde{\mathbf{x}}_{k-1}) \end{aligned} \quad (11)$$

$$L_{\tilde{\mathbf{x}}_k}(\tilde{\mathbf{z}}_k, \delta \hat{\mathbf{x}}_{k-1}) = \delta \hat{\mathbf{x}}_{k-1}^\dagger \cdot \mathbf{m}'[\mathbf{x}_{k-1}] \cdot \{\tilde{\mathbf{z}}_{k-1}^\dagger \cdot \mathbf{s} \cdot \tilde{\mathbf{z}}_k\} \\ + \{\tilde{\mathbf{z}}_k^\dagger \cdot \mathbf{s} \cdot \tilde{\mathbf{z}}_{k-1}\} \cdot \mathbf{m}'[\mathbf{x}_{k-1}] \cdot \delta \hat{\mathbf{x}}_{k-1} \quad (12)$$

$$L_{\tilde{\mathbf{x}}_k}(\tilde{\mathbf{z}}_k, \delta \hat{\mathbf{z}}_{k-1}) = \{\mathbf{m}[\mathbf{x}_{k-1}] \cdot \delta \hat{\mathbf{z}}_{k-1}^\dagger \cdot \mathbf{s}\} \cdot \tilde{\mathbf{z}}_k \\ + \tilde{\mathbf{z}}_k^\dagger \cdot \{\mathbf{s} \cdot \delta \hat{\mathbf{z}}_{k-1} \cdot \mathbf{m}[\mathbf{x}_{k-1}]\} \quad (13)$$

$$\{\tilde{\mathbf{z}}_k^\dagger \cdot \mathbf{s} \cdot \tilde{\mathbf{z}}_{k-1}\} \rightarrow \mathbf{p}_+[s] \quad (14)$$

$$\{\mathbf{s} \cdot \delta \hat{\mathbf{z}}_{k-1} \cdot \mathbf{m}[\mathbf{x}_{k-1}]\} \rightarrow \mathbf{n}[s] \quad (15)$$

#### IV. BASIS SET ILL-CONDITIONING IN ELECTRONIC STRUCTURE

##### A. 3,3 carbon nanotube with diffuse *sp*-function

double exponential (Fig.)

##### B. Water with triple zeta and double polarization

Here's looking at you Jurg...

#### V. IMPLEMENTATION

##### A. Methods

FP, F08, OpenMP 4.0

##### B. A Modified NS Map


##### C. $\delta \mathbf{x}_k$ and $\delta \mathbf{x}_k$ channels

tau= Figure showing channels etc.

##### D. Convergence

Map switching and etc based on TrX

FIG. 2: equation...



#### VI. EXPERIMENTS

##### A. Occlusion Flows


##### B. Comments

$$\delta \mathbf{z}_{k-1} \approx \Delta_{\tau}^{\tilde{\mathbf{z}}_{k-2} \cdot \mathbf{m}[\tilde{\mathbf{x}}_{k-2}]} + \mathbf{z}_{k-2} \cdot \mathbf{m}'[\tilde{\mathbf{x}}_{k-2}] \cdot \delta \mathbf{x}_{k-2} \\ + \delta \mathbf{z}_{k-2} \cdot \mathbf{m}[\tilde{\mathbf{x}}_{k-2}] \quad (16)$$

$$\|\delta \mathbf{z}_{k-1}\| \lesssim \|\mathbf{z}_{k-2}\| ( \tau \|\mathbf{m}[\tilde{\mathbf{x}}_{k-2}]\| \\ + \|\delta \mathbf{x}_{k-2}\| \|\mathbf{m}'[\tilde{\mathbf{x}}_{k-2}]\| ) \quad (17)$$

$$\|\mathbf{z}_k\| \rightarrow \sqrt{\kappa(s)} \quad (18)$$

FIG. 3: equation...



### C. Scaling

### D. Comments

Pictures of the spamm structure

## VII. CONCLUSION

- 
- [1] O. Penrose and J. Lebowitz, Commun. Math. Phys. **184** (1974).
  - [2] J. Voit, *The Statistical Mechanics of Financial Markets*, Theoretical and Mathematical Physics (Springer Berlin Heidelberg, 2006), ISBN 9783540262893, URL [https://books.google.com/books?id=6zUlh\\\_TkWSwC](https://books.google.com/books?id=6zUlh\_TkWSwC).
  - [3] L. Anselin, Int. Reg. Sci. Rev. **26**, 153 (2003), ISSN 01600176, URL <http://irx.sagepub.com/cgi/doi/10.1177/0160017602250972>.
  - [4] J. Hardin, S. R. Garcia, and D. Golan, Ann. Appl. Stat. **7**, 1733 (2013), ISSN 1932-6157, arXiv:1106.5834v4.
  - [5] I. Krishtal, T. Strohmer, and T. Wertz, Found. Comput. ... (2013), ISSN 1615-3375.
  - [6] P. O. Lowdin, Advances in Physics **5**, 1 (1956).
  - [7] A. R. Naidu, p. 1 (2011), 1105.3571.
  - [8] N. J. Higham, *Functions of Matrices* (Society for Industrial & Applied Mathematics, 2008).
  - [9] F. G. Gustavson, ACM Transactions on Mathematical Software (TOMS) **4**, 250 (1978).
  - [10] S. Toledo, IBM J. Res. Dev. **41**, 711 (1997).
  - [11] M. Challacombe, Comput. Phys. Commun. **128**, 93 (2000).
  - [12] D. R. Bowler and T. M. and M. J. Gillan, Comp. Phys. Comm. **137**, 255 (2000).
  - [13] M. Benzi, J. Comput. Phys. **182**, 418 (2002).
  - [14] D. R. Bowler and T. Miyazaki, Reports Prog. Phys. **75**, 36503 (2012).
  - [15] M. Benzi, P. Boito, and N. Razouk, SIAM Rev. **55**, 3 (2013).