Damping per (cell, mass), the inputs for Paper 2 (sec. 4.8.6.4; methods sec. 13): Gamma and Q from the item-4a fit of the seed-averaged ACF, block B

alpha = 0.5 · alpha = 1 · alpha = 2 · alpha = 3 · alpha = 5 · alpha = 7.5 · alpha = 10 · alpha = 15 · alpha = 20 · Q = 5 (the low-Q flag) · open: left out by the group rule, or a stationarity flag

N = 100 · N = 400 · N = 900 · N = 1600

dilute gas-like fluid · dense isotropic fluid · stiff dense isotropic fluid

ideal gas limit · fluid-hexatic coexist · hexatic

Gamma = 2 / tau_r [1 / sigma-time]

Q_k = pi nu_d-bar tau_r

packing fraction eta

data: 261015_s1_fig6_Gamma_and_Q_per_mass_vs_eta_N100-1600.csv
script: cluster/gen3_stage1_261012/s1_figures_261015.py (s1_verdict_261012.analyse on experiments_gen3_stage1_koa_261012, all six groups, the full report's numbers)
N = 100, 400, 900, 1600 disks, N/2 per side; disk radius 0.5; divider thickness 0.05; H = 10 sqrt(N/100) = 10, 20, 30, 40; L0 = N pi / (8 H eta) per side (exact box); gen3 engine, one build (f72d764); 25 seeds per (cell, mass), 10 in block S