δ_M per engine: Report G2 (POST HOC, INFORMATION) and Test R (INFORMATION)

N = 100 disks (50 per compartment), radius 0.5, L0 = 10, H = 10, η = π/8 (dense isotropic fluid); one KOA binary, engine-gen3 a643bc7 (target koa),
run as gen2 and gen3; Test G 400 trajectories per engine and mass (M = 2000: 100), Test R 3600

(a) Report G2 (Test G on KOA): gen2 (c_ref: weighted mean over M ≥ 300)



(b) Report G2 (Test G on KOA): gen3 (c_ref: weighted mean over M ≥ 300)



(c) Test R (its information rows): gen2 (against c_300)



(d) Test R (its information rows): gen3 (against c_300)



c_M = nu_M / x_M, x_M = K(alpha)/(2 pi L_eff); delta_M = c_M / c_ref - 1. The references differ by row: Report G2's c_ref is the inverse-variance weighted mean of c_M over M >= 300 (300, 1500, 2000) of the same engine and estimator; Test R's is c_300, its one mass >= 300.
SE: delete-one-group jackknife (groups of 10), the reference's own error included; error bars 95 % (1.96 SE). Values as printed in Report G2 (main 76f1f115) and in Test R's verdict output (main 7ed5e102, information rows).
Test R at M = 50, nu_d: gen2 -0.389 +- 0.029 %, gen3 -0.428 +- 0.029 %: gen3 - gen2 = -0.039 +- 0.041 % (Report R1).
tables: hspist3/experiments_m1_figures_261011/tables/ (main 752d5754); script: hspist3/experiments_m1_figures_261011/m1_figures_v2_261011.py (main 204c322d)