δ_M per engine from Report G2 (Test G on KOA): POST HOC, INFORMATION

N = 100 disks (50 per compartment), radius 0.5, L0 = 10, H = 10, η = π/8; one KOA binary, engine-gen3 a643bc7 (target koa),
run as gen2 (a) and gen3 (b); 400 trajectories per engine and mass (M = 2000: 100)



c_M = nu_M / x_M with x_M = K(alpha)/(2 pi L_eff) (tests_20260913.x_of, L_eff = 8.975); delta_M = c_M / c_ref - 1, c_ref the inverse-variance weighted mean of c_M over M >= 300 (300, 1500, 2000) of the same engine and estimator.
Estimators: argmax nu (registered); nu_d (sec. 4.4.13 item 4a fit per trajectory); nu_0 = sqrt(nu_d^2 + (1/(2 pi tau_r))^2) (methods sec. 16). SE: delete-one-group jackknife over the trajectories (groups of 10; M = 2000: 5), the reference's own error included.
Error bars 95 % (1.96 SE). Values exactly as Report G2 prints them (sec. 4.7.56, main 76f1f115). Methods sec. 16's -0.424 +- 0.078 % is against a nine-mass reference, not against G2's M >= 300 reference.
tables: hspist3/experiments_m1_figures_261011/tables/ (main b7ac7c59); script: hspist3/experiments_m1_figures_261011/m1_figures_261011.py (main 973c1839)