PREVIEW: c_s(η) at N = 100 by the canonical estimator E0
(INFORMATION, no verdict; never part of stage 1, never in a paper)

N = 100 disks (50 per compartment), radius 0.5, H = 10, wall thickness 0.05; the nine A1 v2 masses, 25 seeds each; the canonical A1 v2 command (methods sec. 8)
the preview's cells, recorded eta (L0): 0.052360 (74.9998), 0.112200 (34.9999), 0.196350 (20.0000), 0.261799 (15.0000), 0.392699 (10.0000), 0.523599 (7.5000), 0.590001 (6.6559)
the axis: the box-corrected eta of methods sec. 14; A1 v2's cells as in its canonical table









E0: per (eta, M) the mean argmax frequency over the seeds; per cell the through-origin slope over the nine masses against x_M = K(alpha)/(2 pi L_eff,true), with the box-truncation correction of methods sec. 14 (eta = eta_rec L0/(L0 - delta/2), L_eff,true = L0 - 2r - t/2 - delta/2); error bar c_s_err_scaled = c_s_err max(1, sqrt(chi2_red)) (PDG scale factor). Theory: the adiabatic mapping c_s^2 = (kT/m)(Z + eta Z' + Z^2) for every EOS; KR compared only for eta <= 0.69. Residuals: KR of the current module at each table's own eta.