The solid: Z per compartment, 0° and turned orientation, in the cells of P4–P6: PILOT (no verdict, no paper use)



Z_c: the dt-weighted mean of each compartment's virial blocks in Q4 = [1.5e4, 2e4) sigma-time of a 2e4 sigma-time hold with the divider held (M = 300); for 'P4 A, first quarter' its window [0, 5000), before any compartment turned. Two compartments per run, 6 seeds per start.
Marker = the orientation MEASURED in that window (P6: psi6 per compartment, every seed; P4, P5: snapshots of seeds 0 and 1 only), not the start's: at N = 1600, eta = 0.740 every compartment of the 0° start reads +-30° in Q4 (it turned).
Grey bands (N = 1600): the compressed starts' compartments in the two Z groups, split at the cut P6 printed (the largest gap); the label gives each group's mean (SE). Box: L0 = N pi / (8 H eta) exactly (--gen3-exact-box), radius 0.5.
Builds, all gen3: P6 engine-gen3 f72d764; P4, P4a and P5 engine-gen3 06e7248. Values as printed in P6's tables (experiments_gen3_p6_261010/p6_tables_output.txt, engine-gen3 c540b746; sec. 4.7.43).
tables: hspist3/experiments_m1_figures_261011/tables/ (main b7ac7c59); script: hspist3/experiments_m1_figures_261011/m1_figures_261011.py (main 973c1839)