SEEDS OF GRACE PRI. SCHOOL

P.O BOX NTENJERU- MUKONO











PRIMARY FIVE SCIENCE BOOKLET





HOME STUDY MATERIAL FOR 2024

NAME:	
SCHOOL:	
CLASS:	
CLASS TEACHER	

Designed By: GOOD PRICE STATIONERY

For inquiries, contact us on 0758 299 602,0783415073, 0753 046 640 **Location**: we are located in Pallisaalong Gognyo road.

Other services offered: printing, photocopying, scanning, computer training & repair, Airtel Money, MTN Mobile money plus all stationery products, Exams & Lesson notes.

'Quality is Our Pride"

TERM I TOPICS

- 1. Keeping birds and bees
- 2. Measures
- 3. Immunization
- 4. Digestive system

TOPICAL BREAK DOWN

- What is poultry
- What is poultry keeping
- Examples of poultry birds
- Importance of poultry keeping
- External structure of poultry birds (Diagram)
- External features and their uses to a chicken
- Comparing external features of different kinds of poultry birds.
- Breeds of chicken
- Types of poultry breeds (local, exotic and cross-breeds)
- Characteristics of each type of chicken
- Systems of keeping poultry (free range, deep litter, Battery, Fold pen)
- Simple diagrams to illustrate each type of system
- Caring and management of poultry e.g. How to improve on local breeds.
- Feeding in poultry (poultry feeds)
- The digestive system of a bird (parts and their functions)
- Reproduction in chicken birds (parts of an egg and their functions)
- Factors that may lead a fertilized eggs from not hatching
- What is incubation? Incubation periods of different birds
- Types of incubation (Natural and Artificial incubation)
- Diagram of an (artificial incubator)
- Comparison between natural and artificial incubation
- What is brooding?
- What is a brooder?
- Simple diagram of a brooder
- Different ways of improving and managing poultry
- Poultry vices
- Diseases and parasites of poultry

KEEPING POULTRY

Poultry: Poultry refers to domestic birds

Poultry keeping: This is the rearing or keeping and management of domestic birds.

Terms used in poultry and their meanings

- **Hen** Adult female chicken.
- **Cock** Adult male chicken.
- **cockerel** Young male from 8 weeks on wards
- **capon** castrated male
- **pulle**t young female
- **chick** very young bird from hatching to 8 weeks
- **Incubation** process by which an egg is given necessary condition to hatch into a chick
- **Incubator** A machine used to hatch eggs into chicks
- Layer Type of bird which kept mainly for egg production
- **Broiler** This is a type of chicken kept mainly for meat production.
- **Brooding** It is the giving of special care to young chicks below 8 weeks.
- **Broody hen** A hen sitting or incubating eggs to hatch them.
- **Brooder** A special structure for keeping chicks below 8 weeks
- **Culling** This is the removing of un wanted chicken e.g. sick or un productive birds from the flock.
- Moulting The process by which birds shed their feathers to replace them

(a) **Importance of poultry keeping**

- Kept for production of Eggs and chicken.
- Some birds are kept as pets.
- Feathers can be used for decoration
- Their droppings are used to make farm yard manure

Types of poultry


- chicken
- ducks
- Turkeys

- Guinea fowl
- Geese

(f) External features of a domestic bird

- The external features of a domestic bird consist of the following major parts namely; comb, beak, wattle, breast, belly, wing, shank, toe, claws,foot, spur, tail feathers, back, nape, earlobe and eye.

Parts of a domestic bird



(g) Functions of each part

- **Beak** picks up food
- **Nostrils** Used for breathing and sense organs for smell
- **Eyes** Are sense organs for sight
- Wings Enable birds to fly
- **Feathers** Protect the body from cold weather
 - Enable birds to fly (quill feathers)
 - Acts as a source of attraction to opposite sex during mating period
- Legs and toes Help in movement
- Toes are also used to scratch for food
 - **Spur** Used for protection

Firm grip during mating

Feathers

Feathers are the outermost covers of the bird's body.

Uses of feathers to birds

- They cover the bird's body and keep it warm
- The feathers enable the birds to fly (Quill feather)
- Give the birds shape and colour for identification

Uses of feathers to people


- For decoration
- For making pillows

Types of feathers

- Quill feathers/flight feathers
- Body feathers/covert feathers
- Down feathers/ contour feathers
- Filoplume feathers/hair feathers

Structure of Quill Feather

- It consists of the following major parts, Quill, after shaft rachis or shaft and vanes
- Used by birds for flying.



Meaning of each part

- shaft The long hard part in the middle of the feather
- vane Flat expanded part of the feather
- Quill The extreme end of the shaft
- Barbs They form the vane

Body feathersor covert feathers

• These cover most of the bird's body and keep the bird warm

Structure of body feather



Thedown feather

- These are the first feathers to appear on a chick
- They help to prevent heat loss from the bird's body

Structure of a down feather.



Filoplume (Hair feathers)

- These are feathers found nearest to the skin of a bird
- They can be clearly seen after removing the body and down feathers.

Structure of a filoplume feather



Differences between a hen and a cock

<u>Cock</u> <u>Hen</u>

- Has a big comb and wattle - Has a small comb and wattle

- It has a long spur - Has a short spur

-Has shiny nape -Has dull nape

-Has long tail feathers -Has short tail feathers

Types of chicken

• A type of chicken is a class of chicken kept for specific purpose.

There are types of chicken.

- Broilers
- Layers
- Dual purpose chicken

(a) **Broilers (table birds)**

These are a type of chicken kept mainly for meat production

Examples of broilers

- Ply mouth rock
- Light Sussex

(b) Layers (light breeds)

These are a type of chicken kept mainly for egg production

Examples of layers

• White leghorn

- Ancona
- Minorca
- Brown egger
- Sykes

(c) **Dual purpose breeds**

These are a type of chicken kept mainly for both meat and egg production

Example of dual purpose chicken

- Rhode Island Red
- New Hampshire
- Black ausralorp

Breeds of chicken

A breed of chicken is a family of chicken with similar characteristics

Breeds of chicken include:

White leghorn

Ancona

Light Sussex

Minorca

Sykes

Types of breeds of poultry

There are three main or major types of poultry breeds namely: local breeds, exotic breeds, cross breeds

Local breeds

They are sometimes referred to as indigenous or native breeds

They are called native or local because they existed in Uganda for a very long time.

N.B These types of breeds are named after their area of origin e.g. enkonko enganda

Characteristics local breeds

- They are more resistant to diseases and parasites
- They take a long period of time to grow
- They lay small and only a few eggs during each laying season
- They can feed on a variety of feeds.
- They are resistant to bad weather conditions.
- They produce less chicken
- Have different colours

Advantages of local breeds over exotic breeds

• Local breeds are more resistant to disease than exotic breeds.

How to improve upon the local breeds of poultry

• By carrying out cross breeding.

(b) Exotic breeds

• These are breeds which were imported into Uganda from other countries e.g white leghorn, Rhode Island red and light Sussex etc

Characteristics of exotic breeds

- They are less resistant to parasites and diseases
- They grow fast and mature in a short time at the same rate
- They laymanyeggs
- They have the same colour, shape and size
- They have the same ancestors
- They produce good quality meat

Advantages of exotic breeds over local breeds

- They grow and mature faster than local breeds.
- They lay more eggs than local breeds
- They produce much meat than the local breeds

Cross breeds

These are types of breeds got by mating exotic breeds with local breeds.

Advantages of cross breeds over local breeds

They produce much meat than local breeds

They grow and mature faster than local breeds

Advantages of cross breeds over exotic breeds

They are fairly resistant to diseases than exotic breeds

Poultry keeping systems (systems of keeping birds)

The systems of poultry keeping depend entirely on the amount of land available

Common systems carried out in Uganda are: free range (extensive) Deep litter systems (intensive), Battery system, Fold system.

(a) Free range system (Extensive)

This is a system where by birds are left free to move (roam) around looking for their own food and in the evening they come back for shelter.

This system of poultry keeping is very common in most parts of Uganda especially in rural areas where land is available.

Simple diagram to illustrate a free range system



Advantages of free range system

- Birds get variety of feeds (balanced diet)
- Birds get enough exercise
- Manure is well spread in the field
- It is a little bit cheap and easy to manage (farmers doesn't need to buy food)
- Birds need little care
- Vices (bad habits) such as cannibalism, egg eating etc are greatly reduced

Disadvantages of free-range system

- Record keeping becomes difficult
- It requires a very big piece of land
- Poultry may eat up farmers' crops
- It is very difficult for a farmer to carry out close supervision
- Eggs may be laid in the bush and end up being stolen or eaten by wild animals
- Birds can be easily pick worms and other parasites
- Birds can easily be stolen or eaten by predators

Deep litter system

Here, the birds are kept in the house whose floor is covered with soft material called **litter**The house where such a practice is carried out is called **a deep litter house**N.B Deep litter system is carried by commercial farmers

Materials used as litter

- Coffee husks
- Crushed maize cobs
- Wood shavings

Qualities of good litter

- It should not be too dusty
- It should not be too sharp

- It should not become wet easily.
- It should be able to absorb moisture from poultry wastes
- It should not be too comfortable for birds

Importance of litter

• To keep the floor of a deep litter house warm and dry

Simple diagram to illustrate deep litter system.



Advantages of deep litter system

- Collection of eggs is easy
- Large numbers of birds are kept in a small area
- It is easy to feed birds in one place
- Birds are protected from predators
- Record keeping is easy
- Birds cannot easily pick parasites
- Chances of disease spread are reduced
- It is easy to collect manure

Disadvantages of deep litter system

- It is expensive to start and maintain the system
- Much attention and care is needed
- If litter stays for long it produces bad smell
- It is tiresome to look after the birds
- Vices such as egg eating, cannibalism, etc may be developed by birds
- Birds may not get enough exercise
- Comfortable litter may lead the birds to become broody.

(c) **Battery (cage) system**

In this system, birds are kept in separate cages (one or two)

The floor slopes slightly so that when eggs are laid they roll into a wire trough Droppings fall under raised cage and make the floor remain clean

Advantages

Food, water and eggs are not easily contaminated

- Sick birds can be detected and removed.
- Many birds can be kept on a small piece of land
- Eggs laid remain clean
- It is very easy to identify poor layers.
- Birds are protected from predators
- Poultry vices are reduced
- It is easy to collect manure
- Record keeping of individual birds can be kept easily
- Diseases and parasites do not spread easily

Disadvantages


- It is very expensive (cages are expensive to construct)
- Birds may not get a balanced diet
- Much labour is needed
- Birds do not get enough exercise

Fold or pen system

In this system, birds are kept in a small structure called a fold or pen or ark The fold or pen and the birds are moved to new places daily.

The fold or pen provides shade and egg laying space for birds.

A simple structure of fold or pen



Advantages of fold or pen system


- Birds are restricted and cannot destroy crops
- Chicken feeds are not wasted
- Manure is well spread on the farm
- Chicken are safe from predators
- Feeding is cheaper and birds get a balanced diet

Disadvantages of fold or pen system

- It needs a very big piece of land.
- Folds get old quickly due to frequent movement
- More labour is needed to move folds every day
- It is not applicable on hilly and swampy areas
- It is very expensive to construct fold
- Only few birds are kept

Birds do not get enough exercise

Simple diagram of alimentary canal



The digestive systems of a bird (Alimentary canal)

• The digestive system of a bird (alimentary canal) consists of the following major parts namely; Beak, Gullet, crop, stomach, small intestine, large intestine, caeca, Gizzard and cloaca (vent), Beale


Functions of each part

- (a) **Beak** It picks food
- (b) **Gullet** It is a food passage to the crop
- (c) Crop It stores, moisture and softens the food before it goes to the stomach
- (d)Stomach- it is where food is mixed with digestive juices.
- (e) **Gizzard** In this organ, food is crushed into small particles to help small stones called **grit or pebbles**
- (f) **Small intestines** Digestion of food is competed here and digested food is absorbed into blood stream
- (g) **Large intestines** absorption of water from undigested food takes place here.
- (h) caeca- digestion of roughage takes place here by the help of useful bacteria
- (i) Cloaca (vent) –it is a passage of eggs and droppings at different times.

Reproduction in poultry

Poultry reproduce by laying eggs

Reproductive system of a bird



Functions of the parts

Ovary: to produce eggs or ova

Funnel'[infundibulum]: it is where egg yolk is produced and added to an egg.

Fertilization also takes place here

Magnum (Albumen producing region)

It is where albumen is added to an egg

Isthmus:it is where shell membrane, water, air space, chalaza and mineral salts are added to an egg

Uterus (egg shell producing region)


It is where egg shell is added to an egg

Vagina: It is where pigments are added to an egg

An egg (fully grown/developed)

• The structure of an egg consists of the following parts; Egg shell ,shell membrane, air space, albumen (egg white), chalaza, embryo or germinal disc

Cross section through an egg



Functions of each part

Egg shell – It is made up of calcium

- It protects the inner parts of an egg
- It's porous to allow gaseous exchange to take place

Shell membrane – To hold the inner contents of an egg.

Albumen (Egg white)

- It is also called egg white
- It provides proteins and water to the growing Embryo

Chalaza

- It holds the yolk and embryo in position
- Carries water and food to the embryo
- Carries oxygen from the air space to the embryo

Air space

- It provides fresh air (Oxygen) to the embryo
- Takes carbon dioxide produced by the embryo

Yolk – It is also a source of proteins and fats for the growing embryo

Abnormalities in eggs

- Yolk less eggs
- Double yolk eggs
- Thin shelled eggs
- Eggs with blood spots
- Eggs with meat spots
- Eggs with soft shells

Factors that may lead fertilized eggs fail to hatch

- Eggs laid with two yolks
- Eggs with a lot of moisture content
- Eggs exposed to too much heat
- Failure of an egg to receive enough oxygen
- Failure of an egg to receive enough warmth

<u>Incubation</u>- This is the process whereby a fertilized egg is given necessary conditions to hatch into a young bird

<u>Incubation period</u> – This is the time taken for a fertilized egg to hatch into a young bird **Incubation of different birds**

- Ducks between 30-31 days
- Turkeys 28 days or 4 weeks
- Pigeon 16 days (2 weeks, 2 days)
- Geese 30 days

Types of incubation

(a) Natural incubation

This is when a mother hen sits on her eggs to enable them hatch.


• The broody hen usually comes off to feed and gets little exercise

Diagram of a hen incubating eggs

A chick hatching from an egg







How to ensure effective natural incubation

- Use good fertilized eggs
- The place for incubation should be with dim light
- Protect the hen from rats, snakes etc which might eat eggs
- Provide a good nest which is clean, dry and comfortable for the broody hen

Advantages and Disadvantages of natural incubation

Advantages

- It is cheap and easy to manage
- Chicks get extra care from the mother hen
- Chicks get protection from the mother hen

Disadvantages of natural incubation

- Fewer chicks are hatch at a time
- Natural incubation can't be used by commercial farmers
- Eggs are not tested for their fertility
- Some eggs are not hatched
- Hens used may not be good for incubation
- The birds can be attacked by snakes, wild cats and mites

Artificial incubation


This is where by a machine called incubator is used to hatch eggs

An incubator may provide all conditions that a broody hen provides

An incubator can use paraffin or electricity to provide enough warmth

A locally designed incubator uses paraffin and modern one uses electricity

An incubator



Advantages of Artificial incubation

- Very many eggs can be hatched at a time
- It is good for commercial farmers
- Eggs are tested for their fertility in the process called candling
- Temperature is maintained properly in an incubator by the use of a thermometer
- All eggs are hatched since they are first tested for their fertility

Disadvantages of Artificial incubation

- It is very expensive method of incubating eggs
- It needs regular supervision
- It needs skilled man power
- It is tiresome
- Eggs can be affected in case of frequent black out or load shading

Brooding

Brooding is a process of giving special care to chicks below 8 weeks

Types of brooding

- Natural brooding
- Artificial brooding


Natural brooding

This is a method where by a hen takes more care of her chicks by providing warmth, security and food to them

Farmers may also support by providing more food and water to the mother hen and its chicks

When the chicks are big enough, they are left on their own

A hen looking after her chicks



Advantages of natural brooding

- The mother hen looks for food for its chicks
- The mother hen provides security for its chicks
- Toes and feather pecking are reduced among chicks

Disadvantages of brooding

- Chicks may die due to poor protection
- Chicks are exposed to predators
- It is not carried out on large scale
- It gives little profit
- A mother hen may be stolen or eaten by predators leaving chicks without care

Artificial brooding

This is where by chicks are kept in a special structure called a brooder

A brooder is a special house /structure designed to keep chicks below 8 weeks


Types of brooder

- (a) Infra red brooder
- (b) Kerosene brooder

<u>Infra – red brooder</u>

Infra – **red brooder:** In this type, electricity is used as a major source of heat Lamps are hung from the ceiling and adjusted (changed) from time to time as chicks grow

A simple diagram of infra-red brooder



Kerosene brooder

In this type, farmers use paraffin lamps (lanterns) as sources of heat.

This type needs careful handling of lamps not to burn chicks and avoid accumulation of soot

The lamp should also be with guards to keep off chicks from being burnt

A layout of kerosene brooder



Management practices

• Management practices are routine activities carried out on farm to increase on the quality of farm animals and their products

Common daily/routine activities include

- Record keeping
- Regular cleaning
- Vaccination
- De-worming

- Culling
- Dusting
- Egg collection
- Debeaking

- Cross breeding
- Housing
- feeding

Feeding poultry

Reasons for good feeding

- We feed chicken so that they grow well
- We feed them in order to get high quality eggs and meat
- Proper feeding boosts the birds' immunity

Classes of food given to birds

• Birds need mash containing all classes of food such as; proteins, vitamins, carbohydrates, minerals, water, green etc.

Proteins: (body building food)

- Enables birds to grow well and put on flesh (meat or chicken)
- Crushed fish (mukene), Soya beans etc are rich source of proteins

<u>Vitamins</u>: keep the body healthy or protect the body from diseases

- They are healthy/protective food
- Green vegetables are rich sources of vitamin

Carbohydrates: (energy giving food)

- Provide the birds with energy
- Maize bran, cassava, rice, millet, sweet potatoes etc are rich source of carbohydrates

Minerals: Calcium makes the bones and eggs shell strong

- Calcium, iron, crushed shells, iodine, dried animal blood m-(iron) etc are examples of minerals
- Oyster shells, Green vegetables, common salts are also rich source of minerals

Water

- Water promotes easy digestion of food
- Maintains body temperature of a bird

Chicken mash

- Chicken are fed on different types of food called **mash.** (mixture of silver fish (Mukene), maize bran (outer layer of the grain) Oyster etc depending on what age the types of birds are;
- Chicken mash is grouped into four (4) types namely; chicken and duck mash, Growers mash, Layers mash and broilers mash

Chickand duck mash

- It is given to chicks and ducklings which are 1 day 8 weeks
- This mash contains high protein content to support fast growth in chicks
- Each chick needs about two grammes (2gm) of mash per day

(b) **Broilers mash**

- It is prepared for chicken that provide meat (chicken)
- It is introduced to broilers from 8 weeks on wards

(c) Growers mash

• It is given to growing birds e.g. layers from 8 weeks to about 16 weeks (8-16) weeks and Broilers between 4-8 weeks

(d) Layers mash

- It is prepared for laying chicken
- It is introduced to layers at 16 weeks

Summary table

Mash	Bird's age
Chick and duck mash	Chicks and ducklings of 1 day to 8 weeks (two
Layers mash	months)
Broilers mash	Layers from 8weeks – Broilers from 4-8 weeks
	Broilers to 8 weeks on wards

Feeding tools

- The mash given to chicken must be always put I a clean containers called feeding troughs
- A good feeding trough should be constructed with a spinning/rotating handle in the middle to prevent birds from stepping on and spoil or contaminate the mash

Simple diagrams of feeding tools


Feeding trough

Drinking trough

Spinning/rotating handle trough

chicken mash

conical drinking water



Wooden feeding trough

General practices in feeding poultry

- Give correct feeds to the right type of a bird according to age and products
- Introduce new feeds gradually (slowly)
- Measure/record and give correct amount of feeds per day
- Contaminated food water should not be given to birds
- Provide fresh greens in the daily diet of birds
- Cull the birds

Housing poultry

Reasons for proper housing

- To prevent loss of eggs
- To prevent the birds from being stolen
- To protect the birds from predators

Qualities of a good poultry house

- It should be well ventilated to allow free exchange of gases
- It should have enough light to enable the birds to see feeds and water
- It should have enough space for the birds
- It should have strong doors
- It should have a roof which does not leak
- It should be free from dampness or moisture
- It should be free from diseases and parasites etc

Selection of birds for breeding

Means choosing the best bird for breeding or reproduction

Select the best quality birds which:

- Grow fast
- Lay many eggs
- Produce good meat

Are resistant to diseases

Cross breeding

This is the process of mating birds that are not related to each other e.g. pure local + pure exotic breed = cross breed

Advantages of cross breeding

- They grow fast and are large like the exotic breeds
- They lay a lot of eggs
- They produce soft meat like exotic breeds
- They are resistant to diseases like local breeds
- They are resistant to droughts etc

Poultry vices

These are bad habits developed in poultry

Common poultry vices are;

- Egg eating
- Feather pecking

- Toe pecking
- Cannibalism

Causes of poultry vices

- Over-crowding of birds (too many birds in a small area)
- Boredom in the house (lack of what to do)
- Failure for a farmer to provide enough and balanced food
- Failure to remove immediately damaged eggs from poultry house
- Failure to collect eggs regularly from the poultry house

Prevention/control of poultry vices

- Carryout de-beaking
- Collect eggs regularly from laying boxes
- Provide the birds with enough food.
- Provide food to the birds in time
- Remove broken eggs from the poultry house as soon as possible

Diseases and parasites of poultry

General causes of diseases and parasites

- Poor feeding of poultry on already contaminated food
- Poor housing structure for birds
- Dirty houses lead to an outbreak of diseases and parasites

Dangers of diseases and parasites

- Death may occur
- Lower the quality of products
- Reduces the bird's rate of growth

Common diseases of poultry

<u>Disease</u> <u>cause</u>

• Coccidiosis Protozoa

• Newcastle virus

• Fowl pox virus

• Fowl typhoid bacteria

• Avian leucosis

• Pneumonia Bacteria

• Marek's disease(fowl paralysis) Virus

Pneumonia

- It is caused by bacteria
- It attacks the lungs

Signs and symptoms

- Difficulty in breathing
- Coughing

- Lack of appetite
- Mucus from the nose

Prevention and treatment

- Keep the poultry house clean
- Make the house well ventilated

Separate diseased birds from healthy ones

Treat early with antibiotics

<u>Parasites</u> - These are living organisms which depend on other living organisms for food and shelter.

Types of parasites

- Ecto parasites (External parasites)
- Endo Parasites (internal parasites)

Ectoparasites

These are parasites which live on the body of a host

Examples of ecto parasites of poultry

Fleas, mites, lice

Endo - parasites

These are parasites that live inside the body of a host.

• Common examples are: Round worm, Tape worm, Threadworm etc

Types of	Examples	Effects
parasites		
Ecto parasite	Fleas	- They create wounds on skin of birds
		- They cause itching and scratching of
		body that may lead to loss of feathers
	mite	- They suck blood from the birds making
		them anaemic

		-	Egg laying is greatly reduced
	fowl lice	-	Mites cause a disease called mange
Endo parasites	Tape worm	-	They cause stunted or poor growth
		-	Egg laying is reduced
	Round worm		

Control/Treatment of parasites in poultry

- Use of pesticides for external parasites
- Carry out regular deworming to kill internal parasites.
- Clean poultry houses regularly.
- Apply Vaseline jelly on affected parts of the body caused by mites and fleas
- Give birds clean water and food
- Give birds water and food in clean troughs

Keeping farm records

Farm records are written information showing activities carried out on a farm.

Types of records

- Health records
 - Feeding records
 - Operation records
 - Production records
 - Sales and expenses records (expenditure) etc

Importance of keeping records on a farm

- It helps the farmer to plan for his farm.
- It helps the farmer to know if the farm is making profits or losses
- Help the farmer to know the birds to cull
- Farmers are able to know the sales and expenditure
- Help a farmer to acquire loans from a bank
- Help a farmer to be taxed fairly by government.

A format of production record

Namirembe Parents' School farm

P.O. Box 29928

Kampala

(a) month: March 2011

(b) Breeds of chicken: white leghorn

(c) Age of the flock: 18 weeks

(d) Number of birds 500

Number of	Date	Morning	Afternoon	Total	Remarks
eggs broken					
50	24.03.11	250	200	450	Poor
					handling
-	25.03.11	300	200	500	Careful
					handling

SOCIAL INSECTS

Social insects are insects that live and work together

Examples of social insects include:

Honey bees

Wasps

Termites

Bee keeping [apiculture]

Bee keeping: is the rearing of bees.

Apiculture: is the rearing of honey bees on a large scale.

Importance of bees to plants and people

People

- People get honey from bees
- People get wax from bees.
- Bee farming creates employment to people


Plants

• Bees help in pollination

The life cycle of a honey bee

- The honey bee undergoes complete metamorphosis
- The Queen lays eggs
- The larvae are developed from the eggs and are fed by the worker bees
- They change in pupae
- The pupae develop into adult bees

Simple diagram to illustrate




Types of honey bees

- There are three types of bees in each colony namely;
- (a) Worker bees (b) Drone bees and (c) Queen bees

Queen bee

- It is head of the colony
- The Queen bee is formed when female grub is fed on royal jelly throughout its life cycle.
- its main function is to lay eggs
- It is a female fertile bee
- It is the biggest in the colony
- It's the master bee in the hive
- It can last 4 5 years
- Its abdomen is longer than others
- Its fed on special food by workers called royal jelly

Simple diagram to illustrate



Worker bees

- These are female sterile bees
- They are the smallest in the hive
- They have hair on the back and on their legs
- They have a sting
- They are many in the hive

Roles of a worker (s) in the hive

- It feeds the Queen bee on a special food called royal jelly
- It feeds the grub
- Protects the Queen and the hive
- They make honey
- they repair the hive
- They clean the hive

• They regulate temperature

Simple diagram of a worker bee



Drone bee

- It is a male bee in the hive
- It is the second biggest in the hive
- Its main work is to mate with the queen bee.
- It makes a buzzing sound
- It's the most hairy bee in the hive

Simple diagram of a drone bee



A swarm – is a large cluster or group of bees

Swarming

• Is the massive movement of bees from one place to another looking for a better hive.

Reasons for swarming in bees

- When another Queen is produced
- For the queen bee to mate.
- When they are overcrowded in the hive
- When the hive is leaking or damaged
- When there is a bad smell around the hive
- Death of the Queen
- Over heating of the hive by direct sun shine
- When they are disturbed by enemies
- When there is shortage of water
- Noisy places can lead to bee swarming

- Area with a lot of smoke
- Dampness of the bee hive

Enemies of bees

(a) Ants – several types of ants attack bees or their hues e.g. the wood ants and safari ants

How to protect bees from ants

- Hang the hive from between poles
- Put or/smear the bases of the poles with grease.
- Place grease on the wires, This can prevent the ants from reaching the hive

(b) Honey Badgers

• These spoil the hives, kill bees and eat the honey

Other pests include

• Hive beetles

Wasps

wax moth

• Pirate ants

Hawk moths

Modern methods of keeping bees

Bees can be kept using two types of bee hives namely

- Traditional bee hive
- Modern bee hive

A hive – This is a structure where beeslive.

(a) Traditional bee hive

This is the type of bee hive made up of local materials such as hollow logs, grass, woven sticks, cow dung, clay etc

When the bee hive is made, it is hung up in places like trees

Examples of local hive

- Kigezi bee-hive
- Dug out log hive

Advantages of traditional (local) bee hive

- It is cheap
- It is easy to make/construct because materials are locally available

Disadvantages of local bee hive

- The honey collected is always dirty
- Colony cannot be inspected
- The hive is damaged when harvesting honey
- The brood is damaged when harvesting honey
- Too much honey can break the hive


(b) Modern bee hive

• These are constructed out of materials and piece of wood


Common examples are: Top bar hive, Box hive, and Tin hive

- They are constructed with chambers called brood and honey chambers
- Two chambers are separated by a metal called **Queen excluder**
- The queen excluder has holes of given sizes through which only workers pass through to look after the Queen
- Once the Queen is put into the brood chamber, it cannot move to the honey chamber because the Queen is too big to pass through the small holes
 - The queen excluder prevents the Queen bee from entering the honey chamber to eatthe honey

Internal structure of a modern bee-hive



Top bar hive Box hive



Advantages of modern bee hive

- The honey collected is always clean
- The eggs, larvae and pupae are not disturbed
- Number of bees continue to multiply
- Honey combs which are ready for harvesting can be removed and those which are not ready can be replaced back
- Honey and wax can be inspected separately
- Modern bee hives are long lasting

Disadvantages of modern bee-hive

• It is very expensive

Harvesting honey

Honey must be or should be harvested in the evening when it is cool and all the bees have settled inside the hive

When harvesting honey, the farmer must have the following equipment

- A bee veil
- A bee keeper's glove
- A bucket
- A suitable protective clothing (Overall)

- A smoker
- Gum boots
- Knife

Uses of each equipment above

Bucket: it is used for collecting honey.

Bee veil: protects the face of a honey harvester from being stung by worker bees

Bee keeper's gloves: protect hands of a honey harvester from being stung by worker bees

Smoker: for smoking the hive to calm the bees

Knife: for cutting honey combs

Gum boots: To prevent worker bees from stinging the feet of a honey harvester

Steps followed when harvesting honey

- Smoking the hive
- Putting the hive down
- Opening the hive
- Cutting the honey combs

A simple diagram of a farmer ready to harvest



Factors to consider when harvesting honey

- Don't frighten the bees while harvesting
- Don't kill bees while harvesting
- Dress properly in suitable clothes

- Use a smoker
- Bees should not be crushed while harvesting honey

Honey extraction

Honey extraction is a process of separating honey from honey combs.

Method of extracting honey

- By floating the wax
- Centrifuging method
- Pressing the wax method

Products got from bees

- Honey
- Bee wax

Uses of honey to people

- honey is eaten as food
- It is used to sweeten tea.
- It is used as medicine for cough.

NB: the food value got from honey is carbohydrates

Products from bee wax

- Candle wax
- Polish for wood, floor and shoes
- Chewing gum

Care and management of bees

Siting the bee hive. This is simply the selecting a suitable place in which to put the hive A place in which a number of hives are kept is called **an apiary**

Factors to consider when siting hives

- Bees should be sited away from houses, roads and animals
- Bee hives should be put in places near water sources
- Places with flowering plants
- Sheltered places
- Places with short grass

(b) **Stocking the hive**

This is a process of encouraging bees to occupy an empty hive

It is done in various ways:

- (a) Use of a catcher box
- (b) Use of a swarm catching net
- (c) Setting up the hive in its permanent place, smear the inside parts with wax and wait bees to come for their own.

NB: the handle of a swarm catching net can be made longer to catch a higher swarm

A swarm catching net



Feeding bees

• Bees naturally feed on nectar from flowers

Conditions that make bees be fed on supplementary food

- (a) When they are new in a hive (new colonies)
- (b) During the drought conditions
- (c) To encourage multiplication

Examples of supplementary feeds

- (a) Syrup (ordinary sugar mixed water)
- (b) Ordinary water put in a container at a distance of 10m from the ordinary hive

Handling of bees by a bee keeper (Three occasions)

- When harvesting honey
- When inspecting the hive either to know whether he can harvest honey or there are problems
- Catching bees for stocking

MEASURES

Regular shaped objects

These are objects which have proper or regular shape. Examples include: - a cube, a cuboids, a brick, blocks, tins and other cylinders.

Measuring regular shaped objects.

(a) Length

Length is measured in the following units

- millimetres (mm)
- centimetres (cm)
- decimetres (dm)
- metres
- decametres (Dm)
- Hectometres (Hm)
- Kilometres (Km)
- The most commonly used units for measuring length are centimtres, metres and kilometres
- The smallest unit for measuring length is millimetre.


- The biggest unit for measuring length is kilometre
- The basic nit for measuring length is metre

Area:

- Area is the total space occupied by a figure (two dimensional figure)

Are is measured in square units

In two dimensional figure, the longer side is the length while the shorter side is the width.




Volume

- Volume is the space occupied by an object.
- Volume is measured in cubic units i.e. cubic metres, cubic centimetres or cubic millimetres.
- Volume s also measured in litres, milliliters and centiliters (cl)

Finding volume of regular shaped objects


The volume of regular objects like, cuboids e.g. boxes blocks bricks etc, can be found after knowing their length, width and height.




Volume = Length x width c Height

Exercise

1. Find the volume of the figures.





- 2. Find the volume of a brick of length 6cm width 4cm and height 3cm.
- 3. Find the area of a cube whose side is 3cm.

Irregular objects

Irregular objects are objects without regular shape.

Examples include stones, keys, needles, soda bottle etc

Finding volume of irregular objects

- The method of finding the volume of irregular objects is **measuring by displacement**.
- There are two instruments used in measuring by displacement i.e.
 - (a) the measuring cylinder
 - (b) the over flow can or eureka ca

Using a measuring cylinder


Step I

Pour water into a measuring cylinder so that it is a half full.

Record the volume of water say 30cc

Step II

- Get a stone and tie some thread around it
- Lower the stone into the measuring cylinder so that the stone is covered by water.
- Record the volume of water again say 35cc.




volume of the stone =
$$2^{nd}$$
 level - 1^{st} level
= $35cc - 30cc$
= $5cc$

The volume of water is 5cc because the amount of water displaced is always equal to the volume of the irregular object.

Using an overflow can (eureka can)

- Fill the overflow can with water so that the water pours out through the spout until it is at the same level of the spout.
- Put a stone whose volume you want to find and lower it into the can.
- The water will overflow and pour into the measuring cylinder.

Find the volume of the stone by reading the level of water in the cylinder.



The volume of the irregular object is 10cc.

Difference between capacity and volume

Capacity is measured in litres while volume is measured in cubic units Capacity refers to amount of a liquid a container can hold while volume is the space occupied by a substance.

Weight

- Weight is the gravitational pull acting on an object.
- Weight of an object depends on three things i.e. its size, the substance form which it is made and the gravitational pull on it.
- Weight changes according to the distance between the centre of the earth and the body of object.

Note: Objects weigh less on the moon that on earth because the moon is for away from the centre of the force of gravity from the earth which pulls it.

- When you lift or pull something you are overcoming the force of gravity.
- It is difficult to push a wheel barrow uphillbecause you are opposing the force of gravity.
- It is easier to push a wheel barrow downhill because **you are helped by the force of gravity**.
- Weight is measured in units called Newtons (N) but (10N = 1kg) or (1N = 100g) Weight is measured using an instrument called **spring balance**

Diagram of a spring balance



Mass

Mass is the quantity of matter an object contains.

Mass is constant because it doesn't depend on the force of gravity

A body can have the same mass on land and on the moon


- Mass is measured in grammes, kilogrammes, milligrams etc
- The basic unit for measuring mass is grammes (grams)

Examples of machines used for measuring mass.

(a) Beam balance



(b) set of scales



(c) scale balance



Differences between mass and weight

Mass Weight

Measured in kilograms

Measured using beam balance

It is constant

Measured in Newtons Measured using a spring balance

Keeps changing from place to place

Density

- Density is the mass of a substance per unit volume
- Density (D) = mass (M)
 - = Volume

$$D=\underline{\boldsymbol{M}}$$

V

Density is measured in grams per cubic centimetre (g/cm3)

Table showing densities of some substances.

Substance	Density
Gold	19.3
Mercury	13.6
Lead	11.3
Silver	10.5
Copper	8.9
Aluminium	2.7
balss	2.6
Brass	8.3
Rubber	1.3
Salt solution	1.2
Water at $4c^0$	1.0
Ice	0.92
Methylated spirit	0.83
Kerosene	0.8
Alcohol	0.8
Pertrol	0.7
Cork	0.24
Iren	7.8
Air	0.0012
Hydrogen	0.00009

- From the above table, aluminium has the least density among metals and that is why it is used. for bodies of aero planes.
- Objects which are less dense than water will always float on water e.g. cork, ice, wood etc
- Substances which are denser than water always sink in water e.g. iron, brass, nails, stones, sand etc

Kerosene, petrol and most oils always float on water because they are less dense than water.

A hydrometer is an instrument used for measuring the density of different liquids e.g, beer, wine, acids in batteries etc

A special instrument used for measuring density of milk is called **lactometer**

Finding Density

Example I

Find the density of an object of mass 150g and volume 3cc.

Solution

$$D = \underline{M}$$
 mass = 150g

$$V$$
 volume = 30cc


$$D = 5g/cc$$

$$D = \underline{150}$$

30

Example 2

If the mass of the cuboid is 48g. Find its density



Now,
$$D = \underline{M}$$
 mass = $480g$
V volume = $48cm^3$

Solution

Here, we are not given the volume so we first find the volume

V = LxWxH

=6x2x4

=48cm

Example 3

Calculate the volume of a stone of mass 48g and density 6g/cc

Solution

We are asked to find volume

M=48g

D=6g/cc

 $V=\underline{M}$


D

<u>48g</u>

6g/cc

<u>=8cc</u>

Note: We can use the triangle below to help us to remember the formula used.



Floating and sinking

D

Floating is when an object is put in water and it stays on top of.

- Objects float on water because they are less dense than water.
- The density of water is 1g/cc and therefore, objects whose density is less than 1gg/cc float on water.

Examples of things that float on water

Cork, plastic, rubber, boats

Wood, sponge, a leaf, feathers etc

Sinking

- Sinking is when an object is put in water and it goes to the bottom of the water.
- Objects sink in water because they are denser than water.
- Objects whose density is more than 1g/cc sink in water.

Examples

- stones, sand, soil, metal, glass, nails etc

Note: A sinking object displaces water equal to its volume while a floating object displaces water equal to its weight

IMMUNIZATION

What is immunization? This is the introduction of vaccines to the body.

Types of immunization

- o Primary immunization this is done to all infants or children below 5 years
- Mass immunization this is done to all people when there is an outbreak of a disease in an area
- o Booster immunization this is done to maintain the required amount of antibodies in the body of a person e.g. administration of T.T vaccine to teenage girls and pregnant mothers.

Importance of immunization

- Protect the people from acquiring dangerous disease
- Reduce infant mortality rate.
- Enables the body to resist diseases

Immunity

This is the body's ability to fight or resist disease or infection.

Types of immunity

- Natural immunity: This is acquired without receiving any vaccine.
- Artificial immunity: This is when vaccines are introduced into one's body.

How the body acquires natural immunity

- Through suffering and recovering from an illness.
- From the mother to her baby through breast feeding

How the body acquires artificial immunity

• Through immunization

Vaccines

Vaccines are medical drugs used for immunization

Methods of vaccine administration

- Oral administration
- Injection method

Types of vaccines

- Toxoid vaccines
- Attenuated/living vaccines
- Dead/ killed vaccines

Examples of vaccines

BCG Vaccines

- It provides immunity against tuberculosis
- It is injected on the right upper arm at birth

- It is administered by injection method
- It is injected once that is it has only one dose

Polio vaccine

- It provides immunity against polio.
- It is administered by putting into the mouth
- Four does are given at different intervals

1st dose at birth (polio 0)

2nd dose at 6 weeks (polio 1)

3rd dose at 10 weeks (polio 2)

4th dose at 14 weeks (polio 3)

Measles vaccine

- It provides immunity against measles.
- It is injected on the left upper arm at nine (9 months)
- One dose of measles vaccine is applied.

Pentavalent vaccine (DPT-HepB+Hib)

- It provides immunity against Diphtheria, whooping cough, Tetanus, Hepatitis B, Haemophilus Influenza B
- o It is injected on the left upper thigh
- o Three doses are given at different intervals
- o 1st dose at 6 weeks
- o 2nd dose at 10 weeks
- o 3rd dose at 14 weeks

Note: **HepB** is the vaccine for hepatitis B while**Hib** is a vaccine for haemophilus influenzaB

HPV Vaccine

- o **HPV in full stands** for Human Papilloma virus
- o -It is used to immunize against cervical cancer.
- o -It is administered to girls from 9 years onwards

Pneumococcal conjugate vaccine (PCV)

- $\circ\hspace{0.1cm}$ -It is immunized against pneumonia and other pneumococcal diseases
- It is administered to babies by injecting on the right outer thigh.
- o -There are four doses of PCV administered to babies
- o First dose at 2 months
- Second dose at 4 months
- o Third dose at 6 months
- o Forth dose at 12 to 15 months

Some effects of PCV after administration

- o Swelling at the injection site
- Mild fever
- o Loss of appetite
- o Vomiting
- o Redness at the injection site

N.B If these symptoms persist visit a medical doctor immediately

Yellow fever vaccine

- o Yellow fever vaccine is used to immunize against yellow fever
- Yellow fever is caused by a virus transmitted by an infected female tiger mosquito
- o Yellow fever vaccine is an example of attenuated vaccine or living vaccine
- Yellow fever vaccine is administered through injection on the right upper arm

People who should be given yellow fever vaccine

- o People who work in research laboratories
- o People who travel to areas with yellow fever
- o Children who are 9 months and above

N.B People who travel to areas with yellow fever should travel after 10 days of immunization

- o They should also be given international certificate of immunization
- The international certificate of immunization becomes valid after 10 days of immunization and expires after 10 years.
- The people who continue to stay in areas with yellow fever after 10 years should be given yellow fever booster vaccine

People who should not receive yellow fever vaccine

- o People who are allergic to eggs
- o People with cancer
- o People with weak immunity
- Children below 9 months
- o Breast feeding mothers

N.B People who are allergic to eggs can be immunized if they are in places affected by yellow fever. But the doctor can give a vaccine in small doses to avoid serious reaction.

Childhood immunisable diseases

- Measles
- Tuberculosis
- Poliomyelitis
- whooping cough

Other immunizable diseases

o Pneumonia

- Hepatitis B
- Haemophilus influenza B
- Diphtheria
- Tetanus

- Yellow fever
- Cervical cancer
- Meningitis
- o Rabies

Tuberculosis (TB)

- o It is caused by bacteria
- o It attacks the respiratory system

Signs of tuberculosis

- o chronic cough
- o loss of weight
- o A lot of sweating at night

Symptoms

- o mild fever
- o pain in the chest or in the upper back

How it is spread

- o It spreads to a healthy person through droplets of mucus from a sick person through air.
- o It can also spread through sharing cutlery, cups and plates with an infected person
- o People can also get it through drinking un boiled milk from tubercular cows

Treatment of Tuberculosis

- o Carry out immunization
- o Treat with antibiotics
- Visit a health worker for correct treatment
- o Avoid sharing cutlery, cups plates etc with a sick person
- o treat the patient in isolation

MEASLES

- o It is caused by a virus
- o It is an air borne disease (spread through air)
- o It affects the skin and the lungs

Signs of measles

- Sores in the mouth
- o runny nose
- o skin rash
- o dry cough
- o red eyes

Symptoms of measles

- o High temperature
- o Itching rash

- o Loss of appetite
- o weakness

Prevention of measles

- Isolate measles victims
- o Immunize children at 9 months

Poliomyelitis (polio)

- o It is caused by a virus
- o It spreads when one drinks contaminated water.
- o It affects bones especially the limbs.

Signs and symptoms

- Paralysis and weakness of one or more limbs
- o lameness
- o fever

Prevention and control

- o Immunize children with polio vaccine.
- Drink safe clean water

Tetanus

- o It is caused by a bacteria found naturally in the soil
- o The bacteria enter the body through fresh cuts or wounds.
- o In new born babies, it can enter through the umbilical cord if it is cut with a dirty or unsterilized instrument like a razor blade.

Signs and symptoms

- o Stiff muscles all over the body
- o Spasms when touched.
- o The baby stops suckling the mother's breasts

Prevention and control

- o Immunization with three doses of DPT vaccine at an interval of one month starting at 6wks or 1 ½ months.
- o The vaccine is administered by injection on the left upper thigh.
- o Girls and women between 15 years and 49 years of age are given tetanus toxoid vaccine to prevent their babies from getting tetanus and also the mothers.
- All pregnant women receive tetanus toxoid in order for their unborn babies to get immunity against tetanus.

Note: DPT vaccine is given to babies at 6 weeks because babies are born with immunity against Diphtheria, pertussis and tetanus that lasts six weeks.

Whooping cough (pertussis)

o It is caused by bacteria and spread through air by coughing

Signs and symptoms

- o Coughing spells which end up in vomiting
- o Gasps for breath
- o Running nose

Prevention and control

- Immunization with DPT vaccine
- o It is administered by injection on the left upper arm
- o The vaccine is given in three doses at an interval of one month beginning at 6 weeks.

Diphtheria

o It is caused by bacteria spread through air by coughing of an infected person.

Signs and symptoms

- Sore throat
- swollen neck

Haemophilus influenza type B

This disease is caused by bacteria and prevented using Hib vaccine

Hepatitis B

It is caused by a virus and spreads through body contact with infected body fluids.

Signs and symptoms

Passing out dark urine

Nausea

Vomiting

Jaundice

Fatigue

Muscle and joint pain

Prevention

Immunize babies with Hep B Vaccine

SUMMARY TABLE

Disease	Vaccine	How it is	Doses	At what age
		administered		
Tuberculosis	BCG	Injection on	1	At birth
		the right upper		
		arm		
Polio	Polio vaccine	Drops in the	4	1 st dose at birth
		mouth		2 nd dose at 6 weeks
				3 rd dose at 10 weeks
				4 th dose at 14 weeks

Diphtheria	DPT-	Injection on	3	1st dose at 6 weeks
Whooping	HepB+	the left thigh		2 nd dose at 10 weeks
cough	Hib(Pentavalent			3 rd dose at 14 weeks
Tetanus	vaccine)			
Hepatitis – B				
Influenza – B				
Haemophilus				
Measles	Measles	Injection on	1	At 9 months
	vaccines	the left upper		
		arm		

Roles of individuals families and communities I immunization

Families

- Taking their children for immunization

Individuals

- Helping to identify children who have not been immunized in the community
- Advising members of the community to Immunize their children
- Taking their children for immunization
- collecting information about immunization in an area and using it to educate people

Communities

- setting up immunization centres
- Encouraging families to take their children for immunization
- contributing money to use on immunization
- Inviting health workers to talk to the members on immunization
- Reporting outbreak f immunisable disease in the area.

Children

Taking their siblings for immunization

Reminding their parents about immunization days

Helping to put up notices about immunization programmes

Child health card

A child health card is a document that contains the child's bio data and immunization status.

Importance of child health

- Monitor the growth of a child.
- It reminds the parent the next date for immunization.
- It helps the doctor to know which dose has been given and what is remaining.

<u>Government and non – governmental organizations which help to carry out</u> immunization activities in Uganda

- Ministry of health
- UNEPI Uganda national Expanded Programme on Immunization

Roles of UNEPI

To organize immunization programmes

To supply vaccines to immunization centres

Setting immunization schedules

Activity

- 1. What is immunization?
- 2. Define the following terms
 - a) Immunity

b) Anti – bodies

c) vaccines

- 3. Name the two types of immunity.
- 4. What is natural immunity?
- 5. Give the meaning of artificial immunity.
- 6. State two ways the body may acquire natural immunity
- 7. How does the body acquire artificial immunity?
- 8. State two methods of administering vaccines in the body
- 9. Name any four immunisable diseases.
- 10. How is polio vaccine administered?
- 11. Name one immunisable disease administered to people at the age of 9 months.
- 12. Name the vaccine which provides immunity against Diphtheria, whooping and Tetanus.
- 13. State any two importance of a child health card.
- 14. Write UNEPI in full.
- 15. Write NIDs in full
- 16. Name any two domestic animals that are vaccinated (Immunized)
- 17. State any two importance of immunizing people
- 18. Why does immunization start at birth?
- 19. State the diseases whose vaccines are given more than one disease.
- 20. Name the vaccine administered orally.
- 21. Why is the first polio vaccine given to a child at the age f 6 months and not at birth?
- 22. Why is a measles vaccine not given soon after birth but given after nine months?

THE DIGESTIVE SYSTEM

The digestive system of human being is made up of tissues and organs which take part in the digestion of food,

It is a system that deals with the breaking of food and its use in the body.

The long muscular tube runs through the body from the mouth to the anus is a part of digestive system called the <u>Alimentary canal</u> which is 10metres long. It is where digestion of food takes place.

DIGESTION

This is a process by which food is broken down into soluble particles which can be absorbed into blood stream

The process by which food is taken into the body is called **ingestion**

Digestion begins in the mouth and ends in the ileum

There are two types of digestion in the alimentary canal namely:

-Mechanical digestion/ physical -Chemical digestion digestion

Mechanical digestion: This is the breaking down of large molecules of food into simpler molecules by the help of the teeth.

Mechanical digestion is called a physical change because the physical nature of food remains un changed

Chemical digestion: This is the breaking down of large food molecules into soluble substances by the help of enzymes

Enzymes

Enzymes are biological catalysts that speed up the rate of chemical digestion

Characteristics of enzymes

- Enzymes work best under suitable temperature range
- They are normally destroyed by heat
- Each enzyme digests only one class of food.
- Enzymes work best under suitable conditions

Conditions under which enzymes work

Alkaline conditions (PH)

Acidic conditions (PH)

N.B Enzymes in the mouth, duodenum and the ileum work under alkaline conditions

2. Enzymes in the stomach work under acidic conditions

Role of enzymes

To speed up (increase) the rate of chemical digestion

The alimentary canal

This is a long muscular tube along which food moves from the mouth to the anus.

How does food move in the alimentary canal?

Food moves along the alimentary canal through peristalsis

Peristalsis is the wave like movement of food through the alimentary canal

Parts of the alimentary canal (tube)


- mouth
- Gullet (oesophagus)
- stomach
- duodenum
- ileum
- colon
- rectum
- anus

The structure of a human digestive system consists of the following parts namely:

- mouth
- gullet (oesophagus)
- liver
- stomach
- spleen
- pancreas
- gall bladder

- bile duct
- Duodenum
- small intestine (ileum)
- large intestine (colon)
- appendix
- rectum
- Anus

The structure of a human digestive system



Activity

1. What is digestion?

- 2. Where does digestion of food
 - a. Begin?

- b. End
- 3. Name the two types of digestion in the alimentary canal
- 4. What is mechanical digestion of food?
- 5. Name the muscular tube of digestive system that runs through the body from the mouth to the Anus.
- 6. Name the chemical compound that spreads the digestion of food.


Digestion of food in the mouth

- food is broken down by the teeth in the mouth
- saliva softens or lubricates food for easy swallowing
- chewed food is rolled into a bolus by the tongue
- The small ball of food is called a bolus.
- Digestion of cooked starch by enzymes called salivary amylase or ptyalin starts from the mouth
- cooked starch is then changed to maltose
 - N.B Chemical digestion of carbohydrates begins in the mouth

Food in the gullet oesophagus

- The food moves from the mouth to the stomach through a tube called Gullet or Oesophagus by peristalsis

Simple diagrams to illustrate (Peristalsis in the Oesophagus)



Food in the stomach

The stomach is a muscular bag where food is temporarily stored (It stays in the stomach for about 3 hours)

Food is churned by the stomach turning it into **chyme**

The food in the stomach which has been mixed with digestive juices is called chyme.

The stomach walls produce gastric juice

Components of gastric juice

Pepsin and rennin

Hydrochloric acid

Pepsin changes proteins to peptides

Rennin clots milk proteins

N.B Rennin is found in babies

Function of hydrochloric acid

To kill germs that escape with food to the stomach

To provide acidic PH for pepsin to digest proteins


To set un favourable conditions for amylase

The lower end of the stomach is kept closed by a circular muscle called the **pyloric sphincter** which allows a little chyme at time to pass the duodenum

NB 1. Absorption of alcohol, simple sugars, common salt and medicine takes place in the stomach.

2. Chemical digestion of proteins begins in the stomach

Simple diagram to illustrate (The stomach)



Food in the duodenum

- The duodenum is a U shaped and 25cm long.
- The duodenum contains two glands
 The liver and the pancreas

The liver

This produces bile juice

Bile juice is stored in the gall bladder

Bile juice is then directed to the duodenum by the bile duct

Bile juice helps to emulsify fats or break down fats for easy digestion

N.B Bile juice contains no enzyme

The pancreas produces pancreatic juice.

Pancreatic juice

This contains three enzymes

- -lipase
- -trypsin
- -amylase

Amylase completes the digestion of carbohydrates (starch) which began in the mouth

Lipase acts on fats and turns them into fatty acids and glycerol

Trypsin acts on peptides and remaining proteins turning them into amino acids

N.B 1. No absorption of food in the body takes place in the duodenum

2. Chemical digestion of fats begins in the duodenum

Food in the ileum

Digestion of food ends in the ileum

The ileum is about 6 metres long

Absorption of food takes place in the ileum

Absorption is a process by which digested food is taken into the blood stream.

It uses the villi to absorb food nutrients

The villi are finger-like structures covering the wall of the ileum

Adaptations of the ileum to absorbing digested food

It has villi hat absorb digested food

It has thin walls for easy diffusion of digested food

It has a network of blood capillaries

It is long and coiledincreasing a surface area for food absorption

The digestive juice produced in the ileum

Succus entericus

Enzymes contained in succus entericus

- maltase

- lipase

lactase

- peptidase (erepsin)

sucrase

Maltase turns maltose to glucose

Lactase turns lactose to glucose

Sucrase turns sucrose to glucose

Lipase turns lipids into fatty acids and glycerol

Peptidase (erepsin) turns peptides to amino acids

Summary

Class of foodEnd product of digestion

Proteins amino acids

Carbohydrates glucose

Fats and oils (lipids) fatty acids and glycerol

Note: The end product in the digestion of protein is amino acids, digestion of fats is fatty acids and glycerol and in the digestion of carbohydrates.

Uses of digested food by our bodies

- The maltose, glucose (from carbohydrates) fatty acids and glycerol (from fats) provide energy to the body
- Amino acids (from proteins) assist in the proper growth and repair of the body tissues.

The colon

- The function of the large intestine (colon) is to absorb water and some salts are removed from their fluid wastes are absorbed into the blood stream.
- The rest of the fluid waste and undigested solid food particles are sent to rectum.

Rectum

- The rectum stores undigested food as waste matter (faeces)
- The rectum also stores worn out cells from the digestive tract.

Anus

Passes out faeces

N.B the process by which un digested food is passed out the body is called egestion

Importance of roughage

It prevents constipation

It stimulates peristalsis

It reduces the risk of getting bowel cancer.

Activity

- 1. Explain the term absorption of food?
- 2. Where does absorption of food take place?
- 3. How is the ileum adapted to food absorption?
- 4. Where does absorption of the following foods take place?
 - a. Alcohol
 - b. Water
- 5. Where does digestion of fats take place in the mouth?
- 6. Write down things that happen to food in the mouth?
- 7. By what process does food move down the gullet?
- 8. Identify the two enzymes produced in the stomach.
- 9. State the major function of the following parts of digestive system.
 - a. Gall bladder
 - b. liver
- 10. Name tee enzyme that breaks down carbohydrates into maltose

- 11. What is the major role of bile produced by the liver in the digestive process?
- 12. Identify the major function of a spleen.
- 13. Identify the digestive juice produced by
 - a. pancreas
 - b. stomach

Summary table of digestive processes

Organ	Gland	Digestive juice	Enzymes	Food changes
Mouth	Salivary	saliva	Ptyalin or	Starch to
	glands		salivary	maltose
			amylase	
Stomach	Gastric glands	Gastric juice	pepsin	Proteins to
				peptides
			Rennin	clot milk in
				babies
Duodenum	liver	bile	No enzyme	Emulsifies the
				fats
	pancreas	Pancreatic	Trypsin	Proteins and
		juice		peptides to
				amino – acids
			Amylase	Starch to
				maltose
			Lipase	fats to fatty
				acids and
				glycerol
Ileum		Intestinal juice	Lactase	Lactose to
		or succus	Maltase	glucose
		entericus		Maltose to
				glucose
			Sucrase	Sucrose to
				glucose
			Lipase	Fats to fatty
				acids and
				glycerol
			peptidase	Peptides to
				amino - acids

colon		Absorption of
		water
Rectum		Storage of
		faeces

<u>Digestive disorders</u>(disturbance to alimentary canal and make it fail to function properly)

Constipation

- This is a condition where by a person finds it difficult to pass out faeces. The faeces are very hard and dry and do not come out easily

Causes

- lack of roughages in one's diet
- having irregular meals
- drinking too little water
- lack of exercise to the body

Indigestion

- This happens when food is not properly digested
- A person feels stomach pains, heart burn (burning in the chest) and tiredness.

Causes

- No chewing food properly
- Eating so hurriedly
- Too much drinking of alcohol

Intestinal obstruction

- Caused when intestines twist or fold themselves causes vomiting, thirst and death if not reported immediately to the doctor.

Vomiting

This is the expulsion of undigested food through the mouth

- This is a sign of very many diseases.
- It disturbs the digestive system and forces the cardiac sphincter to open and the food in the stomach to be ejected through the mouth.

Diseases of the digestive system

The diseases associated with the alimentary canal include;

- Appendicitis

- cholera

- Diarrhoea

- Peptic ulcers

- Typhoid

Appendicitis

- This is the inflammation of the appendix.
- This may be due to stones or other indigestible solids that get trapped in the appendix.
- See the doctor for medical attention.

Peptic ulcers

- These are sores in the stomach wall caused by too much aid.
- The victim experiences a chronic sharp pain in the stomach and frequent heart burn.

Dysentery

- It is caused by a bacterium or Amoeba got by eating uncooked food or dirty food or water.
- The victim visits the toilet very frequently, passing out watery stool with some blood.

Typhoid

- Caused by bacteria which cause inflammation of the intestine and easily lead to death.
- It is spread through drinking contaminated water

Ways through which food is made dirty

- Handling food with dirt hands.
- House flies landing on food with their hairy bodies.
- Use of dirty containers to keep food.
- Leaving food uncovered.
- Preparing food in dirty places.

Good eating habits

- Washing hands before serving or eating food..
- Chewing food properly before swallowing.
- Avoid talking while eating food
- Washing fruits/vegetables before eating them (germs).

How to improve on the working condition of digestive system.

- Have daily physical exercises.
- Avoid eating stale or rotten food.
- Having a balanced diet.
- Avoid drinking too much alcohol.
- Have regular meals.
- Always eat the right quantity of food.
- Avoid eating while talking.
- Drink safe clean water

Activity

- 1. Write down three diseases and three disorders of digestive system.
 - a. diseases
 - b. disorders
- 2. Suggest any two ways through which food we eat gets contaminated.
- 3. Write down at least two examples of food eating habits.
- 4. Why should food be covered immediately after being oared?
- 5. How are house flies adapted to spreading of diseases?