

SECTION A (40 Marks)

1a) • Dangerous liquids (acids)

- sharp objects
- High tables
- Dark rooms

b) - Teacher spelling out rules and regulations
- Learning operation of apparatus before using it
- use of apparatus common to the learner

2

Any two correct responses

3

Any three correct responses

2. B weighs more.

- Iron is denser than water.
- From mass = volume \times density
- Extra mass of B compared to A is, due to extra mass due to higher density of iron.

3a) - Conduction

- Convection
- radiation

b) • lagging using insulators around the container to prevent conduction
• covering to prevent convection
• Using highly polished dishes

3

Award 1 score for each response (correct)

2

Award 1 score for each correct response

4a) • Refraction, as light from the coin bends at the interface and appears to come from a near point.

b) • Reduce on amount of water, then real depth of coin from surface reduces, and hand can reach the coin


3

1 score for stating refraction
1 score for the effect of refraction
1 score for the conclusion
Award 1 score for each correct response


2

5a) Distance, = $80 + 60 = 140\text{m}$

b)




$$x = \sqrt{80^2 + 60^2} = 100\text{m}$$



6 • Sagging so as to allow room for expansion during hot weather or, contraction during cold weather, without risking contact of wires

5

7.



$$\Sigma = 450 \times 5, \quad \Sigma 750 \times x$$

$$G = E$$

$$\Rightarrow 750x = 450 \times 5$$

$$x = 3\text{m}$$

Correct diagram (1 score)

Correct statement of formula (1 score)

Substituting 2 correct answers with unit (3 score)

- - Sharp pointed metals (are spikes),
- Working as lightning conductors,
- To repel electrons due to electrostatic induction with the cloud
- Transfer the electrons via the copper strip and plate to the earth.
- So as to minimise effects of lightning and thunder

5

Award 1 score for each correct response regardless of order.

SECTION B (60 Marks) :-

a) Time taken, From 9:25am to 9:30am, 5 minutes
 $= 5 \times 60 = 300 \text{ s}$ ✓

$$u = \frac{36 \times 1000}{60 \times 60} = 10, \quad v = \frac{144 \times 1000}{60 \times 60} = 40 \quad \checkmark$$

$$a = \frac{v - u}{t} = \frac{40 - 10}{300} = 0.1 \text{ m s}^{-2} \quad \checkmark$$

$$s = \left(\frac{v + u}{2} \right) t = \left(\frac{40 + 10}{2} \right) \times 300 = 7500 \text{ m} \quad \checkmark$$

b) i) A faces;

- Supporting extra load of wheel barrow.
- Experiencing extra effects of friction.

B faces;

- Hard labour, no use of machine
- Bigger health risks due to contact with dusty bricks

(ii) Distance of ;

$$A: \left(\frac{500}{30} \times 2 \right) - 1 \Rightarrow 17 \times 2 - 1 = 33 \text{ times} \quad \checkmark$$

$$D = 33 \times 100 = 3300 \text{ m} \quad \checkmark$$

$$B: \left(\frac{500}{8} \times 2 \right) - 1 \Rightarrow 63 \times 2 - 1 = 125 \text{ times} \quad \checkmark$$

$$D = 125 \times 100 = 12500 \text{ m} \quad \checkmark$$

$$\text{Extra distance} = 12500 - 3300 = 9200 \text{ m} \quad \checkmark$$

7

Conversion from km/hr to m/s is a must.

4

Any 4 correct responses

- 10 a) (i) • Both are upright.
 • Both are same size.

2

- (ii) • Image is virtual, object is real.
 • Image is laterally inverted, object is proper.

2


- (ii) • Image would be smaller than object ✓
 • Image would be nearer to mirror than object. ✓

(i) height = 10cm ✓

(ii) $m = \frac{V}{u} = \frac{20}{20} = 1$ ✓

Nature of Image formed

- Inverted ✓
- Same size as object ✓ Any two correct responses
- real



Subject Teacher Signature:-

1a) $F = 60 \times 10 = 600 \text{ N}$ ✓

(i) $A = 0.002 \times 2 = 0.004 \text{ m}^2$


$P = \frac{F}{A} = \frac{600}{0.004} = 150,000 \text{ Pa}$ ✓

(ii) $A = 0.002$,

$P = \frac{600}{0.002} = 300,000 \text{ Pa}$ ✓

b(i) - Pain due to high pressure, ✓ ✓
- Falling due to unstable equilibrium

(ii) Damage due to high pressure. ✓



- Drilling holes at levels A, B, C of the bottle
- Covering holes with Cellotape.
- Pouring water in bottle above hole C
- Opening the holes by removing Cellotape
- Observing water pouring furthest from A due to high pressure at deeper point.
- Water pouring nearest from C due to low pressure at shallow point.

6.

Score 1 for each correct step with or without a diagram.

12 a) (i) They are longitudinal.

- They cannot pass through a vacuum
- They travel in straight lines
- They are Mechanical

(ii) Frequency of sound by tiptoeing thief is infra (lower) than sound in range of audibility (20 Hz - 20 KHz) but the dog's ear can detect such infra sound.

b(i) - Current flows, hammer hits the gong, sound is heard ✓

(ii) - Air is removed gradually, sound fades, and is not heard at all when all air is removed. Conclusion? ✓

c(i) - Echo as sound is reflected by wall. ✓

(ii) $2d = st$ ✓

$s = \frac{2d}{t} = \frac{2 \times 640}{4} = 320 \text{ m/s}$ ✓

3

1 Score for each correct response

3

Score 1 for each correct response

1

3

3 a(i) - A magnet is a metal that attracts certain metals.

• Magnetic materials are those that can be turned into a magnet.

(ii) - Hammering, midway roughly ✓
- Electrical with a.c. ✓
- Heating strongly. ✓

b) Current flows through the coil, the nail becomes magnetised. ✓

• Current is out of coil, nail loses magnetism

ii) • Once it acquires magnetism, it is not lost even when current is switched off.

Conclusion? ✓
d) • The magnet swings seeking the north and south, being attracted by the earth's magnetic field.

- When it comes to rest, the north pole is attracted to the south pole hence facing geographic north.

- The south pole is attracted to the north pole of the earth hence facing geographic south.

14 a) Covid-19 was a sickness that was more risky when the medical personnel is in contact with patient.

- Temperature gun could take measurements without making contact between user and patient.

(ii) - Readings affected by temperature of the surrounding ✓
- Readings affected by voltage of cells used. ✓

b) (i) - Alcohol ✓
- Mercury. ✓

(ii) Alcohol

- It expands readily ✓
- Low freezing point. ✓

Mercury

- Does not vaporize or distil in glass ✓
- Cohesive, does not stick on glass ✓

(iii) • Bring it in contact with a warm part.
• Liquid expands into the bore.
• Taking length L_1 , L_0 is obtained as length, L_{100} at boiling point and L_0 at the temperature of patient.

$$X = \frac{L_1 - L_0}{L_{100} - L_0} \times 100^\circ \text{C}$$

A learner may also explain the steps in Calibration of Thermometer