


Topics to be covered



- 1) Travelling Wave
- 2) Wave Interference
- 3) Standing Wave



Travelling Wave Equation



Wave vel. → Medium only

Wave freq. → Source only

Wave shape → Source only
(waveform)

$$f(x, t) = f\left(0, t - \frac{x}{v}\right)$$




Eq. of travelling wave towards +ve x-axis →

$$y(x, t) = y(0, t - \frac{x}{v}) = f(t - \frac{x}{v})$$

Eq. of travelling wave towards -ve axis →

$$y(x, t) = y(0, t + \frac{x}{v}) = f(t + \frac{x}{v})$$

Sinusoidal wave →



$$\begin{aligned}y(x, t) &= y(0, t - \frac{x}{v}) \\&= A \sin \left[\omega \left(t - \frac{x}{v} \right) \right]\end{aligned}$$

$$y = A \sin(\omega t - kx)$$

$$\text{where, } k = \frac{\omega}{v}$$

$$v = \frac{\omega}{k}$$

Wave propagation ~~in~~ direction.

$$y = A \sin(\underline{\omega t - kx})$$

$\frac{\omega}{v} = k$

+ve x-axis

$$y = A \sin(\underline{kx - \omega t})$$

+ve x-axis.

$$y = A \sin(\underline{\omega t + kx})$$

-ve x-axis

$$y = -A \sin(\underline{kx + \omega t})$$

-ve x-axis.

QUESTION

Which of the following equations represents a traveling wave?

[24 Feb, 2021 (S-II)]

A

$$y = A \sin x \cos \omega t \text{ } \color{magenta}{\times}$$

B

$$y = A \sin (15x - 2t)$$

C

$$y = Ae^x \cos (\omega t - \theta) \text{ } \color{magenta}{\times}$$

D

$$y = Ae^{-x^2} (vt + \theta) \text{ } \color{magenta}{\times}$$

QUESTION

If $y = (6x - 30t)^2$ then Find speed of waves.

$$\frac{6x}{\downarrow k} - \frac{30t}{\downarrow \omega}$$

$$v = \frac{\omega}{k} = \frac{30}{6} = 5 \text{ m/s}$$

$$f(t - \frac{x}{v})$$

$$f(vt - x)$$

ω = kendnya Vidhyalaya

$$\omega = kv$$

$$y = A \sin(kx - \omega t)$$

$$v = \frac{\omega}{k}$$



QUESTION



If $y = \frac{1}{(4x+3t)^2 + 1}$ then find vel. of wave along with direction.

$$\cancel{4x^2 + 3t}$$

$$\cancel{4x - 3t^2}$$

$$\cancel{4x^2 + 3t^2}$$

$$\frac{4x + 3t}{\bar{k}} = \frac{\omega}{\bar{\omega}}$$

$$V = \frac{\omega}{k} = \frac{3}{4} \text{ m/s}$$

towards - ve
x-axis

QUESTION

The amplitude of wave disturbance propagating in the positive x -direction is given by

$y = \frac{1}{1+(x)^2}$ at $t = 0$ and $y = \frac{1}{1+(x-2)^2}$ at $t = 1\text{s}$, where x and y are in metres. The shape of wave does not change during the propagation. The velocity of the wave will be 2 m/s.

[20 July, 2021 (S-I)]

$$y = \frac{1}{1+(x-2)^2} \quad \text{at } t=1$$

$$v = \frac{\omega}{k} = \frac{2}{1} = 2 \text{ m/s}$$



Wave Parameters



$$y = A \sin(kx - \omega t + \phi)$$

A = Amplitude

k = Angular wave number

ω = Angular frequency

ϕ = Phase const.

$kx - \omega t + \phi$ = Phase

Time period

Wave vel.

Diagram illustrating wave parameters:


- Wavelength (λ)
- Frequency (f)
- Time period (T)

$$\lambda = \frac{2\pi}{k}$$
$$T = \frac{2\pi}{\omega}$$
$$f = \frac{1}{T} = \frac{\omega}{2\pi}$$

$$v = \frac{\omega}{k} = \frac{\lambda}{T} = \lambda f$$



Wave Velocity, Particle Velocity and Slope



$$y = A \sin(kx - \omega t)$$

$$v_{\text{particle}} = \frac{dy}{dt} = -A\omega \cos(kx - \omega t)$$

$$\tan \theta = \frac{dy}{dx} = \frac{\partial y}{\partial x} = \frac{Ak \cos(kx - \omega t)}{\downarrow \text{max. slope}}$$

QUESTION

Marathon Q.

If $y = 4 \sin (\underline{k}x - \frac{\pi}{2}t)$ then
find

1. Amplitude of wave ? = 4 m

2. Wavelength = ? 2π

3. Frequency = ? $\frac{1}{4}$ Hz

4. Wave speed = ? $\frac{1}{2}$ m/s

5. Speed of particle at $x = 1$ at $t = 3$ s.
= 0 m/s.

2) $\lambda = \frac{2\pi}{k} = \frac{2\pi}{\pi} = 2m$

3) $f = \frac{\omega}{2\pi} = \frac{\pi/2}{2\pi} = \frac{1}{4}$ Hz

4) $v = \frac{\omega}{k} = \frac{\pi/2}{\pi} = \frac{1}{2} m/s$

5) $v = \frac{dy}{dt} = 4 \cos(\pi x - \frac{\pi}{2}t) \times -\frac{\pi}{2}$

$v = -2\pi \cos(\pi x - \frac{\pi}{2}t)$

$|v_{x=1, t=3}| = 2\pi \cos(\pi \times 1 - \frac{\pi}{2} \times 3)$
= $2\pi \cos(-\frac{\pi}{2}) = 0 m/s$



QUESTION

If $y = 4 \sin(\pi x - \frac{\pi}{2}t)$ then
find

6. Position of particle at $t = 1$ at $x = 2$.

7) 7) 7. Snapshot of wave at $t = 0.5$ s. 

8. Position of particle at $x = 0.5$ m at any time.

9. Slope of tangent on wave at $x = 1$ at $t = 1$. $= 0$

10. Max. particle velocity. $\Rightarrow V_{max} = A\omega = 4 \times \frac{\pi}{2} = 2\pi \text{ m/s}$

11. Phase diff. between two particles separated by 0.5 m.

$$11) \Delta\phi = \frac{\pi}{2} \times 0.5 = \frac{\pi}{2} A\omega$$

$$6) y_{x=2, t=1} = 4 \sin\left(\pi \times 2 - \frac{\pi}{2} \times 1\right) \\ = 4 \sin\left(\frac{3\pi}{2}\right) \\ = -4 \text{ m}$$

$$7) y_{t=0.5} = 4 \sin\left(\pi x - \frac{\pi}{2} \times \frac{1}{2}\right) \\ = 4 \sin\left(\pi x - \frac{\pi}{4}\right)$$


$$8) y_{x=0.5} = 4 \sin\left(\pi \times \frac{1}{2} - \frac{\pi}{2} t\right) \\ = 4 \sin\left(\frac{\pi}{2} - \frac{\pi}{2} t\right) \\ y_{x=0.5} = 4 \cos\left(\frac{\pi}{2} t\right)$$

$$y = 4 \sin\left(\pi x - \frac{\pi}{2} t\right)$$

g) $\tan \theta = \frac{\partial y}{\partial x} = 4 \cos\left(\pi x - \frac{\pi}{2} t\right) \times \pi$

$$\begin{aligned} (\tan \theta)_{x=1, t=0} &= 4 \pi \cos\left(\pi \cdot 1 - \frac{\pi}{2} \cdot 0\right) \\ &= 0 \end{aligned}$$

Relation b/w Phase difference & Path difference



$$\Delta\phi = \frac{2\pi}{\lambda} \times \Delta x$$

Path Difference

$$\lambda$$

$$2\pi$$

Phase difference

$$1$$

$$\frac{9\pi}{\lambda}$$

$$\Delta x$$

$$\frac{2\pi}{\lambda} \Delta x = \Delta\phi$$

QUESTION



A wave travelling in the positive x-direction having displacement along y-direction as 1 m, wavelength 2π m and frequency of $\frac{1}{\pi}$ Hz is represented by

- A $y = \sin(x - 2t)$
- B $y = \sin(2\pi x - 2\pi t)$
- C $y = \sin(10\pi x - 20\pi t)$
- D ~~$y = \sin(2\pi x + 2\pi t)$~~

$$y = A \sin(kx - \omega t)$$

$$= 1 \sin(1 \times x - 2t)$$

Common mistake

$$\lambda = \frac{2\pi}{k} \Rightarrow k = \frac{2\pi}{\lambda}$$

$$k = \frac{2\pi}{2\pi}$$

$$k = 1$$

$$f = \frac{\omega}{2\pi} \Rightarrow \omega = 2\pi f$$

$$= 2\pi \times \frac{1}{\pi}$$

$$f = 2$$

QUESTION

The wave described by

$$y = 0.25 \sin(10\pi x - 2\pi t),$$

where, x and y are in metre and t in second, is a wave travelling along the

- A negative x -direction with frequency 1Hz
- B positive x -direction with frequency π Hz and wavelength $l = 0.2\text{m}$
- C positive x -direction with frequency 1Hz and wavelength $l = 0.2\text{m}$
- D negative x -direction with amplitude 0.25m and wavelength $l = 0.2\text{m}$

QUESTION

A transverse wave is represented by $y = A \sin(\omega t - kx)$. For what value of the wavelength is the wave velocity equal to the maximum particle velocity?

A $\pi A/2$

$$V = \frac{\omega}{k}$$

$$V_{\max} = A\omega$$

B πA

$$\cancel{\frac{\omega}{k}} = A\cancel{\omega} \Rightarrow k = \frac{1}{A}$$

C $2\pi A$

$$\lambda = \frac{2\pi}{k} = \frac{2\pi}{\left(\frac{1}{A}\right)} = 2\pi A$$

D A

QUESTION

A transverse wave propagating along x-axis is represented by

$$y(x, t) = 8 \sin \left(\underline{0.5\pi x} - \frac{4\pi t}{\omega} - \frac{\pi}{4} \right)$$

where, x is in metre and t is in second. The speed of the wave is

- A** 4π m/s

$$v = \frac{\omega}{k} = \frac{4\pi}{0.5\pi} = 8 \text{ m/s}$$

- B** 0.5π m/s

- C** $\pi/4$ m/s

- D** 8 m/s

QUESTION

A travelling harmonic wave is represented by the equation $y(x, t) = 10^{-3} \sin (\underline{50t} + \underline{2x})$, where x and y are in meter and t is in seconds. Which of the following is a correct statement about the wave?

[JEE Mains 2019]

- A The wave is propagating along the negative x -axis with speed 25 ms^{-1} .
- B The wave is propagating along the positive x -axis with speed 100 ms^{-1} .
- C The wave is propagating along the positive x -axis with speed 25 ms^{-1} .
- D The wave is propagating along the negative x -axis with speed 100 ms^{-1} .

$$\begin{aligned}v &= \frac{\omega}{k} \\&= \frac{50}{2} \\&= 25 \text{ m/s}\end{aligned}$$

QUESTION

A wave travelling in the positive x-direction having displacement along y-direction as 1 m, wavelength 2π m and frequency of $\frac{1}{\pi}$ Hz is represented by

A

$$y = \sin(x - 2t)$$

Repeated

B

$$y = \sin(2\pi x - 2\pi t)$$

C

$$y = \sin(10\pi x - 20\pi t)$$

D

$$y = \sin(2\pi x + 2\pi t)$$

QUESTION

The wave described by

$$y = 0.25 \sin (10\pi x - 2\pi t),$$

where, x and y are in metre and t in second, is a wave travelling along the

reject

- A negative x -direction with frequency 1Hz
- B positive x -direction with frequency π Hz and wavelength $l = 0.2\text{m}$
- C positive x -direction with frequency 1Hz and wavelength $l = 0.2\text{m}$
- D negative x -direction with amplitude 0.25m and wavelength $l = 0.2\text{m}$

QUESTION

The equation of waves is given by $Y = 10^{-2} \sin 2\pi \frac{\omega}{k} t - 0.5x + \pi/4$ where x and Y are in m and t in s . The speed of the wave is _____ km h⁻¹. [11 April, 2023 (S-I)]

$$V = \frac{\omega}{k} = \frac{160}{0.5} = 320 \frac{m}{s}$$

$$320 \times \frac{10^3 \text{ km}}{3600 \text{ hr}}$$

$$320 \times 36 \times 10^3 \frac{\text{km}}{\text{hr}}$$

$$\frac{32 \times 36}{192} \frac{\text{km}}{\text{hr}}$$

$$\underline{\underline{961}} \frac{\text{km}}{\text{hr}}$$

QUESTION



A longitudinal wave is represented by $x = 10 \sin 2\pi (\Omega t - x/\lambda) \text{ cm}$. The maximum particle velocity will be four times the wave velocity if the determined value of wavelength is equal to:

[29 June, 2022 (S-I)]

- A 2π
- B 5π
- C π
- D $5\pi/2$

$$\begin{aligned}x &= 10 \sin \left[2\pi \left(\Omega t - \frac{x}{\lambda} \right) \right] \\&= 10 \sin \left(\underbrace{2\pi \Omega t}_{\omega} - \underbrace{\frac{(2\pi)x}{\lambda}}_{k} \right)\end{aligned}$$

$$\begin{aligned}v_{\text{particle}} &= A\omega \\&= 10 \times 2\pi \Omega \\&= 20\pi \Omega\end{aligned}$$

$$v = \frac{\omega}{k} = \frac{\cancel{2\pi}\Omega}{\cancel{2\pi}} = \Omega \times \lambda$$

$$\begin{aligned}v_{\text{particle}} &= 4 \times v \\20\pi \Omega &= 4 \times \cancel{\Omega} \lambda \\5\pi &= 4 \times \cancel{\Omega} \lambda \\5\pi &= 4 \times \lambda \\5\pi &= \lambda\end{aligned}$$

$$\lambda = 5\pi$$



Wave Velocity



$$y = A \sin(kx \pm \omega t)$$


$$v_{\text{wave}} = \frac{\omega}{k} = \frac{\lambda}{T} = \lambda f$$

Depends on medium only.



Transverse Wave Velocity

in a string



$$v = \sqrt{\frac{T}{\mu}}$$

T = Tension in string

μ = Linear mass density
(mass per unit length)

QUESTION

A Steel wire with mass per unit length 7.0×10^{-3} kg is under tension of 70 N. The speed of transverse waves in the wire will be:

[01 Feb, 2023 (S-I)]

- A 200π m/s
- B 100 m/s
- C 10 m/s
- D 50 m/s

$$v = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{70 \times 10^3}{7.0 \times 10^{-3}}} = \sqrt{10^4} = 100 \text{ m/s}$$

QUESTION

$$\mu = 0.135 \times \frac{10^{-3} \text{ kg}}{1 \text{ m}} = 0.135 \times 10^{-3} \frac{\text{kg}}{\text{m}}$$



The mass per unit length of a uniform wire is 0.135 g/cm . A transverse wave of the form $y = -0.21 \sin(x + 30t)$ is produced in it, where x is in meter and t is in second. Then, the expected value of tension in the wire is $x \times 10^{-2} \text{ N}$. Value of x is _____.
 (Round-off to the nearest integer) [26 Feb, 2021 (S-I)]

$$v = \frac{\omega}{k} = \frac{30}{1} = 30 \text{ m/s}$$

$$v = \sqrt{\frac{T}{\mu}} \Rightarrow 30 = \sqrt{\frac{T}{0.135 \times 10^{-3}}}$$

$$\Rightarrow T = \frac{30^2}{0.135 \times 10^{-3}}$$

$$\begin{aligned} T &= 99 \times 0.135 \times 10^6 \\ &= 12.15 \text{ N} \\ &= 12.15 \times 10^{-2} \text{ N} \\ &\text{Ans} \end{aligned}$$

QUESTION

Adv. level



A uniform string of length 20 m is suspended from a rigid support. A short wave pulse is introduced at its lowest end. It starts moving up the string. The time taken to reach the supports is:

- A $2\sqrt{2} \text{ s}$
- B $\sqrt{2} \text{ s}$
- C $2\pi\sqrt{2} \text{ s}$
- D 2s

$$v = \sqrt{\frac{T}{\mu}}$$

$$v = \sqrt{\frac{\mu x g}{\mu}}$$

$$v = \sqrt{g x}$$

$$\Rightarrow \frac{dx}{dt} = \sqrt{x g}$$

$$T = m_x g$$

$$T = (\mu x) g$$

$$\int_0^{20} \frac{dx}{\sqrt{x}} = \int_0^t \sqrt{g} dt$$


$$\left[\frac{x^{1/2}}{\left(\frac{1}{2}\right)} \right]_0^{20} = \sqrt{g} t$$

$$2\sqrt{2} = \sqrt{g} t$$

$$t = 2\sqrt{2} \lambda$$



[2016]



QUESTION



A uniform thin rope of length 12 m and mass 6 kg hangs vertically from a rigid support and a block of mass 2 kg is attached to its free end. A transverse short wave-train of wavelength 6 cm is produced at the lower end of the rope. What is the wavelength of the wave-train (in cm) when it reaches the top of the rope?

[Sep. 03, 2020 (I)]


A 3

$$f_1 = f_2$$

$$\frac{v_1}{\lambda_1} = \frac{v_2}{\lambda_2}$$

$$\frac{\sqrt{\frac{2g}{\mu}}}{6} = \frac{\sqrt{\frac{8g}{\mu}}}{\lambda_2}$$

$$\lambda_2 = 6 \times \sqrt{\frac{8}{2}} = 12\text{ cm}$$



QUESTION

The percentage increase in the speed of transverse waves produced in a stretched string if the tension is increased by 4%, will be%,

[25 Feb, 2021 (S-II)]

$$v = \sqrt{\frac{T}{\mu}}$$

$$\begin{aligned}\% \Delta v &= \frac{1}{2} \times \% \Delta T \\ &= \frac{1}{2} \times 4\% \\ &= 2\%\end{aligned}$$

$$y = x^n$$
$$\% \Delta y = n \times \% \Delta x$$

(for small % change)



Longitudinal Wave

(Sound)

$$\Delta x = A \sin(kx - \omega t)$$

$$\text{or } s = A \sin(kx - \omega t)$$

$$v = \frac{\omega}{k} = \frac{\lambda}{T} = \lambda f$$



Longitudinal Wave Velocity



$$V \propto \sqrt{\frac{\text{Elasticity}}{\text{Inertia}}}$$

Solid \rightarrow
(solid)

* $V = \sqrt{\frac{Y}{\rho}}$

Y - Young's modulus
of elasticity

Liquid \rightarrow

* $V = \sqrt{\frac{B}{\rho}}$

B - Bulk modulus of
elasticity

Gases →

Speed of sound (v_g)

$$\gamma = \frac{C_p}{C_v} = 1 + \frac{2}{f}$$

Mono → f = 3

Dial. → f = 5

Poly → f = 5
(linear) γ = Adiabatic exponent

Poly → f = 6
(non-linear) P = Pressure of gas
ρ = vol. density

$$v_g = \sqrt{\frac{B}{\rho}} = \sqrt{\frac{\gamma P}{\rho}} = \sqrt{\frac{\gamma RT}{M}}$$

Newton's Law

& Laplace correction

{Depends only
on T}

T = Absolute temp.

M = Molar mass

↓
for const temp.,

Speed of sound does
not depend on pressure
& vol.

Effect of Temp & Humidity



$$v = \sqrt{\frac{RT}{M}}$$

① Temp →

$$v \propto \sqrt{T}$$

② Humidity → Humidity ↑ → v ↑

Atmosphere



QUESTION

Diatomic, f = 5

Find ratio of speed of sound in O₂ & H₂ at same temperature.

$$\frac{V_{O_2}}{V_{H_2}} = \frac{\sqrt{\frac{\gamma R T}{M_{O_2}}}}{\sqrt{\frac{\gamma R T}{M_{H_2}}}} \Rightarrow \frac{V_{O_2}}{V_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}} = \sqrt{\frac{2}{32}} = \sqrt{\frac{1}{16}} = \frac{1}{4} \text{ Ans}$$

QUESTION

Two mono atomic ideal gases 1 and 2 of molecular masses m_1 and m_2 respectively are enclosed in separate containers kept at the same temperature. The ratio of the speed of sound in gas 1 to that in gas 2 is given by [2000S]

A $\sqrt{\frac{m_1}{m_2}}$

$$v_1 = \sqrt{\frac{\gamma RT}{m_1}}$$

B $\sqrt{\frac{m_2}{m_1}}$

$$v_2 = \sqrt{\frac{\gamma RT}{m_2}}$$

C $\frac{m_1}{m_2}$

D $\frac{m_2}{m_1}$




Pressure Wave

Displacement wave $\rightarrow \Delta x = A \sin(\omega t - kx)$

Pressure wave $\rightarrow \Delta P = P_0 \cos(\omega t - kx)$


$$P_0 = BAk$$

Disp. amplitude
Bulk modulus
Ang. wave no.





Energy Transfer in String



$$m = \mu \times \delta = \mu v t$$

Energy of small element = $\frac{1}{2} k A^2$

$$\int dE = \int \frac{1}{2} dm \omega^2 A^2$$

$$E = \frac{1}{2} m \omega^2 A^2$$

$$E = \frac{1}{2} (\mu v t) \omega^2 A^2$$

$$E = \frac{1}{2} \mu v \omega^2 A^2 t$$

Power = $\frac{\text{Energy}}{\text{time}} = \frac{\frac{1}{2} \mu v \omega^2 A^2 t}{t}$

$P \propto A^2$


$\frac{1}{2} \mu v \omega^2 A^2$





Intensity

$$I = \frac{\text{Energy}}{\text{time} \times \text{Area}}$$



$$I = \frac{\text{Power}}{\text{Area}}$$

$$I \propto A^2$$

waves


Wave optics

Amplitude of wave

QUESTION

A point source emits sound equally in all directions in a non-absorbing medium. Two points P and Q are at distance of 2m and 3m respectively from the source. The ratio of the **intensities** of the waves at P and Q is

- A** 9 : 4
- B** 2 : 3
- C** 3 : 2
- D** 4 : 9



$$E_{S_1} = E_{S_2}$$

$$\frac{I_{S_1}}{I_{S_2}} = \frac{\frac{E_{S_1}}{A_1}}{\frac{E_{S_2}}{A_2}} = \frac{A_2}{A_1}$$

$$\frac{I_{S_1}}{I_{S_2}} = \frac{\cancel{E_{S_1}}}{\cancel{E_{S_2}}} \cdot \frac{A_2}{A_1}$$

$$\frac{I_{S_1}}{I_{S_2}} = \frac{\cancel{E_{S_1}}}{\cancel{E_{S_2}}} \cdot \frac{\frac{4\pi R_2^2}{4\pi R_1^2}}{\frac{3^2}{2^2}} = \frac{9}{4}$$

Ans.



Wave Reflection and Refraction




Reflection →


Case I:- Rigid end




Wave will get
inverted after
reflection from
rigid end .




Case II:- Reflection from free end -



Wave does not
get inverted
upon reflection
from free end.






$$\begin{aligned}\mu_1 &< \mu_2 \\ v_1 &> v_2 \\ \lambda_1 &> \lambda_2 \\ A_1 &< A_0 \\ A_2 &< A_0\end{aligned}$$

Denser $\xrightarrow{\hspace{1cm}}$ Rarer



$$\begin{aligned}\mu_1 &> \mu_2 \\ v_1 &< v_2 \\ \lambda_1 &< \lambda_2\end{aligned}$$



QUESTION

Sound waves travel at **350 m/s** through a warm air and at **3500 m/s** through brass.
The wavelength of a 700 Hz acoustic wave as it enters brass from warm air.

10x

$$\lambda = \frac{V}{f}$$

↓ ↓
10x 10x

- A increases by a factor 20
- B increases by a factor 10
- C decreases by a factor 20
- D decreases by a factor 10



Wave Interference



① Waves of same frequency travelling in same direction

② Waves of same frequency and amplitude travelling in opposite direcn

(Standing Waves)

③ Waves of different frequencies travelling in same direcn

(Beat)

(Sound)



Wave Interference



$$\nu_1 = \nu_2$$

$$\omega_1 = \omega_2$$

$$\lambda_1 = \lambda_2 \Rightarrow k_1 = k_2$$

Case 1: Interference of waves of same frequency travelling in same direction:


$$y_1 = A_1 \sin(kx - \omega t)$$

$$y_2 = A_2 \sin(kx - \omega t + \phi)$$

$$y = y_1 + y_2 = A \sin(kx - \omega t + \delta)$$

$$A = \sqrt{A_1^2 + A_2^2 + 2A_1 A_2 \cos \phi}$$

$$\tan \delta = \frac{A_2 \sin \phi}{A_1 + A_2 \cos \phi}$$



$$I \propto A^2$$

$$I \propto A_1^2 + A_2^2 + 2A_1 A_2 \cos \phi$$

$$I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \phi$$

Special Cases① Constructive Interference ($\phi = 0$) \rightarrow

$$y_1 = A_1 \sin(kx - \omega t) \quad \text{--- I}_1$$

$$y_2 = A_2 \sin(kx - \omega t) \quad \text{--- I}_2$$

$$y = y_1 + y_2 = (A_1 + A_2) \sin(kx - \omega t)$$


$$I \propto (A_1 + A_2)^2$$

$$I = (\sqrt{I_1} + \sqrt{I_2})^2$$

$$T = T_1 + T_2 + 2\sqrt{T_1 T_2}$$

$y_1 = A \sin(kx - \omega t)$	I
$y_2 = A \sin(kx - \omega t)$	I
$y = 2A \sin(kx - \omega t)$	$4I$

$\rightarrow A_2 \rightarrow A_1$



② Destructive Interference ($\phi = \pi$) \rightarrow

$$y_1 = A_1 \sin(kx - \omega t) \quad \text{--- } I_1$$

$$y_2 = A_2 \sin(kx - \omega t + \pi) \quad \text{--- } I_2$$

$$= -A_2 \sin(kx - \omega t)$$

$$y = y_1 + y_2 = (A_1 - A_2) \sin(kx - \omega t)$$

$$I \propto (A_1 - A_2)^2$$

$$I = (\sqrt{I_1} - \sqrt{I_2})^2$$

$$I = I_1 + I_2 - 2\sqrt{I_1 I_2}$$



$$y_1 = A \sin(kx - \omega t) \quad I$$

$$y_2 = A \sin(kx - \omega t + \pi) \quad I$$

$$y = 0$$

QUESTION

If $A_1 : A_2 = 3 : 2$ then find the ratio of I_{\max}/I_{\min} upon interference of two waves.

$$\frac{I_{\max}}{I_{\min}} \propto \frac{(A_1 + A_2)^2}{(A_1 - A_2)^2} \Rightarrow \frac{I_{\max}}{I_{\min}} = \frac{(3A + 2A)^2}{(3A - 2A)^2} = \frac{25}{1} \text{ Ans}$$

QUESTION

There harmonic waves having equal frequency ν and same intensity I_0 have phase angles $0, \frac{\pi}{4}$ and $-\frac{\pi}{4}$ respectively. When they are superimposed the intensity of the resultant wave is close to:

A $5.8 I_0$

B $0.2 I_0$

C $3 I_0$

D I_0


$$A_{\text{net}} = A_0 + A_0\sqrt{2}$$

$$A_{\text{net}} = A_0(1 + \sqrt{2})$$

$$I_{\text{net}} \propto A_{\text{net}}^2$$

$$I_{\text{net}} \propto [A_0(1 + \sqrt{2})]^2$$

$$I_{\text{net}} = I_0 (1 + \sqrt{2})^2 \Rightarrow I = (2.4)^2 I_0 \quad A_0$$



QUESTION

$$y_1 = 5 \sin\left(\frac{2\pi}{\lambda}x - 2\pi v t\right)$$

$$\lambda = \frac{2\pi}{2\pi} = 1$$

The equations of two waves are given by:

$$y_1 = 5 \sin 2\pi(x - vt) \text{ cm}$$

$$y_2 = 3 \sin 2\pi(x - vt + 1.5) \text{ cm.} = 3 \sin(2\pi x - 2\pi v t + 3\pi)$$

These waves are simultaneously passing through a string. The **amplitude** of the resulting wave is:

[24 June, 2022 (S-I)]

A 2 cm

B 4 cm

C 5.8 cm

D 8 cm



$$1 \longrightarrow 2\pi$$

$$1.5 \longrightarrow 3\pi$$

QUESTION

Amplitude of


⑥ Ans

Find the resultant wave upon interference of following two moves-

$$y_1 = 6 \sin(\pi x - \pi/2 t)$$

$$y_2 = 6 \sin(\pi x - \pi/2 t + 2\pi/3)$$

120°



$$\begin{aligned}A_{\text{net}} &= \sqrt{A^2 + A^2 + 2A^2 \cos 120^\circ} \\&= \sqrt{2A^2 + 2A^2 \left(-\frac{1}{2}\right)} \\&= A\end{aligned}$$

QUESTION



Two coherent sources produce waves of different intensities which interfere. After interference, the ratio of the maximum intensity to the minimum intensity is 16. The intensity of the waves are in the ratio

[9 Jan, 2019 (S-I)]

A 16 : 9

B ✓ 25 : 9

C 4 : 1

D 5 : 3

$$\frac{\frac{I_{\max}}{I_{\min}} = \frac{16}{1}}{\frac{(\sqrt{I_1} + \sqrt{I_2})^2}{(\sqrt{I_1} - \sqrt{I_2})^2} = \frac{16}{1}}$$

$$\Rightarrow \frac{\sqrt{I_1} + \sqrt{I_2}}{\sqrt{I_1} - \sqrt{I_2}} = 4$$

$$\Rightarrow \sqrt{I_1} + \sqrt{I_2} = 4\sqrt{I_1} - 4\sqrt{I_2}$$

$$\Rightarrow 3\sqrt{I_1} = 5\sqrt{I_2}$$

$$\Rightarrow \sqrt{\frac{I_1}{I_2}} = \frac{5}{3} \Rightarrow \frac{I_1}{I_2} = \frac{25}{9}$$



Wave Interference



Case 2: Interference of waves of same frequency and amplitude travelling in opposite direction: (STANDING WAVE)

Posn. of nodes

$$\sin(kx) = 0$$

$$kx = n\pi$$

$$x = \frac{n\pi}{k}$$

$$y_1 = A \sin(kx - \omega t)$$

$$y_2 = A \sin(kx + \omega t)$$

$$y = y_1 + y_2 = \underbrace{2A \sin(kx) \cos(\omega t)}_{A_s} \quad \text{Eq. of Standing wave.}$$

Posn. of Anti-nodes

$$\sin(kx) = \pm 1$$

$$kx = (2n+1)\frac{\pi}{2}$$

$$x = (2n+1)\frac{\pi}{2k}$$

$$v = \sqrt{\frac{T}{\mu}}$$

Standing Wave in String
fixed at both ends



① First Harmonic (Fundamental Harmonic) →



$$L = \frac{\lambda}{2} \Rightarrow \lambda = 2L$$

$$f = \frac{v}{\lambda} = \frac{v}{2L}$$

fundamental frequency

Total no. of nodes = 2 Points of nodes = 0, L

Total no. of anti-nodes = 1 Points of anti-nodes = $\frac{L}{2}$



② 2nd Harmonic (1st overtone) →



$$\leftarrow \text{L} \rightarrow$$



$$L = 2 \frac{\lambda}{2} \Rightarrow \lambda = L$$

$$f = \frac{v}{\lambda} = \frac{2v}{2L}$$

Total no. of Nodes = 3

Total no. of Anti-nodes = 2

Posn. of nodes = $0, \frac{L}{2}, L$

Posn. of Anti-nodes = $\frac{L}{4}, \frac{3L}{4}$

③ 3rd Harmonic (2nd overtone) →

$$L = \frac{3\lambda}{2} \Rightarrow \lambda = \frac{2L}{3}$$


$$f = \frac{v}{\lambda} \Rightarrow f = \frac{3v}{2L}$$

Total no. of nodes = 4

Total no. of anti-nodes = 3

Posn. of nodes = 0, $\frac{L}{3}, \frac{2L}{3}, \frac{3L}{3}$

Posn. of anti-nodes = $\frac{L}{6}, \frac{3L}{6}, \frac{5L}{6}$




① n^{th} Harmonic ($(n-1)^{\text{*}}$ overtone) \rightarrow

$$L = \frac{n\lambda}{2} \rightarrow \lambda = \frac{2L}{n}$$

$$f = \frac{v}{\lambda} \rightarrow f = \frac{n v}{2L}$$

Total no. of nodes = $n+1$

Total no. of anti-nodes = n



Posn. of nodes = $0, \frac{L}{n}, \frac{2L}{n}, \frac{3L}{n}, \dots, \frac{nL}{n}$

Posn. of anti-nodes = $\frac{L}{2n}, \frac{3L}{2n}, \frac{5L}{2n}, \dots, \frac{(2n-1)L}{2n}$

$$V = \sqrt{\frac{\gamma P}{g}}$$

Standing wave in open organ pipe -

① First Harmonic (Fundamental Harmonic) →

Disp. wave



$$L = \frac{\lambda}{2} \Rightarrow \lambda = 2L$$

$$f = \frac{V}{\lambda} = \frac{V}{2L}$$



Pressure wave



① n^{th} Harmonic $\left[(n-1)^{\text{th}} \text{ overtone} \right] \rightarrow$



$$L = \frac{n\lambda}{2} \Rightarrow \lambda = \frac{2L}{n}$$

$$f_n = \frac{v}{\lambda} = \frac{n v}{2L}$$




$$v = \sqrt{T/\mu}$$

Standing wave in string free at one end
(only odd Harmonics are detected)

① first Harmonic (fundamental Harmonic) →

$$L = \frac{\lambda}{4} \Rightarrow \lambda = 4L$$

$$f = \frac{v}{\lambda} = \frac{v}{4L}$$



Total no. of nodes = 1

Total no. of anti-nodes = 0


Total no. of anti-nodes = 1

Total no. of anti-nodes = L

② 3rd Harmonic (1^{st} overtone) \rightarrow

$$L = \frac{3\lambda}{4} \Rightarrow \lambda = \frac{4L}{3}$$

$$f = \frac{v}{\lambda} = \frac{3v}{4L}$$



Total no. of nodes = 2

Posn. of nodes = 0, $\frac{2L}{3}$

Total no. of anti-nodes = 2

Posn. of anti-nodes = $\frac{L}{3}, \frac{3L}{3}$

③ 5^{th} Harmonic (2^{nd} overtone) \rightarrow

$$L = \frac{5\lambda}{4} \Rightarrow \lambda = \frac{4L}{5}$$


$$f = \frac{v}{\lambda} = \frac{5v}{4L}$$

Total no. of nodes = 3

Total no. of anti-nodes = 3

Posn. of Nodes = 0, $\frac{2L}{5}$, $\frac{4L}{5}$

Posn. of anti-nodes = $\frac{L}{5}$, $\frac{3L}{5}$, $\frac{5L}{5}$




$(2n+1)^{\text{th}}$ Harmonic (n^{th} overtone) \rightarrow



$$n = 0, 1, 2, \dots$$

$$L = (2n+1) \frac{\lambda}{4} \rightarrow \lambda = \frac{4L}{2n+1}$$

$f = \frac{v}{\lambda} = \frac{(2n+1)v}{4L}$



$$\text{Total no. of nodes} = n+1$$

$$\text{Total no. of anti-nodes} = n+1$$

$$\text{Posn. of nodes} = 0, \frac{2L}{2n+1}, \frac{4L}{2n+1}, \dots, \frac{2nL}{2n+1}$$

$$\text{Posn. of anti-nodes} = \frac{L}{2n+1}, \frac{3L}{2n+1}, \frac{5L}{2n+1}, \dots, L$$

$$V = \sqrt{\frac{Y\rho}{\beta}}$$


Standing wave in closed organ pipe →



① First Harmonic (fundamental Harmonic)

$$L = \frac{\lambda}{4} \Rightarrow \lambda = 4L$$

$$f = \frac{V}{\lambda} = \frac{V}{4L}$$



$(2n+1)^{\text{th}}$ Harmonic (n^{th} overtone) —

$$f = \frac{(2n+1)V}{4L}$$

QUESTION

Two travelling waves of equal amplitudes and frequencies move in opposite directions along a string. They interfere to produce a stationary wave whose equation is given by

$$y = \left(\underbrace{10 \cos \pi x}_{\text{A}} \sin \frac{2\pi t}{T} \right).$$

The amplitude of the particle at $x = 4/3 \text{ cm}$ will be ____ cm.

[24 June, 2022 (S-II)]

$$A_x = 10 \cos(\pi x)$$

$$A_{x = \frac{4}{3}} = 10 \cos\left(\pi \times \frac{4}{3}\right)$$

$$\approx 10 \cos(240^\circ)$$

$$\approx |10 \cos 60^\circ|$$

$$\approx |5|$$

QUESTION

4 त्रिकौपिका



A string is clamped at both the ends and it is vibrating in its 4th harmonic. The equation of the stationary wave is $Y = 0.3 \sin(0.157x) \cos(200t)$. The length of the string is: (All quantities are in SI units.)

क्

[9 April, 2019 (S-I)]

A 20 m**B** 80 m**C** 60 m**D** 40 m

$$L = 4 \times \frac{\lambda}{2}$$

$$= 2\lambda$$

$$= 80 \text{ m}$$

$$Y = 0.3 \sin(kx) \cos(\omega t)$$

$$\lambda = \frac{2\pi}{k} = \frac{2\pi}{0.157}$$

$$= \frac{2 \times 3.14}{1.57 \times 10^{-1}}$$

$$\lambda = 40 \text{ m}$$


QUESTION

Two travelling waves produces a standing wave represented by equation, $y = 1.0 \text{ mm} \cos(1.57 \text{ cm}^{-1})x \sin(78.5 \text{ s}^{-1})t$. The node closest to the origin in the region $x > 0$ will be at $x = \underline{\hspace{2cm}}$ cm.

[26 Aug, 2021 (S-I)]

$$y = 1 \cos\left(\frac{\pi}{2}x\right) \sin(\omega t)$$

$$\cancel{\frac{\pi}{\lambda}} x = \cancel{\frac{\pi}{\lambda}}$$
$$x = 1 \text{ cm}$$



QUESTION

The two nearest harmonics of a tube closed at one end and open at other end are 220 Hz and 260 Hz. What is the fundamental frequency of the system?

- A 10 Hz
- B 20 Hz
- C 30 Hz
- D 40 Hz

$$f = \frac{v}{4L} = \frac{40}{2} = 20 \text{ Hz}$$

Diagram showing the first four harmonics of a tube closed at one end and open at the other. The wavelengths are labeled: $\lambda_1 = \frac{v}{4L}$, $\lambda_2 = \frac{3v}{4L}$, $\lambda_3 = \frac{5v}{4L}$, and $\lambda_4 = \frac{7v}{4L}$. The frequencies are: $f_1 = \frac{v}{4L}$, $f_2 = \frac{3v}{4L}$, $f_3 = \frac{5v}{4L}$, and $f_4 = \frac{7v}{4L}$.

QUESTION

ONE OR MORE THAN ONE OPTION(S) MAY BE CORRECT



If we study the vibration of a pipe open at both ends, which of the following statements is/are not true?

- A Open end will be pressure antinode - *false*
- B Odd harmonics of the fundamental frequency will be generated - *false*
- C All harmonics of the fundamental frequency will be generated - *true*
- D Pressure change will be maximum at both ends - *false*

QUESTION

The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is

- A** 12.5 cm
- B** 8 cm
- C** 13.3 cm
- D** 16 cm

$$\frac{v}{2L_1} = \frac{3v}{4L_2}$$

$$\frac{1}{2\pi L_1} = \frac{3}{4 \times 20}$$

$$L_1 = \frac{80^{\circ}}{2 \times 3}$$

$$L_1 = 13.3 \text{ cm}$$

QUESTION


A wave of frequency 100 Hz is sent along a string towards a fixed end. When this wave travels back after reflection, a node is formed at a distance of 10 cm from the fixed end of the string. The speeds of incident (and reflected) wave are

- A** 5 m/s
- B** 10 m/s
- C** 20 m/s
- D** 40 m/s

$$\frac{\lambda}{2} = 10 \text{ cm} \rightarrow \lambda = 20 \text{ cm}$$

$$v = \lambda \times f$$

$$\therefore v = 20 \times 10^{-2} \times 100 \\ \therefore v = 20 \text{ m/s}$$



QUESTION

A closed organ pipe (closed at one end) is excited to support the third overtone. It is found that air in the pipe has

- A** three nodes and three antinodes
- B** three nodes and four antinodes
- C** four nodes and three antinodes
- D** four nodes and four antinodes

Common mistake

$(2n+1)^{\text{th}}$ Harmonic

$$2 \times 3 + 1 = 7^{\text{th}} \text{ Harmonic}$$



1st Harmonic



3rd Harmonic - 1st O.T



5th - 2nd O.T



7th Harmonic - 3rd O.T

QUESTION

9

An organ pipe 40 cm long is open at both ends. The speed of sound in air is 360 ms^{-1} .
The frequency of the second harmonic is ____ Hz.

[08 April, 2023 (S-II)]

$$f_2 = \frac{2v}{\lambda L} = \frac{360}{4 \times 10^{-2}}$$
$$= 900 \text{ Hz.}$$

QUESTION

$$\mu = \frac{m}{L} = \frac{2 \times 10^{-5}}{1} = 2 \times 10^{-5}$$



A string of length 1 m and mass 2×10^{-5} kg is under tension T. When the string vibrates, two successive harmonics are found to occur at frequencies 750 Hz and 1000 Hz. The value of tension T is 5 Newton. [JEE Adv, 2023]

$$750 \text{ Hz} \quad 250 = \frac{v}{2L} \Rightarrow v = 500 \times 1$$

$$1000 \text{ Hz} \quad v = 500 \frac{\text{m}}{\text{s}}$$

$$\sqrt{\frac{T}{\mu}} = 500$$

$$\frac{T}{2 \times 10^{-5}} = (500)^2 \Rightarrow T = 2 \times 10^{-5} \times 25 \times 10^4 \\ = 50 \times 10^{-1} = 5 \text{ N}$$

$$\left[\begin{array}{l} \frac{v}{2L} \longrightarrow 1^{\text{st}} \\ \frac{2v}{2L} \longrightarrow 2^{\text{nd}} \\ \frac{3v}{2L} \longrightarrow 3^{\text{rd}} \end{array} \right]$$

Ans.

QUESTION

A guitar string of length 90 cm vibrates with a fundamental frequency of 120 Hz. The length of the string producing a fundamental frequency of 180 Hz will be ____ cm.

[08 April, 2023 (S-II)]

$$f = \frac{v}{2L} \Rightarrow v = f \times 2L$$
$$= 120 \times 2 \times 90 \text{ cm/s}$$

$$v = 24 \times 9 \text{ m/s}$$

$$180 = \frac{24 \times 9}{2L}$$

$$L = \frac{24 \times 9}{180 \times 2} \text{ m}$$

$$= \frac{24 \times 9}{180 \times 2} \times 10^{-2} \text{ m} = 60 \text{ cm}$$

QUESTION

For a certain organ pipe, the first three resonance frequencies are in the ratio of 1 : 3 : 5 respectively. If the frequency of fifth harmonic is 405 Hz and the speed of sound in air is 324 ms⁻¹ the length of the organ pipe is ___ m. [12 April, 2023 (S-I)]

$$f_5 = \frac{5v}{4L}$$

$$405 = \frac{5 \times 324}{4 \times L}$$


$$L = \frac{5 \times 324}{4 \times 405} = 1$$



Resonance



$$f_{\text{ext}} = f_{\text{standing wave}}$$



QUESTION

A string is stretched between fixed points separated by 75.0 cm. It is observed to have resonant frequencies of 420 Hz and 315 Hz. There are no other resonant frequencies between these two. The lowest resonant frequency for this strings is:

- A 155 Hz
- B 205 Hz
- C 10.5 Hz
- D 105 Hz

$$\begin{aligned}f &= f_n - f_{n-1} \\&\approx 420 - 315 \\&= 105 \text{ Hz}\end{aligned}$$

QUESTION

A wire of length L and mass per unit length $[6.0 \times 10^{-3} \text{ kgm}^{-1}]$ is put under tension of 540 N . Two consecutive frequencies that it resonates at are: 420 Hz and 490 Hz . Then L in meters is:

[9 Jan. 2020 (II)]

- A 2.1 m
- B 1.1 m
- C 8.1 m
- D 5.1 m

$$f = 490 - 420 = 70 \text{ Hz}$$

$$L = \frac{3\pi}{140} = 2.1 \text{ m}$$

$$\frac{g}{2L} = 70$$

$$\frac{1}{2L} \sqrt{\frac{g}{\mu}} = 70$$

$$L = \frac{1}{140} \times \sqrt{\frac{540 \cdot 9.8}{6 \times 10^{-3}}}$$


$$\frac{\sqrt{9 \times 10^4}}{140}$$



Sonometer



$$f_0 = \frac{v}{2L}$$



QUESTION

The length of the string of a musical instrument is **90 cm** and has a fundamental frequency of **120 Hz**. Where should it be pressed to produce fundamental frequency of

180 Hz?

A 75 cm

B 60 cm

C 45 cm

D 80 cm

$$f = \frac{v}{2L}$$

$$v = 2fL$$

$$f_1 L_1 = f_2 L_2$$

$$\cancel{120} \times \cancel{90} = \cancel{180} \times L_2$$

$$L = 60 \text{ cm}$$

QUESTION



If n_1 , n_2 and n_3 are the fundamental frequencies of three segments into which a string is divided, then the original fundamental frequency n of the string is given by

- A** $\frac{1}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$
- B** $\frac{1}{\sqrt{n}} = \frac{1}{\sqrt{n_1}} + \frac{1}{\sqrt{n_2}} + \frac{1}{\sqrt{n_3}}$
- C** $\sqrt{n} = \sqrt{n_1} + \sqrt{n_2} + \sqrt{n_3}$
- D** $n = n_1 + n_2 + n_3$

$$n_1 = \frac{v}{2L_1} \Rightarrow L_1 = \frac{v}{2n_1}, \quad n_2 = \frac{v}{2L_2}, \quad n_3 = \frac{v}{2L_3}$$



$$L = L_1 + L_2 + L_3$$

$$\frac{v}{\rho n} = \frac{v}{\rho n_1} + \frac{v}{\rho n_2} + \frac{v}{\rho n_3}$$

$$\frac{1}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}$$

QUESTION

A stretched string resonates with tuning fork of frequency **512 Hz** when length of the string is **0.5 m**. The length of the string require to vibrate resonantly with a tuning fork of frequency **256 Hz** would be

- A** 0.25 m
- B** 0.5 m
- C** 1 m
- D** 2 m

$$f = \frac{v}{2L} \Rightarrow v = 2fL$$
$$\cancel{512} \times 0.5 = \cancel{256} \times L$$
$$L = 1 \text{ m}$$

QUESTION

In an experiment with sonometer when a mass of 180 g is attached to the string, it vibrates with fundamental frequency of 30 Hz. When a mass m is attached, the string vibrates with fundamental frequency of 50 Hz. The value of m is 50 g.

[13 April, 2023 (S-II)]



$$f = \frac{v}{2L} = \frac{1}{2L} \sqrt{\frac{T}{\mu}}$$

$$f = \frac{1}{2L} \sqrt{\frac{mg}{\mu}}$$

$$\frac{f_1}{f_2} = \sqrt{\frac{m_1}{m_2}} \Rightarrow \frac{30}{50} = \sqrt{\frac{180}{m}}$$

$$\frac{9}{25} = \frac{180}{m}$$

$$m = \frac{180 \times 25}{9}$$

QUESTION




9

L

A wire of density $8 \times 10^3 \text{ kg/m}^3$ is stretched between two clamps 0.5m apart. The extension developed in the wire is 3.2×10^{-4} . If $Y = 8 \times 10^{10} \text{ N/m}^2$, the fundamental frequency of vibration in the wire will be ____ Hz.

[11 April, 2023 (S-II)]



$$\frac{f}{A} = \frac{Y \Delta \ell}{L}$$

$$f = \frac{AY \Delta \ell}{L}$$

$$v = \sqrt{\frac{T}{\mu}}$$

$$v = \sqrt{\frac{F}{\rho A}}$$

$$v = \sqrt{\frac{AY \Delta \ell}{\rho \times \rho A}} = \sqrt{\frac{Y \Delta \ell}{\rho L}}$$

$$m = \mu L = \rho \times A \times L$$

$$\mu = \rho A$$

$$f = \frac{v}{2L}$$

$$f = \frac{v}{2L}$$

$$= \frac{1}{2 \times 0.5} \sqrt{\frac{8 \times 10^1 \times 3.2 \times 10^{-4}}{8 \times 10^3 \times 0.5}}$$

$$\sqrt{6.4 \times 10^3}$$

$$\sqrt{6400} = 80 \text{ Hz}$$

QUESTION

$$\frac{\rho_1}{\rho_2} = \sqrt{\frac{1}{4}}$$



$$m = \mu \ell = \rho A \ell$$

$$\mu = \rho A$$



Two wires W_1 and W_2 have the same radius r and respective densities ρ_1 and ρ_2 such that $\rho_2 = 4\rho_1$. They are joined together at the point O , as shown in the figure. The combination is used as a sonometer wire and kept under tension T . The point O is midway between the two bridges. When a stationary waves is set up in the composite wire, the joint is found to be a node. The ratio of the number of antinodes formed in W_1 to W_2 is :

- A 1 : 1
- B 1 : 2
- C 1 : 3
- D 4 : 1

$$f_1 = f_2$$


$$\frac{\mu_1}{\mu_2} = \frac{\rho_1 \cancel{A}}{\rho_2 \cancel{A}}$$

$$\frac{n_1 \sqrt{\cancel{A}}}{\cancel{\mu_1}} = \frac{n_2 \sqrt{\cancel{A}}}{\cancel{\mu_2}}$$

$$n_1 \sqrt{\frac{1}{\mu_1}} = n_2 \sqrt{\frac{1}{\mu_2}}$$

$$\frac{n_1}{n_2} = \sqrt{\frac{\mu_1}{\mu_2}} = \sqrt{\frac{\rho_1}{\rho_2}} = \sqrt{\frac{1}{4}} = \boxed{\frac{1}{2}}$$


[Online April 8, 2017]





Resonance Tube

$$f_0 = \frac{(2n+1)\nu}{4L}$$




Closed organ pipe .

QUESTION

In a resonance tube experiment when the tube is filled with water up to a height of **17.0 cm** from bottom, it resonates with a given tuning fork. When the water level is raised the next resonance with **the same tuning fork** occurs at a height of **24.5 cm**. If the velocity of sound in air is **330 m/s**, the tuning fork frequency is: [Sep. 05, 2020 (I)]

- A 2200 Hz
- B 550 Hz
- C 1100 Hz
- D 3300 Hz



$$|L_2 - L_1| = \frac{\lambda}{2}$$

$$24.5 - 17 = \frac{\lambda}{2}$$

$$\lambda = 7.5 \times 2 = 15 \text{ cm}$$

$$f = \frac{v}{\lambda} = \frac{330}{15 \times 10^{-2}} = 22000 \text{ Hz}$$

QUESTION

A cylindrical resonance tube open at both ends, has a fundamental frequency f , in air. If half of the length is dipped vertically in water, the fundamental frequency of the air column will be

- A $2f$
- B $3f/2$
- C f
- D $f/2$

$$f = \frac{v}{2L}$$



$$f = \frac{v}{4(L/2)}$$




$$f = \frac{v}{2L}$$

QUESTION

The number of possible natural oscillations of air column in a pipe closed at one end of length **85 cm** whose frequencies lie below **1250 Hz** are (velocity of sound = 340 m^{-1})

- A** 4
- B** 5
- C** 7
- D** 6



$$\begin{aligned}f &= \frac{v}{4L} = \frac{340}{4 \times 85 \times 10^{-2}} \\&= \frac{340 \times 100}{340} = 100 \text{ Hz}\end{aligned}$$

$f_1, f_3, f_5, f_7, f_9, f_{11}, f_{13}$

QUESTION

A tuning fork with frequency **800 Hz** produces resonance in a resonance column tube with upper end open and lower end closed by water surface. Successive resonance are observed at length 9.75 cm, 31.25 cm and 52.75 cm. The speed of sound in air is

- A** 500 m/s
- B** 156 m/s
- C** 344 m/s
- D** 172 m/s

$$\begin{array}{l} 9.75 \text{ cm} \\ 31.25 \text{ cm} \\ 52.75 \text{ cm} \end{array} \quad \left. \begin{array}{l} \xrightarrow{\lambda/2} \\ \xrightarrow{\lambda/2} \end{array} \right\} = \begin{array}{l} 31.25 - 9.75 \\ \frac{\lambda}{2} = 21.5 \text{ cm} \\ \lambda = 43 \text{ cm} \end{array}$$

$$\begin{aligned} V &= \lambda \times f \\ &= 43 \times 10^{-2} \times 800 \\ &= 344 \text{ m/s.} \end{aligned}$$



Beats



Interference of waves having some frequency diff.

$$f_1$$

$$f_2$$

$$f_{\text{beat}} = |f_1 - f_2|$$

$$\begin{aligned} f_1 &= 200 \text{ Hz} \\ f_2 &= 204 \text{ Hz} \end{aligned} \rightarrow f_{\text{beat}} = 4 \text{ Hz}$$

QUESTION

Two sound waves with wavelengths 5 m and 5.5 m respectively, each propagate in a gas with velocity 330 m/s . We expect the following number of beat per second

A 12

$$f_1 = \frac{v}{\lambda} = \frac{330}{5} = 66\text{ Hz}$$

B zero

$$f_2 = \frac{v}{\lambda} = \frac{330}{5.5} = 60\text{ Hz}$$

C 1

$$f_{\text{beat}} = |f_2 - f_1| = 6\text{ Hz}$$

D 6

QUESTION

Two vibrating tuning forks produce progressive waves given by

$$y_1 = 4 \sin 500\pi t \text{ and}$$

$$y_2 = 2 \sin 506\pi t. \text{ Number of beat produced per minute is}$$

A 360

B 180

C 3

D 60

$$f_1 = \frac{\omega_1}{2\pi} = \frac{500\pi}{2\pi} = 250 \text{ Hz}$$

$$f_2 = \frac{506\pi}{2\pi} = 253 \text{ Hz}$$

$$f_{\text{beat}} = 3 \text{ Hz}$$

$$\begin{aligned}\text{No. of beats} &= f_{\text{beat}} \times t \\ &= 3 \times 60 \\ &= 180\end{aligned}$$

QUESTION

Two sources of sound placed closed to each other, are emitting progressive wave given by $y_1 = 4 \sin 600\pi t$ and $y_2 = 5 \sin 608\pi t$. An observer located near these two sources of sound will hear $f_1 = \frac{600\pi}{2\pi} = 300\text{Hz}$ $f_2 = \frac{608\pi}{2\pi} = 304\text{Hz}$.



$$\frac{I_{\max}}{I_{\min}} = \frac{(A_1 + A_2)^2}{(A_1 - A_2)^2}$$

$$= \frac{(4+5)^2}{(4-5)^2}$$

$$= 81 : 1$$

- A** 4 beat/s with intensity ratio 25 : 16 between waxing and waning
- B** 8 beat/s with intensity ratio 25 : 16 between waxing and waning
- C** 8 beat/s with intensity ratio 81 : 1 between waxing and waning
- D** 4 beat/s with intensity ratio 81 : 1 between waxing and waning

QUESTION

Two waves of wavelength **50 cm** and **51 cm** produce **12 beat/s.** The speed of sound is

A 306 m/s

$$f_1 = \frac{v}{\lambda_1} \quad f_2 = \frac{v}{\lambda_2}$$

B 331 m/s

$$f_1 - f_2 = 12$$

C 340 m/s

$$v \left[\frac{1}{\lambda_1} - \frac{1}{\lambda_2} \right] = 12$$

D 360 m/s

$$v = \frac{12}{\left[\frac{1}{50} - \frac{1}{51} \right]}$$




$$\begin{aligned} v &= \frac{12}{\left[\frac{1}{50} - \frac{1}{51} \right]} \\ &= \frac{6 \times 51}{1} = 306 \text{ m/s} \end{aligned}$$

QUESTION

$$f_{\text{beat}} = 2 \text{ Hz} \quad T = \frac{1}{f} = \frac{1}{2} \text{ s}$$



The correct figure that shows, schematically, the wave pattern produced by superposition of two waves of frequencies 9 Hz and 11 Hz is: [10 April, 2019 (S-II)]

A**B****C****D**





Doppler's Effect


(JEE Adv.)

Operating System
↓
Observer Source


$$f = f_0 \left[\frac{c + v_o}{c - v_s} \right]$$



c = speed of sound
in air



$$f = f_0 \left[\frac{c + v_o}{c + v_s} \right]$$



QUESTION

Two cars moving in opposite directions approach each other with speed of 22 m/s and 16.5 m/s respectively. The driver of the first car blows a horn having a frequency 400 Hz . The frequency heard by the driver of the second car is [velocity of sound 340 m/s]

A $\cancel{350 \text{ Hz}}$

B $\cancel{361 \text{ Hz}}$

C 411 Hz

D $\checkmark 448 \text{ Hz}$



$$f = f_0 \left[\frac{c + v_o}{c - v_s} \right]$$

$$= 400 \left[\frac{340 + 16.5}{340 - 22} \right] = 400 \times \frac{356.5}{318}$$

QUESTION

A siren emitting a sound of frequency 800 Hz moves away from an observer towards a cliff at a speed of 15 ms^{-1} . Then, the frequency of sound that the observer hears in the echo reflected from the cliff is (Take, velocity of sound in air = 330 ms^{-1})

- A 800 Hz
- B 838 Hz
- C 885 Hz
- D 765 Hz

$$f_0 = 800 \text{ Hz}$$

$$f_1 = f_0 \left[\frac{c + o}{c - v} \right] = f_0 \times \frac{c}{c - 15}$$

$$f_2 = f_1 \left[\frac{c + 15}{c} \right]$$

$$f_2 = 800 \times \frac{330}{315} \times \frac{c + 15}{c}$$

$$f_2 = 800 \times \frac{345}{315} \times \frac{609}{603}$$



Sound Properties



Loudness →

$$\text{Loudness} = 10 \log \frac{I}{I_0}$$

decibel
dB

$$I_0 = 10^{-12} \frac{\text{W}}{\text{m}^2}$$

QUESTION




A small speaker delivers **2W** of audio output. At what distance from the speaker will one detect **120 dB** intensity sound? [Given reference intensity of sound as 10^{-12}W/m^2] [12 April 2019 II]

A 40 cm

B 20 cm

C 10 cm

D 30 cm



$$I = \frac{E}{A} = \frac{2}{4\pi r^2}$$

$$120 = 10 \log_{10} \frac{I}{10^{-12}}$$

$$\rightarrow 10^{12} = \frac{I}{10^{-12}} \rightarrow I = 10^{24}$$

$$4\pi r^2 = \frac{2}{10^{-12}}$$

$$r = \sqrt{\frac{10^{-12}}{4\pi}} = \frac{10^{-6}}{\sqrt{2\pi}}$$

$$\Rightarrow \frac{2}{4\pi r^2} = 10^{24} = \frac{10^{-6}}{2.5} = 40$$

To get MDCAT important
Stuff and Official Updates
join our Group.

Whatsapp at 0314-3200007



MDCAT Study Buddy

Group · 2000 participants



Group call



Add



Search

MDCAT Entry Test Preparation.
Stay connected and get benefits.

Created by You, 11/06/2023, 2:47 pm