Candidate's Name :	• • • • • •	• • • • •	•••••	•••••	•••••	•••••	•••••	••••
C!		Ran	dom	No.		Pers	sonal	No.
Signature:								

(Do not write your School / Centre Name or Number anywhere on this booklet.)
P525/3
CHEMISTRY
(Practical)
Nov./Dec. 2022
3½ hours



UGANDA NATIONAL EXAMINATIONS BOARD

Uganda Advanced Certificate of Education

CHEMISTRY (PRACTICAL)

Paper 3

3 hours 15 minutes

INSTRUCTIONS TO CANDIDATES:

Answer all questions. Use blue or black ink. Any work done in pencil will not be marked except drawings.

All your answers must be written in the spaces provided.

Mathematical tables and silent non-programmable scientific calculators may be used.

Reference books (i.e. text books, booklets on qualitative analysis etc.) should **not** be used.

You are **not** allowed to start working with the apparatus for the first **15 minutes**. This time is to enable candidates read the question paper and make sure they have all the apparatus and chemicals that they may need.

For Examiners' Use Only						
Q.1	Q.2	Q.3	Total			
	1					

© 2022 Uganda National Examinations Board

Turn Over

1. You are provided with the following:

FA1, which is a solution of hydrochloric acid.

Metal X.

Substance Y, which is an oxide of X, with a formula XO.

You are required to determine the enthalpy change for the reduction of the oxide of X and comment on your answer.

PART I

Procedure

Weigh accurately about 1.2 g of X.

Using a measuring cylinder, transfer 100 cm³ of FA1 into a plastic beaker or cup. Read and record its initial temperature, in table 1.

Add at once the 1.2 g of X into FA1 in the plastic beaker or cup and at the same time, start the stop clock or watch.

Stir gently with the thermometer and record the temperature of the mixture after every half-minute in table 1, up to the fourth minute.

Results

Mass of weighing container $+ \mathbf{X} = \dots g$	$(\frac{1}{2} mark)$
Mass of weighing container alone =g	(½ mark)
Mass of X used =g	$(\frac{1}{2} mark)$

Table 1

Time (minutes)	0	1/2	1	1½	2	2½	3	3½	4
Temperature of solution (°C)				¥					73

 $(4\frac{1}{2} marks)$

PART II

Procedure

Weigh accurately about 2.0 g of Y.

Using a measuring cylinder, transfer 100 cm³ of FA1 into a plastic beaker or cup. Read and record its initial temperature, in table 2.

Add at once the 2.0 g of Y into FA1 in the plastic beaker or cup and at the same time, start the stop clock or watch.

Stir gently with the thermometer and record the temperature of the solution after every half-minute in table 2, up to the fourth minute.

Results


Mass of weighing container $+ Y = \dots g$	(½ mark)
Mass of weighing container alone =g	$(\frac{1}{2} mark)$
Mass of Y used =g	$\binom{1}{2}$ mark)

Table 2

Time (minutes)	0	1/2	1	1½	2	21/2	3	31/2	4
Temperature of solution (°C)	*: 		u r	-		11,,			

 $(4\frac{1}{2} \text{ marks})$

(a) (i) Plot on the same axes, graphs of temperature against time for results obtained in both **Part I** and **Part II**. (07 marks) (Graph paper is provided on page 4.)



				ım temperature chan
	for each reactio	n.		(03 mar
• • • • • • • • • • • • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •		
	• • • • • • • • • • • • • • • • • • • •			
		7 J		
	•••••	••••••		
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	•••••		
(iii)	Calculate the he	eat of reacti	ion in Part I an	d Part II.
	[Specific heat c	anacity of i	he solution =	$4.2 \text{ Jg}^{-1} \text{ K}^{-1}$ and its
		_		of the reactions of
			case, equation	of the reactions of 1
	and Y are as fo	niows:		
	$X_{(s)} + 2 HCl_{(a)}$	$q) \longrightarrow X$	$(Cl_{2(aq)} + H_{2(g)})$	
	$XO_{(s)} + 2HCl$	$(aq) \longrightarrow X$	$(Cl_{2(aq)} + H_2O)$	<i>[a]</i>
Part I:				(01 ma
	The state of the state of the state of		and the second of	Company to the same of the same of the same
	• • • • • • • • • • • • • • • • • • • •		1 2 7	
		, ()	dimension of the	
		.,\ \		
		.,\ ,		
Part II:				

	(1V)	($\mathbf{X} = 24$, $O = 16$)	ctions in 1 art 1 ar	(03 marks)
• • • •				
••••	••••••			
••••	•••••			
••••				
(b)	Deter	mine the heat energy change for	the reaction.	
	XO (s)	$+ H_{2(g)} \longrightarrow X_{(s)} + H_{2O(l)}$		(02 marks)
(c)		nent on your answer in (b).		(01 mark)
				•••••
•••••				

You are provided with substance **D** which contains **two** cations and **two** anions. You are required to carry out the tests in table 3 to identify the cations and anions in **D**. Identify any gas(es) evolved.

Record your observations and deductions in the table.

Table 3 (30 marks)

	TESTS	OBSERVATIONS	DEDUCTIONS
]	To two spatula end-fuls of D in a test tube, add dilute nitric acid until there is no further change.		
(b)	To two spatula end-fuls of D in a test tube, add about 5 cm ³ of distilled water, shake well and filter. Keep both the filtrate and the residue.		The hitse and histories of the form of the first of the firs
(c)	Divide the filtrate into three parts. (i) To the first part, add silver nitrate solution followed by dilute nitric acid.		
	(ii) To the second part, add aqueous iron(II) chloride solution.		

TESTS	OBSERVATIONS	DEDUCTIONS
(iii) Use the third part of the filtrate to carry out a test of your own choice to confirm one of the anions in D .		
Test:		
(d) Wash the residue twice with water. Transfer it into a test tube, add dilute nitric acid and warm. Add dilute sodium hydroxide solution drop-wise until in excess. Filter and keep both the filtrate and the residue.		
(e) To the filtrate from part (d), add dilute nitric acid drop-wise until the solution is just acidic. Divide the solution into three parts.		

THE CITY OF	OBSERVATIONS	DEDUCTIONS
(i) To the first part of the acidified filtrate, add dilute sodium hydroxide solution drop-wise until in excess.		
(ii) To the second part of the acidified filtrate, add dilute ammonia solution drop-wise until in excess.		
(iii) Use the third part of the acidified filtrate to carry out a test of your own choice to confirm one of the cations in D .		
Test:		
(f) Dissolve the residue in part (d) in a minimum amount of dilute nitric acid. Divide the resultant solution into three parts.		

TESTS	OBSERVATIONS	DEDUCTIONS
(i) To the first part, add dilute sodium hydroxide solution drop-wise until in excess.		
(ii) To the second part, add dilute ammonia solution drop-wise until in excess.		
(iii) Use the third part to carry out a test of your own choice to confirm the second cation in D . Test:		

(g) (i)	The cations in D are	and	
410			
(ii)	The anions in D are.	and	

3. You are provided with substance L, which is an organic compound. You are required to carry out the tests in table 4 to determine the nature of L. Record your observations and deductions in the table.

Table 4

(20 marks)

	TESTS	OBSERVATIONS	DEDUCTIONS
(a)	Burn a small amount of L on a spatula-end or in porcelain dish.	the state of the s	
(b)	To about 1 cm ³ of L, add about 2 cm ³ of distilled water. Shake the mixture and test with litmus. Divide the mixture into four parts. (i) To the first part, add 2-3 drops of 2, 4 - dinitrophenyl		Compared Control of the compared Control of the con
	hydrazine solution. (ii) To the second part, add a half a spatula endful of solid sodium		Albertage College
	hydrogencarbonate. (iii) To the third part, add 2-3 drops of neutral iron(III) chloride		
	(iv) To the fourth part, add 2-3 drops of acidified potassium dichromate solution and heat.		

an I		UCTIONS
ne		
