



# JINJA JOINT EXAMINATIONS BOARD

## MOCK EXAMINATIONS 2022

P530/1

### PROPOSED MARKING GUIDE

#### SECTION A

1 B	2 A	3 D	4 D	5 B	6 A	7 B	8 C	9 B	10 A
11 D	12 B	13 C	14 B	15 B	16 C	17 A	18 C	19 D	20 B
21 A	22 B	23 A	24 A	25 C	26 D	27 A	28 D	29 A	30 C
31 A	32 D	33 D	34 B	35 C	36 C	37 B	38 A	39 C	40 B

40 marks/ 1 mark each

#### SECTION B

41. (a)

Distance between parts	Line
Two poles of the cell	P;
A chromosome and a pole	R;
Two identical chromosomes	Q;

1 ½ marks / @ ½

(b) (i) 15 minutes;

1 mark

(ii) Distance between two identical chromosomes rapidly increase;

1 mark

(c) (i) line P

From 0 to 25 minutes, distance between pole remains constant at 40µm; because during metaphase and anaphase; centrosomes (centrioles) have reached opposite poles and stop moving;

1 ½ marks / @ ½

(ii) line Q

From 0 to 15 minutes, distance between two identical chromosomes is 0µm; because sister chromatids are still held together by the centromeres during metaphase;

From 15 to 25 minutes, distance between identical chromosomes rapidly increases; because spindle fibres split the chromatids and pull them towards opposite poles during anaphase;

2 marks / @ ½ (iii)  
line R

From 0 to 15 minutes, distance between chromosome and a pole remains constant at 20 µm; because the cell is in metaphase stage when chromosomes are at equator not moving;

From 15 to 25 minutes, the distance between chromosome and a pole gradually decreases; because after splitting during anaphase; the spindle fibres pull the chromosomes towards the poles;

3 marks / @ ½

42. (a) (i) species with population number (size) so low that they are considered to be in danger of becoming extinct (if the cause of their decline continues to operate); 1 mark

(ii) permanent condition of a species of having no living representative in the wild following the death of the last surviving individual of the species; 1 mark

- (b) -hunting (and poaching, overfishing);  
-deforestation destroying habitats;  
-industrialisation producing poisonous gases that pollute environments;  
-massive spraying of pesticides;  
-swamp reclamation for settlement and agriculture;  
-land fragmentation by road constructions destroying habitats;

(Any 4 suggestions) 4 marks / @ 1

- (c) -legally protect endangered species;  
-recycle waste materials like paper, glass bottles etc to avoid pollution;  
-prohibit release of non-native animals and plants into an area;  
- restrict trade in endangered species;  
- provide breeding programs for endangered species;  
- establish sperm banks and seed stores to maintain biodiversity;

- establish national parks, and nature reserves for protecting endangered species;
- restricting urban and industrial developments in natural habitats;

**(Any 4 suggestions) 4 marks / @ 1**

43. (a) direction away from 5 to the right i.e.(  $\longrightarrow$  ) ; **1 mark**

(b) (i) 1,2,3,4,5; **1 mark**

(ii) 6,7,8,9; **1 mark**

(c) Fully extended cilia move backwards more quickly; exerting greater resistance against water generating a forward force; that propels the organism forward; **3marks / @ 1**

(d) Used to create water currents that move food into gullets of paramecia during feeding;

Locomotion in ciliated microorganisms like paramecia;

Respiratory systems of man like the trachea contain cilia, which trap and remove germs and dust in inhaled air;

Ciliated cells in female human oviducts move eggs towards uterus for fertilization and implantation;

Ciliated male gametes in some lower plants like ferns use cilia to swim towards female gametes;

**(Any 4 suggestions) 4 marks / @ 1**

44. (a) species E because; E starts photosynthesizing at low(er) light intensity; and E reaches its maximum rate at low(er) light intensity;

**3 marks / @ 1**

**OR**

/ E steep(er) increase in rate of photosynthesis (with small increase in light intensity) ;

/ E has a , higher / greater / faster , rate of photosynthesis (than D) at low light intensities;

(b) shade leaf will have;

- 1 larger chloroplast(s) ;
- 2 more chloroplast(s) ;
- 3 more grana / thylakoids (in chloroplast) ;
- 4 larger surface area (of leaves) ;
- 5 more palisade mesophyll cells;
- 6 leaves with thinner lamina;
- 7 leaves with thinner cuticles;
- 8 leaves with smooth leaf lamina (non hairy leaves) ;

**(Any 4 comparisons) 4 marks / @ 1**

(c) Light energy excites electrons in chlorophyll molecules; in photosystems to higher energy levels to generate ATP molecules in the electron transport chain reactions;

Light energy splits; water molecules during photolysis; to produce hydrogen ions that are used to produce reduced NADP; and electrons to replace excited electrons from photosystem II;

**3 marks / @ ½**

45. (a) (no increase until after 0.8 – 1% ) / increasing CO<sub>2</sub> percentage from 0 to 1.5 % Slowly (slightly) increases rate of breathing;
- Increasing CO<sub>2</sub> from 1.5 to 5.4% gradually increases rate of breathing;
- Then increasing from 5.4 to 6% rapidly increases rate of breathing;

**3 marks / @ 1**

(b) An increase in the concentration of carbon dioxide in the blood; stimulates chemoreceptors; in the walls of the carotid artery and the aorta; sending impulses via vagus (sensory) nerve; to respiratory (ventilation) centre in medulla; More impulses (from medulla) ; to diaphragm / intercostals (muscles) ;

Increasing rate of (muscle) contraction (ventilation/breathing);

4 marks / @ ½

(c) During mouth-to-mouth resuscitation expired air contains about 4% (more) CO<sub>2</sub>; and this stimulates an increase in the patient's respiratory rate; enhancing quick recovery;

Pressing on chest wall will cause atmospheric air with only 0.04% ; (much lower) CO<sub>2</sub> to enter the patient's lungs which is not sufficient enough to stimulate the patient's respiratory rate; and recovery is therefore slower;

3 marks / @ ½

46. (a) (i) GgX<sup>R</sup>X<sup>r</sup> ; 1 mark

(ii) If it were recessive all flies of 3 and 4 would be grey ;

OR 3 and 4 grey parents produce black (fly) 9;

OR Grey parents produce black (fly) ;

1 mark

(b) 1. Flies 3 and 4 produce black fly 9 ;  
if (fly 3) X chromosome carried the gene for grey body colour and (Fly) 3 would pass dominant allele to 9;

2 marks / @ 1

OR

2. (Fly) 2 and 1 produce 5/grey (fly) ;(Black female produces grey male) ;

(Fly) 5 could not be grey as (Fly) 5 would receive recessive allele from 2 if it was carried on X chromosome;

(c) Genotypes of parents: GgX<sup>r</sup>X<sup>r</sup> × ggX<sup>R</sup>Y;

Genotypes of offspring GgX<sup>R</sup>X<sup>r</sup>, ggX<sup>R</sup>X<sup>r</sup>, GgX<sup>r</sup>Y and ggX<sup>r</sup>Y ;

Phenotypes of offspring: Grey-bodied red-eyed female, black-bodied red-eyed female, grey-bodied

white-eyed male, black-bodied white  
eyed male. ;

Ratio of phenotypes: 1 : 1 : 1 : 1 ;

4 marks / @ 1

*If 1, 2 and 3 incorrect allow one mark for correct gametes from incorrect dihybrid parental genotypes.*

(d)  $p^2 + 2pq = 0.64$   $q^2 = 1 - 0.64$  ;  
 $q^2 = 0.36$   
 $q = 0.6$   
 $p = 0.4$  ;  
Heterozygous flies =  $2pq = 2 \times 0.4 \times 0.6 = 0.48$  ;  
= 48% ;

2 marks / @ ½

END