

Candidate's Name:

Signature:

| Random No. | | | | | Personal No. | | |
|------------|--|--|--|--|--------------|--|--|
| | | | | | | | |

(Do not write your School /Centre Name or Number anywhere on this booklet.)

P525/1
CHEMISTRY
Paper 1
(Theory)
Nov./Dec. 2024
2¾ hours



UGANDA NATIONAL EXAMINATIONS BOARD

Uganda Advanced Certificate of Education

CHEMISTRY

Paper 1
(Theory)

2 hours 45 minutes

INSTRUCTIONS TO CANDIDATES:

This paper consists of **two** Sections; **A** and **B**.

Section **A** is **compulsory**. Attempt **six** questions from Section **B**. Any additional question(s) attempted will **not** be marked.

All questions **must** be answered in the spaces provided. Use **blue** or **black** ink. Any work done in pencil, **except** drawings, will not be marked.

The Periodic Table, with relative atomic masses, is attached at the end of the paper.

Mathematical tables (3-figure tables) are adequate or silent non-programmable scientific electronic calculators may be used.

Illustrate your answers with equations where applicable.

Where necessary, use the following:

Molar gas constant, $R = 8.31 \text{ JK}^{-1} \text{ mol}^{-1}$.

Molar volume of gas at s.t.p. is 22.4 litres.

Standard temperature = 273 K.

Standard pressure = 101325 Nm^{-2} .

| For Examiners' Use Only | | | | | | | | | | | | | | | | | |
|-------------------------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|-------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | Total |
| | | | | | | | | | | | | | | | | | |

SECTION A: (46 MARKS)*Answer all the questions in this section.*

1. (a) Define the term **standard electrode potential**. (01 mark)

.....

.....

.....

.....

- (b) The standard electrode potentials of some half cells reactions are given below:

| | E^{θ} (V) |
|--|------------------|
| $Co^{2+}(aq) + 2e \longrightarrow Co(s)$ | -0.28 |
| $Zn^{2+}(aq) + 2e \longrightarrow Zn(s)$ | -0.76 |
| $Cl_2(aq) + 2e \longrightarrow 2Cl^{-}(aq)$ | +1.36 |
| $MnO_4^{-}(aq) + 8H^{+}(aq) + 5e \longrightarrow Mn^{2+}(aq) + 4H_2O(l)$ | +1.52 |

- (i) Identify the strongest reducing agent and the strongest oxidising agent. (01 mark)

.....

- (ii) Write the cell notation of the cell constructed by combining the half cells $Co^{2+}(aq) / Co(s)$ and $Zn^{2+}(aq) / Zn(s)$. (01 mark)

.....


- (iii) Write the equation for the overall cell reaction for the cell in b(ii). (01 mark)

.....

- (iv) Calculate the e.m.f. of the cell in b(ii). (01 mark)

.....

.....



condensation polymerisation.

(a) State the meaning of the term condensation polymerisation.

(01 mark)

.....

.....

.....

(b) (i) Write the structural formula(e) and name(s) of the monomers of nylon 6, 6. (03 marks)

.....

.....

.....

(ii) State **one** use of nylon 6, 6.

(01 mark)

.....

3. Draw the structure and name the shape of each of the following ions in Table 1:

(4½ marks)

Table 1

| Ion | Structure | Name of shape |
|--------------------|-----------|---------------|
| NO_3^- | | |
| SO_3^{2-} | | |
| NH_4^+ | | |

4. Complete the following equations and name the major organic product:



Name



Name



Name



Name

5. The vapour pressure of propanone is 37330 Nm^{-2} at 30°C . When 33.4 g of cane sugar were dissolved in 120 g of propanone, the vapour pressure reduced by 1760 Nm^{-2} .

(a) Calculate the molar mass of cane sugar. (2½ marks)

.....

(b) Explain why the vapour pressure of the solution is lower than that of propanone. (1½ marks)

.....

.....

.....

.....

.....

6. (a) State what would be observed and write an equation for the reaction that takes place when dilute hydrochloric acid is added to aqueous potassium manganate(VI) solution.

Observation:

(1½ marks)

.....
.....

Equation:

(1½ marks)

.....
.....

- (b) Explain why potassium manganate(VII) is **not** acidified using hydrochloric acid. (2½ marks)


.....
.....
.....

7. Complete the following equations and in each case write the mechanism for the reaction(s):



Mechanism:

.....
.....
.....
.....



Mechanism:

.....
.....
.....
.....

8. (a) Write an equation for the reaction that takes place when methylamine is dissolved in water. (1½ marks)

.....

- (b) The hydrogen ion concentration of a 1M methylamine solution is $2.5 \times 10^{-13} \text{ mol l}^{-1}$. Calculate the base dissociation constant K_b , of methylamine. (3½ marks)

(The ionic product of water, $K_w = 10^{-14} \text{ mol}^2 \text{ l}^{-2}$)

.....

.....

.....

.....

.....

.....

.....

.....

.....

9. The first ionisation energies and atomic radii for some elements of group(II) of the Periodic Table are given in Table 2.

Table 2

| Element | Beryllium | Magnesium | Calcium | Strontium | Barium |
|---|-----------|-----------|---------|-----------|--------|
| First ionisation energy (kJ mol ⁻¹) | 899 | 738 | 590 | 550 | 503 |
| Atomic radius (nm) | 0.089 | 0.136 | 0.174 | 0.191 | 0.198 |

- (a) State how the first ionisation energy varies with atomic radius. (01 mark)

.....

.....

- (b) Explain your answer in (a). (04 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SECTION B: (54 MARKS)

Answer any six questions from this section.

Any additional question(s) answered will not be marked.

10. The molecular formula of a compound **P** is $C_3H_6Cl_2$.

- (a) Write the structural formulae of all the possible isomers of **P**. (02 marks)

.....

.....

.....

- (b) When **P** was boiled with aqueous sodium hydroxide, a compound **R** which reacted with hydroxylamine (NH_2OH) in the presence of an acid and reduced Fehling's solution, was formed.

- (i) Identify compounds **P** and **R**. (02 marks)

.....

.....

- (ii) Write a mechanism for the reaction between R and hydroxylamine. (05 marks)

.....

.....

.....

.....

.....

.....

11. (a) State two factors which affect the magnitude of lattice energy. (01 mark)

.....

- (b) Some thermochemical data are shown below:

Enthalpy of atomisation of chlorine = $+122 \text{ kJ mol}^{-1}$.

Enthalpy of atomisation of magnesium = $+148 \text{ kJ mol}^{-1}$.

First ionisation energy of magnesium = $+738 \text{ kJ mol}^{-1}$.

Second ionisation energy of magnesium = $+1451 \text{ kJ mol}^{-1}$.

Enthalpy of formation of magnesium chloride = -641 kJ mol^{-1} .

First electron affinity of chlorine = -364 kJ mol^{-1} .

- (i) Construct an energy level diagram for the formation of magnesium chloride. (03 marks)

.....

.....

.....

.....

.....

.....

(ii) Determine the lattice energy of magnesium chloride. (02 marks)

.....

.....

(iii) Calculate the enthalpy of solution of magnesium chloride. (03 marks)

(The hydration energies of magnesium ions and chloride ions are -1891 and -381 kJ mol^{-1} respectively.)

.....

.....

.....

12. (a) One of the characteristics of transition elements, is formation of complexes. Explain how transition metal ions form complexes. (1½ marks)

.....

.....

(b) The formulae of some complex ions are shown in Table 3. Write the name of the complex ion, its oxidation state and the co-ordination number of the central metal ion. (03 marks)

Table 3

| Complex ion | Name of ion | Oxidation state | Co-ordination number |
|---|-------------|-----------------|----------------------|
| $[(\text{Co}(\text{SCN})_4)]^{2-}$ | | | |
| $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$ | | | |

(c) The molecular mass of a salt, $\text{Cu}(\text{NH}_3)_x\text{Cl}_y \cdot z\text{H}_2\text{O}$ is 276. When a solution containing 1.38 g of salt was reacted with excess silver nitrate solution, 2.87 g of silver chloride was formed. Calculate the number of moles of chloride ions in 1 mole of the salt. (02 marks)

- (d) When a solution containing 1.38g of the salt was mixed with aqueous sodium hydroxide and heated, the ammonia liberated completely neutralised 10 cm^3 of a 1.0 M hydrochloric acid.


Calculate the number of moles of;

- (i) ammonia molecules in 1 mole of the salt. (1½ marks)

- (ii) water in 1 mole of the salt. (01 mark)


13. Write the equations to show how the following compounds can be synthesised and in each case, indicate the conditions for the reaction.

- (a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ from propanone. (04 marks)

- (b)  from benzene. (3½ marks)

- (c) $\text{CH}_3\text{CH}=\text{CH}_2$ from 2-bromopropane. (1½ marks)

14. Sulphur dioxide and oxygen can react to form sulphur trioxide according to the following equation:



3 moles of sulphur dioxide and 2 moles of oxygen were heated at 450 °C in a 1 dm³ vessel at 50 atmospheres. At equilibrium the vessel contained 20 % sulphur dioxide.

- (a) (i) Write the expression for the equilibrium constant, K_p . (½ mark)
- (ii) Calculate the value of the equilibrium constant, K_p for the reaction at 450 °C. (04 marks)

(b) State, giving reasons, how the concentration of sulphur trioxide at equilibrium would be affected if;

(i) pressure is increased. (1½ marks)

(ii) temperature is increased. (1½ marks)

(iii) an inert gas is added at constant pressure. (1½ marks)

15. (a) State the meaning of the term **bond energy**. (01 mark)

.....

.....

(b) The bond dissociation energies of fluorine, chlorine, bromine and iodine are 158, 242, 193 and 151 kJ mol⁻¹ respectively.

(i) State the trend in the bond dissociation energies of the elements. (1½ marks)

.....

.....

(ii) Explain your answer in (b)(i). (3½ marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Write an equation to show how cold dilute sodium hydroxide solution reacts with;

(i) fluorine. (1½ marks)

.....

(ii) chlorine. (1½ marks)

.....

.....

16. Propan-1-ol (boiling point 97°C) and water (boiling point 100°C) are miscible in all proportions. A mixture of the two liquids containing 72 % propan-1-ol boils at 88°C .

(a) Sketch a labelled boiling point–composition diagram for the mixture of propan-1-ol and water. (03 marks)

.....

.....

.....

.....

.....

.....

.....

.....

(b) Briefly explain;

(i) why propan-1-ol and water form a minimum boiling point mixture. (04 marks)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) what would happen when a mixture containing 30 % propan-1-ol is fractionally distilled. (02 marks)

.....

.....

.....

.....

.....

17. Tin and lead are elements in group(IV) of the Periodic Table. Describe the reactions of the elements with;

- (a) cold water. (03 marks)

.....

.....

.....

- (b) sodium hydroxide solution. (3½ marks)

.....

.....

.....

- (c) hot concentrated sulphuric acid. (2½ marks)

.....

.....

THE PERIODIC TABLE

| | | | | | | | | | | | | | | | | | |
|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
| 1 | 2 | | | | | | | | | | | 3 | 4 | 5 | 6 | 7 | 8 |
| 1.0 H 1 | | | | | | | | | | | | | | | | 1.0 H 1 | 4.0 He 2 |
| 6.9 Li 3 | 9.0 Be 4 | | | | | | | | | | | 10.8 B 5 | 12.0 C 6 | 14.0 N 7 | 16.0 O 8 | 19.0 F 9 | 20.2 Ne 10 |
| 23.0 Na 11 | 24.3 Mg 12 | | | | | | | | | | | 27.0 Al 13 | 28.1 Si 14 | 31.0 P 15 | 32.1 S 16 | 35.5 Cl 17 | 40.0 Ar 18 |
| 39.1 K 19 | 40.1 Ca 20 | 45.0 Sc 21 | 47.9 Ti 22 | 50.9 V 23 | 52.0 Cr 24 | 54.9 Mn 25 | 55.8 Fe 26 | 58.9 Co 27 | 58.7 Ni 28 | 63.5 Cu 29 | 65.7 Zn 30 | 69.7 Ga 31 | 72.6 Ge 32 | 74.9 As 33 | 79.0 Se 34 | 79.9 Br 35 | 83.8 Kr 36 |
| 85.5 Rb 37 | 87.6 Sr 38 | 88.9 Y 39 | 91.2 Zr 40 | 92.9 Nb 41 | 95.9 Mo 42 | 98.9 Tc 43 | 101 Ru 44 | 103 Rh 45 | 106 Pd 46 | 108 Ag 47 | 112 Cd 48 | 115 In 49 | 119 Sn 50 | 122 Sb 51 | 128 Te 52 | 127 I 53 | 131 Xe 54 |
| 133 Cs 55 | 137 Ba 56 | 139 La 57 | 178 Hf 72 | 181 Ta 73 | 184 W 74 | 186 Re 75 | 190 Os 76 | 192 Ir 77 | 195 Pt 78 | 197 Au 79 | 201 Hg 80 | 204 Tl 81 | 207 Pb 82 | 209 Bi 83 | 209 Po 84 | 210 At 85 | 222 Rn 86 |
| 223 Fr 87 | 226 Ra 88 | 227 Ac 89 | | | | | | | | | | | | | | | |
| | | | 139 La 57 | 140 Ce 58 | 141 Pr 59 | 144 Nd 60 | 147 Pm 61 | 150 Sm 62 | 152 Eu 63 | 157 Gd 64 | 159 Tb 65 | 162 Dy 66 | 165 Ho 67 | 167 Er 68 | 169 Tm 69 | 173 Yb 70 | 175 Lu 71 |
| | | | 227 Ac 89 | 232 Th 90 | 231 Pa 91 | 238 U 92 | 237 Np 93 | 244 Pu 94 | 243 Am 95 | 247 Cm 96 | 247 Bk 97 | 251 Cf 98 | 254 Es 99 | 257 Fm 100 | 256 Md 101 | 254 No 102 | 260 Lw 103 |