# P425/2 APPLIED MATHEMATICS

Paper 2

Nov. /Dec. 2022

3 hours



### UGANDA NATIONAL EXAMINATIONS BOARD

## Uganda Advanced Certificate of Education

#### APPLIED MATHEMATICS

Paper 2

. Francis different Sell Con-

3 hours

## INSTRUCTIONS TO CANDIDATES:

Answer all the eight questions in section A and any five questions from section B.

Any additional question(s) answered will not be marked.

All necessary working must be shown clearly.

Begin each answer on a fresh sheet of paper.

Graph paper is provided.

Silent, non-programmable scientific calculators and mathematical tables with a list of formulae may be used.

In numerical work, take acceleration due to gravity g, to be  $9.8~\mathrm{ms}^{-2}$ .

# SECTION A (40 MARKS)

Answer all the questions in this section.

- 1. A rough plane is inclined at an angle of sin <sup>-1</sup> (<sup>3</sup>/<sub>5</sub>) to the horizontal. A car of mass 1500 kg moves up the plane with a steady speed. The coefficient of friction between the car and the plane is <sup>1</sup>/<sub>4</sub>. Calculate the driving force exerted by the engine of the car.

  (05 marks)
- 2. The table below shows the number of people in each of the 100 cars that passed through a check point on a certain day.

Number of people	1	2	3	/ 4	5
Number of cars	41	33	18	6	2

Find:

(a) the mean number of people in each car.

(02 marks)

(b) the variance of the data.

(03 marks)


3. Use the trapezium rule with seven (7) ordinates to estimate

$$\int_{0}^{2} \frac{1}{3+4x^{2}} dx$$

correct to three decimal places.

(05 marks)

4. The diagram below shows a mass of 5 kg lying on a rough plane inclined at 30° to the horizontal. From the 5 kg mass, a light inextensible string passes up the line of greatest slope and over a smooth fixed pulley carrying a mass of 3 kg hanging freely. The tension in the string is just sufficient to prevent the 5 kg mass from sliding down the slope.



Determine the coefficient of friction between the 5 kg mass and the rough plane.

(05 marks)

- Two events A and B are such that  $P(A) = \frac{1}{2}$ ,  $P(B) = \frac{7}{12}$  and  $P(A' \cap B) = \frac{1}{2}$ . Find  $P(B' \cap A)$ .
- 6. The table below shows the amount of Euros equivalent to US dollars.

US dollars .	79	97	105
Euros	64	78	85

Using linear interpolation/extrapolation, calculate the amount of:

(a) dollars equivalent to 92 Euros.

(03 marks)

(b) Euros equivalent to 85 dollars.

(02 marks)

- 7. A river which is 50 m wide flows with a speed of 3 ms<sup>-1</sup>. A boat must be set to cross the river to a point directly opposite the starting point. The speed of the boat relative to the river is 4 ms<sup>-1</sup>. Determine the;
  - (a) velocity with which the boat sets off to cross the river. (03 marks)
  - (b) time taken by the boat to cross the river.

(02 marks)

- 8. Bag A contains 7 red balls and 4 blue balls. Bag B contains 5 red balls and 8 blue balls. A ball is randomly transferred from A to B. A ball is then randomly removed from bag B. Find the probability that:
  - (a) a red ball was removed from bag B.

(03 marks)

(b) the transferred ball is blue given that a red ball is removed from B.

(02 marks)

## SECTION B (60 MARKS)

Answer any five questions from this section.

All questions carry equal marks.

9. The table below shows the marks of eight students (A to H) in two tests.

Student	A	В	C	D	E	F	G	H
Test 1	67	73	50 ~	75	42 /	44	95	40
Test 2	82	70	64	78	52	64	80	64

- (a) (i) Plot a scatter diagram for the given data.
  - (ii) Draw a line of best fit on the scatter diagram.
  - (iii) Use your scatter diagram to obtain a mark in Test 2 which corresponds to a mark of 63 in Test 1. (06 marks)

- (b) (i) Calculate the rank correlation coefficient for the data.
  - (ii) Comment on your result in b (i).

(06 marks)

10. A particle of mass 4 kg is initially at rest at the point A (2, -2, 3) metres. The particle is acted upon by a force of  $F = \begin{pmatrix} 4t \\ t^2 \\ 5 \end{pmatrix}$  Newtons, where t is time in

seconds. Find the;

- (a) acceleration at time t. (02 marks)
- (b) speed of the particle after 3 seconds. (05 marks)
- (c) displacement of the particle after 3 seconds. (05 marks)
- 11. (a) Two numbers x and y are approximated by X and Y with errors  $\Delta x$  and  $\Delta y$  respectively. Show that the relative error in approximating  $\frac{x}{y}$  by  $\frac{X}{y}$  is  $\left|\frac{\Delta x}{X}\right| + \left|\frac{\Delta y}{Y}\right|$  (06 marks)
  - (b) The values in the expression  $T = \frac{673.16}{40.345}$  are rounded off correct to the given decimal places. Determine the interval within which the exact value of T can be expected to lie. (06 marks)
- 12. A discrete random variable X has a probability distribution given by

$$f(x) = \begin{cases} k x^2, & x = 1, 2, 3 \\ k(7-x)^2, & x = 4, 5, 6 \\ 0, & elsewhere \end{cases}$$

Find the:

- (a) (i) value of the constant k.
  - (ii) mean of X.
  - (iii) variance of X.

(09 marks)


- (b) (i) Determine the cumulative distribution function, F(x). (01 mark)
  - (ii) Sketch the graph of F(x).

(02 marks)

- Forces of magnitude 2N, 2N, 3N, 4N,  $2\sqrt{2}$  N and  $\sqrt{2}$  N act along sides  $\overrightarrow{AB}$ ,  $\overrightarrow{BC}$ ,  $\overrightarrow{DC}$ ,  $\overrightarrow{AD}$ ,  $\overrightarrow{AC}$  and  $\overrightarrow{DB}$  respectively. Where  $\overrightarrow{ABCD}$  is a square of side 2 m. Find the:
  - (a) resultant force. (09 marks)
  - (b) equation of the line of action of the resultant force. (03 marks)
- 14. The taxation system in a company is such that the tax T based on the amount A earned falls in three categories as shown in the table below:

AMOUNT EARNED  A (DOLLARS)	TAX PAID T (DOLLARS)
A < 2000	T=0
$2000 \le A \le 5000$	$T = \frac{2A}{100}$
A > 5000	$T = 60 + \frac{5}{100}(A-5000)$

Parts of a flow chart for computing the tax are given below.



- (a) (i) Arrange the given parts to form a complete logical flow chart.
  - (ii) State the purpose of the flow chart. (08 marks) (01 mark)
- (b) Perform a dry run for the flow chart and complete the table below:

  (03 marks)

. <b>N</b>	A	T
1	1500	_
2	3500	_
3	9000	

- The masses of packets of tea leaves are normally distributed with mean mass 600 g and standard deviation 20 g.
  - (a) If a packet of tea leaves is chosen at random, find the mass exceeded by 7% of the packets.

    (05 marks)
  - (b) If 1,000 packets of tea leaves were sold, find the number of packets that weighed less than 545 g. (07 marks)
  - 16. Two points A and B are on the same horizontal ground. A particle P is projected from point A towards point B with an initial speed of 49 ms<sup>-1</sup> at an angle of 30° to the horizontal. Two seconds later, another particle Q is projected from point B towards A with the same speed and at the same angle of elevation as P.

If the particles collide at a height h m above the ground, determine the:

(a) value of h. (06 marks)

(b) distance between the points A and B. (06 marks)