

When the Clinic Closes, Who Floods the ER?

A Demand–Supply Analysis of ED Overcrowding
DataFest 2026 Final Report

Team Analysis
May 2026

Executive Summary

Using 947,685 patients, 7.7M encounters, and the NYU ED Algorithm on 202K ED visits, we find that **55.6% of ED visits are broadly avoidable**. On the demand side, patients are pushed into the ED by temporal lockout (8.7× weekend surge), lack of PCP connection (2.3× ED rate), and lack of direction (social resource scarcity). On the supply side, the primary care workforce cannot absorb redirected demand—110 core PCPs operate at AAFP ceiling while specialists sit underutilized. Our solutions target both: expanding clinic hours, upgrading triage with Licensed Clinical Staff, distributing community resource guides with MyChart tutorials, and rebalancing specialist workloads toward PCP support.

Contents

1	Problem Statement: The ED Is Overwhelmed with Avoidable Visits	3
1.1	Supply Context: PCP Overload Limits Absorption Capacity	4
2	Demand Side: Why Patients Flood the ED	5
2.1	Demand Driver 1: Temporal Lockout—Evenings and Weekends	5
2.2	Demand Driver 2: Lack of PCP—Minor Issues Escalate to Emergencies	6
2.3	Demand Driver 3: Lack of Direction—Patients Don’t Know Where to Go	7
2.3.1	A. The Full Symptom Spectrum Arrives at ED	7
2.3.2	B. Repeat Visitors—Social Resource Scarcity	8
2.4	Demand Summary: Patient Flow from Vulnerability to ED	9
3	Supply Side: Medical Resource Maldistribution	10
3.1	PCP Overload	10
3.2	Specialist Workload Maldistribution	11
4	Solutions	12
4.1	Demand-Side Solutions	12
4.1.1	Solution D1: Dynamic Clinic Hours—Weekend/Extended Clinics	12
4.1.2	Solution D2: Upgraded ED Triage with Licensed Clinical Staff	13
4.1.3	Solution D3: Community Resource Booklet with MyChart Tutorial	13
4.2	Supply-Side Solution	15
4.2.1	Solution S1: Specialist Rebalancing to PCP Support	15
4.3	Solution Priority Matrix	15
5	Integrated Logic: How Solutions Address Root Causes	16

6	Validation & Predictive Support	16
6.1	Random Forest Model (AUC = 0.953)	16
6.2	Cluster Validation	17
7	Study Context & Methods	17
7.1	Limitations	18
8	Conclusion	18

1 Problem Statement: The ED Is Overwhelmed with Avoidable Visits

The NYU ED Algorithm—the gold standard for classifying ED visit necessity—reveals that **55.6% of 202,000 ED visits are broadly avoidable**:

NYU Category	Wt. %	Interpretation	Examples from Our Data
Non-Emergent	25.4%	Does not require treatment within 12h	Upper respiratory infection (URI), acute pharyngitis, nausea & vomiting, viral infection, otitis media
PCP Treatable	7.3%	Needs treatment but PCP could handle	Cystitis (UTI), dorsalgia (back pain), headache, dizziness
<i>Subtotal: Clinic-Appropriate</i>	32.7%	<i>Should never reach the ED</i>	
Preventable/Avoidable	22.9%	Would not have occurred with outpatient mgmt	Asthma exacerbation, diabetes complications, chronic abdominal pain, influenza with dehydration
<i>Total Broadly Avoidable</i>	55.6%		
Truly Emergent	18.6%	Requires immediate ED care	Sepsis, respiratory failure, myocardial infarction, pulmonary embolism, cerebrovascular accident (stroke)
Injury	21.1%	Trauma, fractures, open wounds	Fracture of femur, intracranial injury, burn & corrosion, poisoning
Mental Health/Substance	4.8%	Specialized psychiatric need	Alcohol use crisis, drug overdose, acute psychosis

Table 1: NYU ED Algorithm classification of 202K ED visits (probability-weighted) with representative diagnoses from our dataset. Over half are avoidable—not because patients are irrational, but because the system failed to catch them earlier.




Figure 1: NYU classification breakdown. Panel D: weekday vs. weekend acuity is virtually identical—weekend patients aren’t sicker, just locked out of clinics.

1.1 Supply Context: PCP Overload Limits Absorption Capacity

Resolving avoidable ED visits requires redirecting patients to primary care. But the primary care system is itself at capacity. After filtering out out-of-state physicians and those without valid Kansas ZIP codes, we retain **809 PCPs**:

PCP Engagement Tier	Count	%	Role
Not in system (0 encounters)	384	47.5%	Affiliated only, private practice
Name-only (1–10 enc)	230	28.4%	Minimal system interaction
Part-time (11–100 enc)	49	6.1%	Occasional referrals
Active (101–1K enc)	36	4.4%	Part-time system physicians
Full-time core (>1K enc)	110	13.6%	System-deployable workforce

Table 2: PCP engagement tiers after geographic filtering (809 PCPs in Kansas). Only 110 (13.6%) are full-time core, serving 177K patients at AAFP panel ceiling (~2,500/physician).




Figure 2: PCP distribution after geographic filtering (809 retained from 2,499). Log-transform reveals bimodal structure: a large “affiliated-only” group and a small “full-time core” group. Only 110 (13.6%) are system-deployable.

The 110 full-time core PCPs average 2,504 patients each—at the AAFP ceiling. **Any solution that redirects patients from ED must simultaneously expand PCP capacity**, or it simply shifts the bottleneck. This supply constraint motivates our supply-side solution (Section 4.2).

2 Demand Side: Why Patients Flood the ED

We identify three demand-side drivers that push patients into the ED:

2.1 Demand Driver 1: Temporal Lockout—Evenings and Weekends

Primary care clinics operate Monday–Friday, roughly 8am–5pm. The system is **closed 76% of the week** (128/168 hours). When it closes, ED becomes the only open door.




Figure 3: Weekend effect: ED proportion surges from 2.6% to 22.5% (8.7×). Weekend clinic volume = 4.3% of weekday. Patients aren’t sicker—just locked out.




Figure 4: Bivariate geographic map of night and weekend visit concentration. Red = high night rate, blue = high weekend rate, **purple = high both**. Purple areas—predominantly rural counties and underserved urban blocks—are the highest-priority targets for Solution D1 (dynamic clinic hours). The geographic pattern confirms that temporal lockout disproportionately affects areas with fewer healthcare alternatives.

Key Evidence: NYU acuity is identical on weekdays vs. weekends (clinic-appropriate: 33.4% vs. 30.8%). Weekend ED patients have the **same** needs as weekday patients—they simply have no alternative venue.

2.2 Demand Driver 2: Lack of PCP—Minor Issues Escalate to Emergencies

Patients without a PCP relationship have no medical home. Minor issues go unaddressed until they escalate into emergencies requiring ED visits.




Figure 5: No PCP relationship = 2.3× ED rate (6.23% vs 2.67%). A large fraction of patients have never been connected to any PCP.

The compound effect: No PCP + Weekend = 28.8% ED rate vs. PCP + Weekday = 2.0% (14× gap).




Figure 6: PCP connection × weekend interaction. The temporal lockout hits hardest for patients without a PCP.

The Medicare natural experiment at age 65 proves this mechanism: when patients gain insurance and PCP access, ED usage drops 35% overnight—with no education campaign, just access.




Figure 7: Medicare at 65: ED drops 35%, outpatient visits increase 7.6/person. All new visits go to clinics. Access itself is the intervention.

Metric	Age 55–64	Age 65–74	Change
ED utilization rate	5.18%	3.36%	↓ 35%
Total visits/person	23.7	31.3	↑ +7.6
Non-ED visits/person	23.0	30.6	↑ +7.6
PCP connection rate	51.6%	55.2%	↑ +3.6pp

Table 3: Medicare doesn’t reduce healthcare seeking—it redirects it from ED to outpatient.

2.3 Demand Driver 3: Lack of Direction—Patients Don’t Know Where to Go

Many patients default to the ED not because their condition is emergent, but because they **lack guidance** on appropriate care venues. This manifests in three ways:

2.3.1 A. The Full Symptom Spectrum Arrives at ED

The NYU Algorithm’s ICD-10 mapping confirms that a wide range of primary-care-manageable conditions present at ED:




Figure 8: Top ED diagnoses. Under-65 patients present with pain, URI, nausea—conditions readily manageable in primary care. Over-65 patients present with sepsis and respiratory failure—appropriate ED use. The under-65 pattern reflects misdirection to ED, not medical necessity.

2.3.2 B. Repeat Visitors—Social Resource Scarcity

SDOH (Social Determinants of Health) screening reveals that patients with social barriers—particularly material deprivation—cycle back to the ED repeatedly.




Figure 9: SDOH domain correlations. Transport/Food/Financial: $\rho > 0.98$ —one life condition, not three separate risks. Social isolation/Stress: $\rho = 0.92$. These collapse into a single syndrome of **social resource scarcity**.

Social Resource Scarcity = One Syndrome

The 11 SDOH domains collapse into 2 actionable dimensions:

- 1. **Material deprivation** (transport + food + financial): $\rho > 0.98$. Having one problem = having all three.
- 2. **Social isolation + stress**: $\rho = 0.92$. Strongest individual readmission predictor (RR = 2.04).

Patients with these barriers don't return to the ED by choice—they return because **unresolved social barriers** prevent them from accessing appropriate follow-up care.




Figure 10: 30-day readmission relative risk by SDOH domain. Social isolation is the strongest predictor (RR=2.04). Material deprivation domains cluster tightly.

2.4 Demand Summary: Patient Flow from Vulnerability to ED

The Sankey diagram below traces 349,440 patients through the complete pathway, showing how demand-side barriers channel patients toward ED use and readmission:

Patient Flow: Social Vulnerability → Primary Care → Emergency Use → Readmission
 Red pathways indicate high-risk flows; SDOH + No PCP group shows disproportionate ED use and readmission




Figure 11: Patient flow from SDOH risk → PCP connection → ED use → readmission. **SDOH + No PCP** patients (5.0%) show 41.4% ED rate and 15.1% readmission—the highest-risk pathway. Solutions D2 (triage) and D3 (resource booklet) target this population.

3 Supply Side: Medical Resource Maldistribution

The demand-side pressures are compounded by a supply-side failure: the physicians who *could* absorb demand are themselves overloaded or maldistributed.

3.1 PCP Overload

The 110 full-time core PCPs operate at AAFP panel ceiling (2,504 patients/physician). PCP overload creates a feedback loop:




Figure 12: Overloaded PCPs → patients with worse SDOH ($\rho = 0.31$) + more after-hours work. Overload feeds back into reduced access for the most vulnerable.




Figure 13: PCP overload deep-dive. Right panel: Family Medicine dominates the overloaded tier (>P90), confirming it serves as the system’s safety net.

3.2 Specialist Workload Maldistribution

After the same geographic filtering (Kansas ZIP only), we analyzed 19,425 non-PCP physicians. The pattern mirrors PCP but is even more extreme:

Specialist Engagement Tier	Count	%
Not in system (0 encounters)	17,375	89.4%
Affiliated only (≤10 enc)	1,393	7.2%
Part-time (11–100 enc)	317	1.6%
Active (101–1K enc)	119	0.6%
Full-time core (>1K enc)	221	1.1%
Total active (>0 enc)	2,050	10.6%

Table 4: Non-PCP specialist tiers after geographic filtering. Only 2,050 of 19,425 specialists are active in the system.

Among active specialists, workload distribution is highly skewed. In 29 of 30 specialties, **mean encounters** > **median encounters**—indicating a small number of physicians carry the majority of the load while many are underutilized:

Specialty	Active N	Median Enc	Mean Enc	Mean/Med Ratio
Anesthesiology	16	2	5,128	2,564×
Orthopedic Surgery	46	6	2,928	488×
Radiology	24	4	1,876	469×
General Surgery	28	12	2,341	195×
Cardiology	35	18	1,892	105×
Emergency Medicine	35	6,150	5,795	0.94×

Table 5: Selected specialties showing extreme mean/median ratios. Most specialists are nearly idle in the system while a few are overloaded. Emergency Medicine is the only specialty where workload is evenly distributed (ratio <1)—because ED physicians *must* work shifts regardless.




Figure 14: Specialist workload violin plot (log-scale). Red violins = severely skewed specialties (mean/median $>5\times$): most physicians are idle while few carry extreme loads. The shaded zone below 1,000 encounters represents **underutilized capacity available for PCP rebalancing**. Of 32 specialties analyzed, 22 are severely maldistributed—a structural reservoir of clinical workforce.

Supply-Side Insight

The imbalance is structural, not coincidental. Of 32 specialties with ≥ 5 active physicians, **22 are severely maldistributed** (mean/median ratio $>5\times$). Specialists in PCP-adjacent fields (Internal Medicine subspecialties, Geriatrics, Preventive Medicine) have the clinical training to support PCP workflows but sit at extremely low utilization within the system. Meanwhile, PCP and EM physicians operate at or above capacity. This represents a **redeployable workforce** that can relieve PCP overload without new hiring.

4 Solutions

Design Principle

All solutions are scoped to what a hospital system can implement **immediately** with existing resources. We do not recommend policy changes (Medicaid expansion), massive hiring, or expensive technology—only operational adjustments validated by our data.

4.1 Demand-Side Solutions

4.1.1 Solution D1: Dynamic Clinic Hours—Weekend/Extended Clinics

Problem addressed: Temporal lockout ($8.7\times$ weekend surge; 55.6% avoidable ED visits)

Implementation:

- Open 1–2 clinic rooms on Saturday mornings (9:00–12:00) and/or extend weekday hours to 7:00 PM
- Staff: 1 NP/PA + 1 front desk per session
- Prioritize: high-frequency ED users (system-identifiable), patients without PCP
- Dynamically adjust hours based on ED volume data—expand Saturday hours during flu season, reduce in summer

Evidence:

- Weekend ED proportion: 22.5% vs weekday 2.6% ($8.7\times$)
- NYU: 32.7% of ED visits are clinic-appropriate
- Projected: divert 10% of weekend ED volume $\approx$ 5,000 visits/year
- Cost: \$1,200/ED visit vs \$150/clinic visit $\rightarrow$ **\$5.25M annual savings**

4.1.2 Solution D2: Upgraded ED Triage with Licensed Clinical Staff

Problem addressed: Lack of direction—patients arrive at ED by default

Implementation:

- Upgrade the ED triage desk: in addition to triage nurses, deploy **Licensed Clinical Staff** (Physician Assistants, Nurse Practitioners) with prescribing authority and clinical judgment capability
- These Licensed Clinical Staff apply the **NYU ED Algorithm framework** at the point of entry to classify presenting complaints into:
 1. **Non-emergent / PCP-treatable:** Provide immediate basic treatment (OTC medications, wound care, prescriptions for antibiotics/analgesics) and **discharge with scheduled PCP follow-up**
 2. **Preventable/Avoidable:** Redirect to appropriate outpatient specialty clinic; schedule appointment before patient leaves
 3. **Repeat visitors:** Identify patients with ≥ 3 ED visits in 90 days; route to care coordinator for clinic-based follow-up plan
- This is NOT denial of care—it is **provision of more appropriate care at lower acuity and cost**

Evidence:

- NYU Algorithm: 32.7% of ED visits are non-emergent or PCP-treatable
- Weekday vs weekend acuity identical (33.4% vs 30.8% clinic-appropriate)—confirming that redirected patients are safe for outpatient management
- High-risk cycling group (58K patients): screened $27\times$ but never effectively redirected

4.1.3 Solution D3: Community Resource Booklet with MyChart Tutorial

Problem addressed: Social resource scarcity (SDOH $\rho > 0.98$) + digital access gap

Implementation:

- Create a concise community resource booklet (bilingual English/Spanish) placed at:
 - ED waiting room
 - Clinic discharge desk
 - Community health centers and pharmacies
- Content:
 1. **SDOH resource directory:** 211 hotline, food banks, community transport services, financial assistance programs—consolidated because $\rho > 0.98$ means one need = all needs
 2. **MyChart setup tutorial:** Step-by-step guide with screenshots for downloading, registering, and using three key features (view results, message provider, schedule appointments)

3. “Where should I go?” decision tree: Simple symptom-based flowchart guiding patients to appropriate venue (ED vs Urgent Care vs PCP vs self-care)

- The MyChart component is data-validated: activated users show **63.6% lower ED rate** and **3.6× lower readmission rate**

Evidence for MyChart effectiveness:




Figure 15: MyChart activated patients: 63.6% lower ED rate, 154% more total visits—care redirected from ED to planned outpatient. Portal provides after-hours access (messaging, scheduling).




Figure 16: Patients who actively declined MyChart: readmission 4.73% vs 1.33% for activated (3.6×). Digital access is a protective factor.

Metric	MyChart Activated	Not Activated / Declined
ED utilization rate	Low (baseline)	+63.6% higher
Total visits/person	154% higher	Baseline
30-day readmission	1.33%	4.73% (Declined)
After-hours access	Messaging + scheduling	None

Table 6: MyChart activation as a protective factor. The portal provides 24/7 channel access even when clinics are physically closed.

4.2 Supply-Side Solution

4.2.1 Solution S1: Specialist Rebalancing to PCP Support

Problem addressed: PCP overload (110 core PCPs at AAFP ceiling) + specialist maldistribution (29/30 specialties mean > median)

Mechanism: Our data shows that in most specialties, the *majority* of physicians are minimally active in the system (median near zero) while a small number carry extreme loads. This means there exists a pool of clinically trained physicians with available capacity.

Implementation:

- Identify specialists in **PCP-adjacent fields** (Internal Medicine subspecialties, Geriatrics, Preventive Medicine, Sports Medicine) whose system workload is below the specialty median
- Offer structured half-day PCP support sessions (1–2 per week) focused on:
 - Routine follow-ups for stable chronic conditions (hypertension, diabetes management)
 - Post-discharge check-ins for patients without established PCP
 - Saturday morning clinic staffing (see Solution D1)
- This leverages existing credentialed physicians without new hiring
- Compensation: pro-rata clinical time or RVU credit within existing contracts

Evidence:


- 29/30 specialties show mean > median (Table 5)—structural underutilization
- Anesthesiology: 16 active, median 2 encounters vs mean 5,128—most have capacity
- Orthopedics: 46 active, median 6 vs mean 2,928—same pattern
- Even reassigning 20–30 specialists to 1 half-day/week of PCP support adds the equivalent of **5–7 full-time PCPs**—a 5% increase in core PCP capacity
- Simultaneously reduces specialist overload for the few carrying extreme volumes

4.3 Solution Priority Matrix

Solution			Cost	Difficulty	Impact	Key Evidence		
D1:	Dynamic	clinic	Low	Low	High	8.7×	surge;	55.6%
	hours					avoidable		
D2:	Upgraded	triage	Med	Med	High	NYU	32.7%	redi-
	(LCS)					rectable		
D3:	Resource	booklet +	Low	Low	Med	63.6%	ED	reduction;
	MyChart					$\rho > 0.98$		
S1:	Specialist	→ PCP	Low	Med	High	29/30 specialties under-		
	support					utilized		

Table 7: Solution priority matrix. D1 and D3 are immediately deployable with minimal cost. D2 and S1 require modest operational coordination but yield high impact.

5 Integrated Logic: How Solutions Address Root Causes



Solutions target both sides: D1–D3 reduce demand on ED; S1 expands PCP capacity. Together they break the overload cycle.

6 Validation & Predictive Support

6.1 Random Forest Model (AUC = 0.953)

Our predictive model confirms that the factors we target are indeed the strongest drivers of ED use and readmission:




Figure 17: Top 15 features for 30-day readmission prediction. #1: Hospitalization count (0.547)—past system failure predicts future failure. SDOH screening count ranks #6—the system *identifies* high-risk patients but fails to act.




Figure 18: Effect sizes for all 18 significant predictors (Bonferroni-corrected $p < 0.003$). Hospitalization count and SDOH domains dominate.

6.2 Cluster Validation

Three distinct patient populations confirm our demand-supply framing:

Cluster	N	Avg Visits	ED Rate	SDOH Screens	Readmit
High-freq, Low-SDOH	70,826	44.5	2.9%	0.3	2.49%
High-risk Cycling	58,266	52.2	3.5%	27.2	4.85%
Low-freq Young	230,668	6.4	9.2%	0.1	0.34%

Table 8: Three patient clusters. The “High-risk Cycling” group (58K patients) is screened 27× more but readmits 14× more than the young group—the system sees them but fails to help. Solution D2 (triage redirect) and D3 (resource booklet) directly target this population.

7 Study Context & Methods

- **System:** Large health system in Kansas; 50/105 counties fully rural; many HPSAs
- **Data:** 947K patients, 7.7M encounters, 1.65M SDOH screenings, 1.53M diagnoses (2022–2025)
- **Geographic scope:** 94.89% of encounters in Kansas (ZIP 66000–67999 + KC Metro 64000–64199)
- **PCP filtering:** Removed out-of-state and no-ZIP physicians; 2,499 → 809 Kansas-based PCPs
- **External data:** NYU ED Algorithm (ICD-10 severity mapping), Kansas insurance pricing, AAFP panel standards

- **Statistical methods:** Mann-Whitney U, Kruskal-Wallis, Chi-squared, Spearman correlation, Bonferroni correction ($p < 0.003$), Cohen’s d , Cramér’s V
- **Models:** KMeans clustering ($k = 3$, Silhouette-optimized), Random Forest (5-fold CV, AUC = 0.953)
- **Policy context:** Kansas has NOT expanded Medicaid; ~39,000 adults in coverage gap; uninsured rate 8.3%

7.1 Limitations

1. **Observational design:** All analyses are correlational. Medicare at-65 approximates a natural experiment but is not a true RCT.
2. **MyChart selection bias:** Health-engaged patients may be more likely to activate portals. Effect size (63.6%) is large enough that bias alone is unlikely to explain it.
3. **PCP affiliation ambiguity:** Cannot precisely distinguish “retired” from “independent practice affiliated.” Geographic filtering mitigates but does not eliminate this.
4. **Age binning:** Data uses 5-year bins; Medicare threshold effect spans 62–67 age band.
5. **Specialist rebalancing feasibility:** Assumes willing participation; actual deployment depends on credentialing and departmental cooperation.

8 Conclusion

The Complete Narrative

The ED overcrowding in this Kansas health system is simultaneously a demand problem and a supply problem—but **neither is a patient behavior problem**.

On the demand side, three forces push patients into the ED:

1. **Temporal lockout**: Clinics close 76% of the week. Weekend ED surges $8.7\times$ with identical acuity—patients aren't sicker, just locked out.
2. **No PCP connection**: Without a medical home, minor issues escalate. No PCP = $2.3\times$ ED rate. Medicare proves: provide access $\rightarrow$ ED drops 35%.
3. **No direction**: Patients lack guidance on where to go. 32.7% of ED visits don't need ED-level care. Social resource scarcity ($\rho > 0.98$) traps patients in a cycle of return visits.

On the supply side, the workforce is maldistributed:

1. 110 core PCPs at AAFP ceiling with no absorption capacity
2. 29/30 specialist fields show extreme workload skew (most idle, few overloaded)
3. Family Medicine carries disproportionate burden as the safety net

Our solutions are operational, not aspirational:

1. **Dynamic clinic hours** break the temporal lockout
2. **Licensed Clinical Staff at triage** redirect 32.7% of ED visits to appropriate care
3. **Resource booklets + MyChart tutorials** address direction failure and social resource scarcity
4. **Specialist rebalancing** adds 5–7 FTE-equivalent PCPs without new hiring

Every contributing factor is independently actionable. The proof: when even one barrier is removed (Medicare at 65, MyChart activation, PCP connection), ED usage drops dramatically and automatically.

Technical Notes: Analysis conducted in Python (pandas, scipy, scikit-learn, matplotlib, plotly). All comparisons Bonferroni-corrected $p < 0.003$. Effect sizes: Cohen's d , Cramér's V . External classification: NYU ED Algorithm (47K ICD-10 codes mapped to severity categories). Geographic filtering applied to provider analyses (Kansas ZIP + KC Metro only).