2.2.1

For any natural numbers a, b, c, we have (a + b) + c = a + (b + c).

Proof. Induct on b by keeping a and c fixed. Consider the base case b = 0. In this case, LHS= (a + 0) + c = a + c and RHS= a + (0 + c) = a + c. Now suppose that (a + b) + c = a + (b + c). We need to show that (a + (b + +)) + c = a + ((b + +) + c):

LHS =
$$(a + (b + +)) + c = ((a + b) + +) + c = (a + b + c) + +$$
,
RHS = $a + ((b + +) + c) = a + ((b + c) + +) = (a + b + c) + +$.

Thus both sides are equal to each other, and we have closed the induction. \Box

2.2.2

Let a be a positive number. Then there exists exactly one natural number b such that b + + = a. (I'm assuming that it meant a is a positive natural number.)

Proof. Induct on a. Since 0 is not positive, we consider the base case a = 1. We have b++=b+1=0+1=1. Cancellation law tells us that b=0, which is unique. Now suppose that there exists exactly one natural number b_0 such that $b_0++=a$, we need to show that there exists exactly one natural number b such that b++=a++. By Cancellation law, we have $b=a=b_0++$. Since b is the successor of b_0 and b_0 is unique, b is also unique. Thus we have closed the induction.

2.2.3

(a)

 $a \ge a$.

Proof. There exists a natural number 0 such that a + 0 = a. Thus, $a \ge a$.

(b)

If $a \ge b$ and $b \ge c$, then $a \ge c$.

Proof. Since $a \ge b$, there exists a natural number m such that b+m=a. Since $b \ge c$, there exists a natural number n such that c+n=b. Then c+(n+m)=(c+n)+m=b+m=a. Therefore, $c \ge a$.

Thus, if $a \ge b$ and $b \ge c$, then $a \ge c$.

(c)

If $a \ge b$ and $b \ge a$, then a = b.

Proof. Since $a \ge b$, there exists a natural number m such that b+m=a. Since $b \ge a$, there exists a natural number n such that a+n=b. Then we have a+n=(b+m)+n=b+(m+n)=b. By Cancellation law, we have m+n=0 which leads to m=0, n=0. Thus, a=a+0=b.

Thus, if $a \ge b$ and $b \ge a$, then a = b.

(d)

 $a \ge b$ if and only if $a + c \ge b + c$.

Proof. First, we need to show that $a \ge b \Rightarrow a+c \ge b+c$. Since $a \ge b$, there exists a natural number n such that b+n=a. Then we have b+n+c=b+c+n=(b+c)+n=a+c. Thus, $a+c \ge b+c$. Then, we need to show that $a+c \ge b+c \Rightarrow a \ge b$. Since $a+c \ge b+c$, there should be a natural number n such that b+c+n=b+n+c=(b+n)+c=a+c. By Cancellation law, we have b+n=a. Thus, $a \ge b$.

Thus, if $a \ge b$ and $b \ge a$, then a = b.

(e)

a < b if and only if $a + + \leq b$.

Proof. First, we need to show that $a < b \Rightarrow a + + \leq b$. a < b means there exists a natural number n such that a + n = b, particularly, $a \neq b$. Then n must not be zero. So n is the predecessor of a natural number, denote it as m. Then we have a + n = a + (m + +) = (a + m) + + = (a + +) + m = b. Therefore, $a + + \leq b$. Then we need to show that $a + + \leq b \Rightarrow a < b$. There exists a natural number n such that (a + +) + n = b. (a + +) + n = (a + n) + + = a + (n + +) = b. Since n + + is the successor of n, n + + must not be equal to 0. If a = b, there will be $a + (n + +) = a \Rightarrow n + + = 0$, contradiction. Therefore, $a \neq b$.

Thus,
$$a < b$$
 if and only if $a + + \le b$.

(f)

a < b if and only if b = a + d for some positive number d.

Proof. First, we need to show that $a < b \Rightarrow b = a + d$ for some positive number d. There exists some natural number d such that a + d = b, $a \neq b$. By Cancellation law, d must not be zero. Therefore, d is positive. Then, we need to show that a + d = b for some positive $d \Rightarrow a < b$. We only need to prove $a \neq b$. If a = b, we have a + d = a = a + 0. By Cancellation law, d = 0 which contradicts to d is positive. Therefore, $a \neq b$.

Thus,
$$a < b$$
 if and only if $b = a + d$ for some positive number d .

2.2.4

Justify the three statements marked in the proof of Proposition 2.2.13.

(a)

 $0 \le b$ for all b.

Proof. By definition of addition, we have 0+b=b. Thus, $0 \le b$.

(b)

If a > b, then a + + > b.

Proof. By Proposition 2.2.12.e, we have $a > b \Rightarrow a \ge b + +$. And by Proposition 2.2.12.d, $a + 1 \ge (b + +) + 1$ that is equivalent to $a + + \ge b + 2$. Since 2 is positive, by Proposition 2.2.12.f, we have a + + > b.

(c)

If a = b, then a + + > b.

Proof. We know from Proposition 2.2.12.a that $a \ge a$, so $a \ge a = b$. And again by Proposition 2.2.12.d, we have $a++=a+1 \ge b+1$. Since 1 is positive, by Proposition 2.2.12.f, a++>b.

2.2.5

Proposition 2.2.14 (Strong principle of induction). Let m_0 be a natural number, and let P(m) be a property pertaining to an arbitrary natural number m. Suppose that for each $m \geq m_0$, we have the following implication: if P(m') is true for all natural numbers $m_0 \leq m' < m$, then P(m) is also true. Then we can conclude that P(m) is true for all natural numbers $m \geq m_0$.

Proof. Let Q(n) be the property that P(m) is true for all $m_0 \leq m < n$. Induct on n. Consider the base case n = 0. This is vacuously true. In fact, Q(n) is vacuously true for all $n \leq m_0$. So we can assume $n > m_0$ to see if the implication stands. Suppose Q(n) is true, that is, P(m) is true for all $m_0 \leq m < n$. We want to show that Q(n+1) is also true. As stated in Proposition 2.2.14, if Q(n) is true, then P(n) is also true. So P(m) is true for all $m_0 \leq m \leq n$. Hence, P(m) is true for all $m_0 \leq m < n + 1$. $(m \leq n \Leftrightarrow m < n + 1 \text{ can be shown using prop 2.2.12.})$ Thus, Q(n+1) is true. This closes the induction.

2.2.6

Let n be a natural number, and let P(m) be a property pertaining to the natural numbers such that whenever P(m++) is true, then P(m) is true. Suppose that P(n) is also true. Prove that P(m) is true for all natural numbers $m \leq n$. (Principle of backwards induction.)

Proof. Apply induction to n. For the base case n=0, suppose P(0) is true. In this case, m can only be 0 ($m+k=0 \Rightarrow m=0, k=0$). Since P(0) is true, the base case is proved. Next, suppose if P(n) is true then P(m) is true for all natural numbers $m \leq n$. We want to show that if P(n++) is true then P(m) is true for all natural numbers $m \leq n++$. $m \leq n$ means there exists a natural number a such that m+a=n++. a is either 0 or a positive number. If a is 0, m=n++. If a is positive, m < n++ (by prop 2.2.12.f), this is equivalent to $m \leq n$ (can be shown using prop 2.2.12.). For m=n++, P(m) is true because of the assumption. For each $m \leq n$, P(m) is also true by induction hypothesis. Therefore, P(m) is true for all natural numbers $m \leq n++$. And we have closed the induction.

In the above proofs, n + + and n + 1 got mixed up because n + + = n + 1 has been illustrated on Page 26 (and the +1 version is a little easier). But ++ is a more desirable expression since it stands for the successor in a general way.