

Cambridge International AS & A Level

CANDIDATE
NAME


--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/23

Paper 2 Further Pure Mathematics 2

October/November 2021

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

- 1 Find the Maclaurin's series for $e^x \tan x$ from first principles up to and including the term in x^2 . [5]

[illegible]

(b) Find the value of $\frac{d^2y}{dx^2}$ at the point $(-1, 1)$. [5]

[illegible]

(b) Use a similar method to find, in terms of N , a lower bound for $\sum_{r=1}^N \frac{\ln r}{r^2}$. [3]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

5 Find the particular solution of the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 4\cos x,$$

given that, when $x = 0$, $y = -4$ and $\frac{dy}{dx} = 3$. [11]

[illegible]

(b) Hence obtain the roots of the equation

$$x^5 - 10x^4 + 40x^2 - 32 = 0$$

in the form $\operatorname{cosec}(q\pi)$, where q is rational. [4]

[illegible]

- 7 (a) Show that an appropriate integrating factor for

$$\sqrt{x^2-1} \frac{dy}{dx} + y = x^2 - x\sqrt{x^2-1}$$

is $x + \sqrt{x^2 - 1}$.

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

for which $y = 1$ when $x = \frac{5}{4}$. Give your answer in the form $y = f(x)$. [7]

This image shows a full page of a document template designed for handwriting practice or general writing. It consists of approximately 28 evenly spaced, horizontal dotted lines extending across the entire width of the page. The background is plain white, and there are no margins, headers, footers, or other markings present.

[illegible]

$$x = 2 \cosh 2t + 3t, \quad y = \frac{3}{2} \cosh 2t - 4t, \quad \text{for } -\frac{1}{2} \leq t \leq \frac{1}{2}.$$

(b) (i) Show that $A = 10\pi \int_{-\frac{1}{2}}^{\frac{1}{2}} (2 \cosh 2t + 3t) \cosh 2t \, dt$. [4]

[illegible]

(ii) Hence find A in terms of π and e .

[7]

[illegible]

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.