

Cambridge International AS & A Level

CANDIDATE
NAME


--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/21

Paper 2 Further Pure Mathematics 2

October/November 2023

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

BLANK PAGE

- 2 Find the roots of the equation $(z + 5i)^3 = 4 + 4\sqrt{3}i$, giving your answers in the form $r \cos \theta + i(r \sin \theta - 5)$, where $r > 0$ and $0 < \theta < 2\pi$. [5]

[illegible]

- 3 Find the first three terms in the Maclaurin's series for $\tanh^{-1}\left(\frac{1}{2}e^x\right)$ in the form $\frac{1}{2}\ln a + bx + cx^2$, giving the exact values of the constants a , b and c . [6]

[illegible]

[Turn over

[illegible]

[illegible]

- (c)** Find the particular solution of the differential equation

$$\frac{dy}{dx} + y \tanh x = \sinh^2 2x,$$

given that $y = 4$ when $x = 0$. Give your answer in the form $y = f(x)$. [7]

[illegible]

$$\mathbf{A} = \begin{pmatrix} -6 & 2 & 13 \\ 0 & -2 & 5 \\ 0 & 0 & 8 \end{pmatrix}.$$

- (a) Find a matrix $\mathbf{P}$ and a diagonal matrix $\mathbf{D}$ such that $\mathbf{A}^{-1} = \mathbf{PDP}^{-1}$. [7]


This image shows a full page of a handwriting practice worksheet. It consists of approximately 20 horizontal dotted lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

[illegible]

- 8 (a)** State the sum of the series $1+z+z^2+\dots+z^{n-1}$, for $z \neq 1$. [1]

(b) By letting $z = \cos \theta + i \sin \theta$, where $\cos \theta \neq 1$, show that

$$1 + \cos \theta + \cos 2\theta + \dots + \cos (n-1)\theta = \frac{1}{2} \left(1 - \cos n\theta + \frac{\sin n\theta \sin \theta}{1 - \cos \theta} \right). \quad [7]$$



The diagram shows the curve with equation $y = \cos x$ for $0 \leq x \leq 1$, together with a set of n rectangles of width $\frac{1}{n}$.

(c) By considering the sum of the areas of these rectangles, show that

$$\int_0^1 \cos x \, dx < \frac{1}{2n} \left(1 - \cos 1 + \frac{\sin 1 \sin \frac{1}{n}}{1 - \cos \frac{1}{n}} \right). \quad [4]$$

.....

.....

.....

.....

.....

.....

(d) Use a similar method to find, in terms of n , a lower bound for $\int_0^1 \cos x \, dx$. [3]

.....

.....

.....

.....

.....

.....

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.