References

- [1] G. Si, T. Wu, Q. Ouyang, Y. Tu. *A Pathway-based Mean-field Model for Escherichia coli Chemotaxis*. Phys Rev Lett. **109(4)** (2012): doi: 10.1103/PhysRevLett.109.048101
- [2] J. E. Keymer, R. G. Endres, M. Skoge, Y. Meir, N. S. Wingreen. Chemosensing in Escherichia coli: Two regimes of two-state receptors. PNAS 103(6) (2006) 1786–1791: doi: 10.1073pnas.0507438103
- [3] H. C. Berg, D. A. Brown. *Chemotaxis in Escherichia coli analysed by three-dimensional tracking*.

 Nature **239** (1972) 500 504
- [4] N. Vladimirov, L. Løvdok, D. Lebiedz, V. Sourjik. Dependence of Bacterial Chemotaxis on Gradient Shape and Adaptation Rate. PLoS Computational Biology 4(12) (2008): e1000242. doi:10.1371/journal.pcbi.1000242
- [5] G. Micali, R. Colin, V. Sourjik, R. G. Endres. *Drift and Behavior of E. coli Cells*. Biophysical Journal 113 (2017) 2321–2325: https://doi.org/10.1016/j.bpj.2017.09.031
- [6] R. Karmakar, R. V. S. U. Bhaskar, R. E. Jesudasan, M. S. Tirumkudulu, K. V. Venkatesh. Enhancement of Swimming Speed Leads to a More-Efficient Chemotactic Response to Repellent. Appl Environ Microbiol 82 (2015) 1205–1214: doi:10.1128/AEM.03397-15
- [7] J. Tatikos, H. Stark, V. Zaburdaev. How the Motility Pattern of Bacteria Affects Their Dispersal and Chemotaxis. PLoS ONE 8(12) (2013): e81936. doi:10.1371/journal.pone.0081936
- [8] S. Neumann, N. Vladimirov, A. K. Krembel, N. S. Wingreen, V. Sourjik. Imprecision of adaptation in Escherichia coli chemotaxis. PloS One 9 (2014):e84904
- [9] Y. V. Kalinin, L. Jiang, Y. Tu, M. Wu. Logarithmic Sensing in Escherichia coli Bacterial Chemotaxis. Biophysical Journal **96** (2009) 2439–2448: doi: 10.1016/j.bpj.2008.10.027

- [10] J. D. Murray. (2000) Mathematical Biology: I. An Introduction, Third Edition, Springer, New York City
- [11] J. Saragosti, P. Silberzan, A.Buguin. *Modeling E. coli Tumbles by Rotational Diffusion*. *Implications for Chemotaxis*. PLoS ONE **7(4)** (2012): e35412. doi:10.1371/journal.pone.0035412
- [12] W. W. Tso, J. Adler. Negative Chemotaxis in Escherichia coli. Journal of Bacteriology 118(2) (1973) 560-576
- [13] M. Descerno. One-dimensional difiusion on a finite region. Department of Chemistry and Biochemistry, UCLA (2004)
- [14] L. Jiang, Q. Ouyang, Y. Tu. Quantitative Modeling of Escherichia coli Chemotactic Motion in Environments Varying in Space and Time. PLoS Comput Biol 6(4) (2010): e1000735. doi:10.1371/journal.pcbi.1000735
- [15] Y. Yang, A. M. Pollard, C. Höfler, G. Poschet, M. Wirtz, R.Hell, V. Sourjik. *Relation between chemotaxis and consumption of amino acids in bacteria*. Molecular Microbiology **96(6)** (2015) 1272–1282: doi:10.1111/mmi.13006
- [16] K. Sona, F. Menolascinac, R. Stocker. Speed-dependent chemotactic precision in marine bacteria. PNAS 113(31) (2016) 8624–8629
- [17] G. Micali, R. Colin, V. Sourjik, R. G. Endres. Supplementary Information: Drift and behavior of E. coli cells. Biophysical Journal 113 (2017): doi: 10.1529/biophysj.106.090944
- [18] G. Si, T. Wu, Q. Ouyang, Y. Tu. Supplemental Material for "A Pathway-based Mean-field Model for Escherichia coli Chemotaxis". Phys Rev Lett. 109(4) (2012): doi: 10.1103/PhysRevLett.109.048101
- [19] H. C. Berg, D. A. Brown. Temporal Stimulation of Chemotaxis in Escherichia Coli. Proc. Nat. Acad. Sci. 71(4) (1974) 1786–1791

[20] R. G. Endres, O. Oleksiuk, C. H. Hansen, Y. Meir, V. Sourjik, N. S. Wingreen. *Variable sizes of Escherichia coli chemoreceptor signaling teams*. Molecular Systems Biology **4(211)** (2008): doi:10.1038/msb.2008.49