



Literature review

PhD candidate: François Leroy

Programme: Environmental Earth Sciences

Department: Spatial Sciences

*"Mapping biodiversity changes across
spatio-temporal scales"*

Advisor: doc. Ing. Petra Šímová, Ph.D.

Consultant: Mgr. Petr Keil, PhD

Beginning of study: October 2020

Contents

| | |
|--------------------------------|-----------|
| Outline | 1 |
| Dashboard | 2 |
| 1 Introduction | 3 |
| 2 Materials and Methods | 5 |
| Supplementary materials | 20 |
| References | 30 |

Outline

Literature review about the link between biodiversity facets trends and spatial/temporal scales.

The idea is to take every paper that talk about biodiversity trends (so far using just the species richness seems already a lot of paper) and to list **1)** which biodiversity metric they use **2)** which taxon/taxa they use, **3)** the spatial scale, **4)** the temporal scale and **5)** what is the dynamic (does the biodiversity metric increase/decrease/doesn't change over time/unclear).

Make a table of all these papers and `group_by(taxa) %>% order_by(spatial_scale | temporal_scale)`. Then see if for each taxa we can find a trend (a bit like in Chase *et al.* 2019 Oikos paper | Jarzyna *et al.* 2015 but here I am not making the analysis, just taking the analysis from papers). Best example found so far: [Hill & Hamer 2004](#)

I am using the “Advanced Research” tab of Web of Science which allows me skim through the entire literature using a convenient syntax. For instance:

```
AB = ((biodiversity OR species richness OR diversity) AND  
(temporal trend* OR dynamic*) AND  
(bird* OR avia*))
```

And

```
AB = ((biodiversity change index) AND (bird* OR avia*) AND trend*)
```

And

```
AB = ((species richness) AND (bird* OR avia*) AND trend*)
```

And

```
ALL=(birds AND species richness AND temporal trend)
```

Dashboard

Reference paper

- 05/07/2021: research was made with the literature review filter for the first query (stopped at #13) and created the second query (stopped at #2)
- 07/07/2021: questions to Petr: **1)** can the geometric mean of relative abundance + the weighted goodness of fit be used as biodiversity trend index, **2)** can the Farmland Bird Indicator (FBI) be used as biodiversity trend (for me it is more biodiversity health, Chiron et al 2013) **3)** what about the Red List Index trend? **4)** what about Multispecies population indexes?
- 08/07/2021: stopped at the article 41 for research #2.
- 12/08/2021: stopped at article 4 for research #4
- 13/08/2021: stopped at article 8 for research #4
- 17/08/2021: stopped at article 15 for research #4
- 18/08/2021: stopped at article 30 for research #4
- 19/08/2021: stopped at article 46 for research #4
- 20/08/2021: stopped at article 64 for research #4
- 01/09/2021: verifying spatial scales → stopped at Dittrich 2019

1. Introduction

Human life quality is intrinsically linked to ecosystems state that he is living in. Indeed, ecosystems services extend in a large spectrum of mechanisms including nutrient cycle, food production, or climate and water cycle regulation (Pereira et al.). Some of those ecosystem functions are managed by bird biodiversity such as seed dispersal, controls pests or pollinate plant. Unfortunately, anthropogenic stressors like habitat loss, over exploitation, pollution or introduction of invasive species could lead biodiversity to its sixth mass extinction (Barnosky et al.).

Biodiversity erosion is now known from everyone and political decisions has been stated in order to limit it (e.g. The Convention on Biological Diversity, 2010, 2002). However, these objectives have been so far not reached due mainly to our confusion and misunderstanding about biodiversity dynamic and how to determine it.

As a matter of fact, studying biodiversity can be confusing, especially because several choices must be done. Firstly, the level at which you are looking at the biodiversity must be chosen (e.g. species, functional, phylogenetic diversity). Secondly, one must decide which metric is the most appropriate for his study. There are many facets of biodiversity that can be measured by different metrics depending on the objective of your study. Measures of static biodiversity are commonly used such as species richness or α diversity (i.e. number of species, Whittaker, b), the Shannon index (Shannon), the Simpson index (Simpson) or the Hill number (Hill). The later three biodiversity indexes take into account the relative abundances of the species and can be considered as the *quality* of the biodiversity. On an other hand, the spatial and temporal β diversity will measure the species turnover and can be measured thanks to Whittaker's (Whittaker, a), Sørensen's (Sørensen) or Jaccard's (Jaccard) dissimilarity indexes (e.g. Keil et al.).

However, overall biodiversity (i.e. taking into account species of every taxa) may not be relevant for one's case study. Thus, several multi-species indicators have also been created, taking into account the abundances of indicator species giving information on the ecosystem health. The most known ones are the Red List Index (Butchart et al., b,a,c) or the Biodiversity Change Index (Normander et al.).

Using all the metrics cited above, we now know that the loss of global biodiversity is unprecedented. However, current scientific literature has also shown that temporal trends in local changes of biodiversity

can be opposite to trends at larger scales (*e.g.* [Chase et al.](#)). Thus, current changes in biodiversity is far more complex than a simple global decrease: most of the ecosystems undergo alterations of their communities with changes in species composition ([Blowes et al.](#); [Dornelas et al.](#)). Wonders persist about how the trend of these different metrics of biodiversity are link to the spatial and temporal scales used when measured.

In order to investigate this link between spatial scales and biodiversity metrics, birds is a relevant taxon. Thanks to the many ornithological monitoring and surveys, we now have a large number of long, high-quality time series on bird populations ([Bejček and Stastný](#)). Birds are easy to observe, easy to identify and thus many volunteers are motivated to conduct standardized sampling. Given their ability to change quickly of locations, their presence is also a good indicator for ecosystem health and thus several standardized metrics have been created to assess their populations. For instance, the geometric mean of relative abundances or the goodness-of-fit statistic ([Studený et al.](#)) are some of the baseline. Other multi-species indicators have also been created specifically for birds, such as the Farmland Bird Indicator ([Gregory et al., a](#)), the Forest Bird Indicator ([Gregory et al., b](#)) or the Wild Bird Indicator ([Gregory and Strien](#)).

Here, I propose to review articles assessing the temporal trends of different avian biodiversity metrics and to look at which spatial scales these studies have been done. Summarizing the trends of these qualitative and/or quantitative avian biodiversity indexes along with their spatial and temporal scales will help to see more clearly how the trends of biodiversity are linked to spatio-temporal scales. It is also important to demonstrate that the information about the sampling plan (*i.e.* spatial scale, time span, temporal scales etc) is not systematically indicated in the scientific literature and can bring confusion to the analysis and comparisons of their trends. I believe that this review can help to have a better overview of the current knowledge on the trend of biodiversity metrics of bird populations.

2. Materials and Methods

For this review, articles of interest were the ones assessing temporal trends of the most common indicators (*i.e.* metrics) of avian biodiversity and specifying spatial and temporal scales. For this, I used the “*advanced search*” tool of the ISI Web of Science Core collection database with these four following queries:

1. AB = ((biodiversity OR species richness OR diversity) AND (temporal trend* OR dynamic*) AND (bird* OR avia*)) which resulted in 1346 references.
2. AB = ((biodiversity change index) AND (bird* OR avia*) AND trend*) which resulted in 60 references.
3. AB = ((species richness) AND (bird* OR avia*) AND trend*) which resulted in 313 references.
4. ALL=(birds AND species richness AND temporal trend) which resulted in 88 references.

For each query, the title and abstract of the articles were reviewed. When the temporal trend was explicitly specified (either visually or literally), the material and method part was read in order to collect the *spatial grain* of the trend (*i.e.* the area at which the trend is assessed), its *temporal grain* (*i.e.* the time span at which data have been gathered on the field), the *spatial extent* (*i.e.* the entire area at which the study applies), the *temporal extent* and the *beginning and ending years* of the study as well as the *general trend* of the metric (Tab. 2.1).

Concerning the trend assessment, some papers contained the *p-value* or directly specified the significant trend of the metric. However, a portion of papers gives only visual representations of the trend. For those, the standard error was used when displayed. For the very few only giving the trend, **the rule of thumb was applied**. Information can be found in the column *Note* of the Tab. 2.2 of the supplementary material. Moreover, the final trend retained (*i.e.* either *Increase*, *Stable* or *Decrease*) doesn't reflect all the fluctuations of the metric through time but rather the difference between the starting and ending points.

Moreover, [Pilotto et al.](#) conducted a meta-analysis in which they computed and summarized the trend of four biodiversity metrics (namely, species richness, species diversity, abundance and temporal turnover). Some of them were concerning bird communities. For those latter, I used their code and data on the

[github repository](#) of their paper in order to compute the trends of these four metrics for the bird datasets.

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness,

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|----------------------|----------------------------|----------------------|----------------|-----------------------------------|-----------------|-----------|---------|----------|
| Barnagaud et al. | SR | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Increase |
| Barnagaud et al. | Abundance | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Decrease |
| Barnagaud et al. | Evenness | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Increase |
| Barnagaud et al. | Functional richness | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Increase |
| Barnagaud et al. | Functional dispersion | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Stable |
| Barnagaud et al. | Functional evenness | 0.5 Km ² | 1 year | 9,834 millions km ² | 41 years | 1970-2011 | USA | Increase |
| Roels et al. | SR | 0.04 Km ² | 1 year | 0.04 Km ² | 5 years | NA | Panama | Increase |
| Roels et al. | Bird activity | 0.04 Km ² | 1 year | 0.04 Km ² | 5 years | NA | Panama | Increase |
| Wretenberg et al. | SR | 0.03 Km ² | 1 year | 1800 km2 | 11 years | 1994-2004 | Sweden | Decrease |
| Ram et al. | SR | 1.6 Km ² | 1 year | 350 000 Km2 | 18 years | 1998-2015 | Sweden | Increase |
| Ram et al. | SR | 1.6 Km ² | 1 year | 350 000 Km2 | 18 years | 1998-2015 | Sweden | Stable |
| Ram et al. | SR | 1.6 Km ² | 1 year | 350 000 Km2 | 18 years | 1998-2015 | Sweden | Increase |
| Ram et al. | Multi-species indicator | 1.6 Km ² | 1 year | 350 000 Km2 | 18 years | 1998-2015 | Sweden | Increase |
| Ram et al. | Multi-species indicator | 1.6 Km ² | 1 year | 350 000 Km2 | 18 years | 1998-2015 | Sweden | Increase |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

∞

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|------------------------|------------------------|------------------------|----------------|----------------------------|-----------------|-----------|----------|----------|
| Harrison et al. (b) | Geometric mean | 10 000 Km ² | 0.5 year | 242 495 km ² | 20 years | 1994-2013 | UK | Increase |
| Harrison et al. (b) | GoF ($\lambda = -1$) | 10 000 Km ² | 0.5 year | 242 495 km ² | 20 years | 1994-2013 | UK | Stable |
| Harrison et al. (b) | GoF ($\lambda = -2$) | 10 000 Km ² | 0.5 year | 242 495 km ² | 20 years | 1994-2013 | UK | Stable |
| Doxa et al. | FBI | 4 Km ² | 1 year | 643 801 km ² | 8 years | 2001-2008 | France | Increase |
| Doxa et al. | FBI | 4 Km ² | 1 year | 643 801 km ² | 8 years | 2001-2008 | France | Stable |
| Doxa et al. | FBI | 4 Km ² | 1 year | 643 801 km ² | 8 years | 2001-2008 | France | Stable |
| Arnold et al. | SR | 0.02 Km ² | 1 year | 1000 km ² | 100 years | NA | Trinidad | Stable |
| Arnold et al. | Shannon | 0.02 Km ² | 1 year | 1000 km ² | 100 years | NA | Trinidad | Stable |
| Arnold et al. | Simpson | 0.02 Km ² | 1 year | 1000 km ² | 100 years | NA | Trinidad | Stable |
| Xu et al. | SR | 6.56 Km ² | 1 year | 6.56 Km ² | 12 years | 2002-2013 | China | Decrease |
| Jiguet et al. | GBI | 4 Km ² | 1 year | 643 801 km ² | 22 years | 1989-2009 | France | Increase |
| Jiguet et al. | WBI | 4 Km ² | 1 year | 643 801 km ² | 22 years | 1989-2009 | France | Increase |
| Jiguet et al. | UBI | 4 Km ² | 1 year | 643 801 km ² | 22 years | 1989-2009 | France | Increase |
| Jiguet et al. | FBI | 4 Km ² | 1 year | 643 801 km ² | 22 years | 1989-2009 | France | Increase |
| Jiguet et al. | EU bird directive | 4 Km ² | 1 year | 643 801 km ² | 22 years | 1989-2009 | France | Increase |
| Jiguet et al. | RLI (Red list Index) | NA | 1 year | 10 180 000 km ² | 22 years | 1989-2009 | France | Decrease |
| Keten | SR | 1.7 Km ² | 1 year | 1.7 Km ² | 11 years | 2006-2016 | Turkey | Stable |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|---------------------|------------------------|-----------------------|----------------|---------------------------------|-----------------|-----------|-------------|----------|
| Davey et al. | Simpson | 1 km ² | 1 year | 242 495 km ² | 13 years | 1994-2006 | UK | Increase |
| Davey et al. | SR | 1 km ² | 1 year | 242 495 km ² | 13 years | 1994-2006 | UK | Increase |
| Davey et al. | Evenness | 1 km ² | 1 year | 242 495 km ² | 13 years | 1994-2006 | UK | Increase |
| Christian et al. | SR | 15.4 Km ² | NA | 15.4 km ² | 209 ans | 1898-2006 | France | Increase |
| Dittrich et al. | SR | 0.053 km ² | 0.33 year | 53 Km ² | 3 years | 2010-2012 | Spain | Increase |
| Dittrich et al. | SR | 0.053 km ² | 1 year | 53 Km ² | 3 years | 2010-2012 | Spain | Increase |
| Dittrich et al. | SR | 0.083 km ² | 0.33 year | 53 Km ² | 3 years | 2012-2014 | UK | Stable |
| Dittrich et al. | SR | 0.083 km ² | 1 year | 53 Km ² | 3 years | 2012-2014 | UK | Stable |
| Sirami and Monadjem | SR | 0.38 km ² | 1 year | 430 Km ² | 21 years | 1998-2018 | Swaziland | Decrease |
| García-Navas et al. | spatial beta-diversity | 267 Km ² | 1 year | 267 Km ² | 20 years | 1999-2018 | Switzerland | Decrease |
| McGeoch et al. | RLI (Red list Index) | NA | 1 year | 148,939,063.133 km ² | 11 years | 1998-2008 | Worldwide | Decrease |
| Ellis et al. | SR | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Stable |
| Ellis et al. | SR | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Stable |
| Ellis et al. | SR | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |
| Ellis et al. | SR | 0.48 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Stable |
| Ellis et al. | Shannon | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |
| Ellis et al. | Shannon | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Decrease |
| Ellis et al. | Shannon | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |
| Ellis et al. | Shannon | 0.48 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|------------------|-------------------------|--------------------------|----------------|-------------------------|-----------------|-----------|-------------------------|----------|
| Ellis et al. | Simpson | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |
| Ellis et al. | Simpson | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Decrease |
| Ellis et al. | Simpson | 0.16 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Increase |
| Ellis et al. | Simpson | 0.48 Km ² | 1 year | NA | 21 years | 1994-2014 | Oregon, USA | Decrease |
| Sicarella et al. | Occurence (%) | 17 370.3 Km ² | 1 year | 23 844 km ² | 22 years | 1992-2013 | Lombardy, Italy | Stable |
| Sicarella et al. | Occurence (%) | 1 403.9 Km ² | 1 year | 23 844 km ² | 22 years | 1992-2013 | Lombardy, Italy | Stable |
| Sicarella et al. | Occurence (%) | 6 461.9 Km ² | 1 year | 23 844 km ² | 22 years | 1992-2013 | Lombardy, Italy | Increase |
| Nally | SR | 0.49 Km ² | 1 day | 10 Km ² | 3 years | 1994-1996 | Australia | Increase |
| Latta et al. | SR | 0.000942 Km ² | 2 years | NA | 14 years | 1994-2007 | Ecuador | Decrease |
| Latta et al. | SR | 0.000942 Km ² | 2 years | NA | 14 years | 1994-2007 | Ecuador | Decrease |
| Scarton | SR | 0.55 Km ² | 2 years | 0.55 Km ² | 25 years | 1990-2014 | Lagoon of Venice, Italy | Increase |
| Scarton | Shannon | 0.55 Km ² | 2 years | 0.55 Km ² | 25 years | 1990-2014 | Lagoon of Venice, Italy | Increase |
| Scarton | Temporal beta-diversity | 0.55 Km ² | 2 years | 0.55 Km ² | 25 years | 1990-2014 | Lagoon of Venice, Italy | Increase |
| Scarton | Temporal beta-diversity | 0.55 Km ² | 2 years | 0.55 Km ² | 25 years | 1990-2014 | Lagoon of Venice, Italy | Increase |
| Chiron et al. | FBI | 4 Km ² | 1 year | 643 801 Km ² | 14 years | 2007-2020 | France | Decrease |
| Chiron et al. | FBI | 4 Km ² | 1 year | 643 801 Km ² | 14 years | 2007-2020 | France | Decrease |
| Chiron et al. | FBI | 4 Km ² | 1 year | 643 801 Km ² | 14 years | 2007-2020 | France | Decrease |
| Chiron et al. | FBI | 4 Km ² | 1 year | 643 801 Km ² | 14 years | 2007-2020 | France | Decrease |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|-----------------------------|------------------------|-------------------------|----------------|-------------------------|-----------------|-----------|-------------------|----------|
| Eglinton and Pearce-Higgins | FBI | 1 Km ² | 1 year | 242 495 km ² | 39 years | 1970-2008 | UK | Decrease |
| Harrison et al. (a) | Geometric mean | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Decrease |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Increase |
| Harrison et al. (a) | Geometric mean | 62 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Increase |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 62 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 62 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Increase |
| Harrison et al. (a) | Geometric mean | 16 000 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 16 000 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 16 000 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Stable |
| Harrison et al. (a) | Geometric mean | 130 000 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Decrease |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|------------------------|------------------------|-------------------------|----------------|-------------------------|-----------------|-----------|-------------------|----------|
| Harrison et al. (a) | GoF ($\lambda = -1$) | 131 000 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Decrease |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 132 000 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Stable |
| Harrison et al. (a) | Geometric mean | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Increase |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 10 000 Km ² | 1 year | 200 000 km ² | 18 years | 1994-2011 | Great Britain, UK | Decrease |
| Harrison et al. (a) | Geometric mean | 14 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Increase |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 14 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Increase |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 14 000 Km ² | 1 year | 77 933 Km ² | 18 years | 1994-2011 | Scotland, UK | Stable |
| Harrison et al. (a) | Geometric mean | 32 300 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Increase |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 32 300 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Decrease |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 32 300 Km ² | 1 year | 130 279 Km ² | 18 years | 1994-2011 | England, UK | Stable |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|------------------------|---------------------------------|-------------------------|----------------|---------------------------|-----------------|-----------|--|----------|
| Harrison et al. (a) | Geometric mean | 3 116 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -1$) | 3 116 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Stable |
| Harrison et al. (a) | GoF ($\lambda = -2$) | 3 116 Km ² | 1 year | 20,779 Km ² | 18 years | 1994-2011 | Wales, UK | Increase |
| Juslén et al. | RLI (Red list Index) | 338 440 km ² | 1 year | 338 440 km ² | 10 years | 2001-2010 | Finland | Decrease |
| Normander et al. | BCI (Biodiversity Change Index) | 84 266 km ² | NA | 1 260 663 km ² | 16 years | 1990-2005 | Finland, Sweden, Norway, Denmark and Iceland | Decrease |
| Normander et al. | BCI (Biodiversity Change Index) | 529 831 km ² | NA | 1 260 663 km ² | 16 years | 1990-2005 | Finland, Sweden, Norway, Denmark and Iceland | Stable |
| Normander et al. | BCI (Biodiversity Change Index) | 163 131 km ² | NA | 1 260 663 km ² | 16 years | 1990-2005 | Finland, Sweden, Norway, Denmark and Iceland | Decrease |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|-----------------|----------------|--------------------|----------------|----------------------------|-----------------|-----------|---------------------|----------|
| Schipper et al. | Geometric mean | 32 Km ² | 5 years | 21 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Geometric mean | 32 Km ² | 5 years | 21 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Geometric mean | 32 Km ² | 5 years | 21 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Geometric mean | 32 Km ² | 5 years | 22 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Geometric mean | 32 Km ² | 5 years | 23 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Stable |
| Schipper et al. | SR | 32 Km ² | 5 years | 24 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | SR | 32 Km ² | 5 years | 25 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | SR | 32 Km ² | 5 years | 26 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | SR | 32 Km ² | 5 years | 27 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | SR | 32 Km ² | 5 years | 28 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Stable |
| Schipper et al. | Shannon | 32 Km ² | 5 years | 29 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|-----------------|---------------------|--------------------|----------------|----------------------------|-----------------|-----------|------------------------|----------|
| Schipper et al. | Shannon | 32 Km ² | 5 years | 30 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Shannon | 32 Km ² | 5 years | 31 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Shannon | 32 Km ² | 5 years | 32 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Shannon | 32 Km ² | 5 years | 33 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Simpson | 32 Km ² | 5 years | 34 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Simpson | 32 Km ² | 5 years | 35 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Stable |
| Schipper et al. | Simpson | 32 Km ² | 5 years | 36 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Simpson | 32 Km ² | 5 years | 37 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Simpson | 32 Km ² | 5 years | 38 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional richness | 32 Km ² | 5 years | 39 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional richness | 32 Km ² | 5 years | 40 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|-----------------|-----------------------|--------------------|----------------|----------------------------|-----------------|-----------|---------------------|----------|
| Schipper et al. | Functional richness | 32 Km ² | 5 years | 41 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional richness | 32 Km ² | 5 years | 42 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional richness | 32 Km ² | 5 years | 43 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Functional evenness | 32 Km ² | 5 years | 44 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional evenness | 32 Km ² | 5 years | 45 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional evenness | 32 Km ² | 5 years | 46 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Functional evenness | 32 Km ² | 5 years | 47 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional evenness | 32 Km ² | 5 years | 48 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional divergence | 32 Km ² | 5 years | 49 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Functional divergence | 32 Km ² | 5 years | 50 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional divergence | 32 Km ² | 5 years | 51 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|-----------------|-------------------------|----------------------|----------------|----------------------------|-----------------|-----------|---------------------|----------|
| Schipper et al. | Functional divergence | 32 Km ² | 5 years | 52 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Functional divergence | 32 Km ² | 5 years | 53 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Schipper et al. | Functional dispersion | 32 Km ² | 5 years | 54 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional dispersion | 32 Km ² | 5 years | 55 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional dispersion | 32 Km ² | 5 years | 56 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional dispersion | 32 Km ² | 5 years | 57 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Increase |
| Schipper et al. | Functional dispersion | 32 Km ² | 5 years | 58 792 000 km ² | 40 years | 1971-2010 | Canada, USA, Mexico | Decrease |
| Pilotto et al. | SR | 1402 Km ² | 0.08 year | 1402 Km ² | 27 years | 1991-2017 | Belgium | Decrease |
| Pilotto et al. | Simpson | 1402 Km ² | 0.08 year | 1402 Km ² | 27 years | 1991-2017 | Belgium | Increase |
| Pilotto et al. | Abundance | 1402 Km ² | 0.08 year | 1402 Km ² | 27 years | 1991-2017 | Belgium | Decrease |
| Pilotto et al. | Temporal beta-diversity | 1402 Km ² | 0.08 year | 1402 Km ² | 27 years | 1991-2017 | Belgium | Stable |
| Pilotto et al. | SR | 509 Km ² | 1 year | 509 Km ² | 42 years | 1976-2017 | Bulgaria | Increase |
| Pilotto et al. | Simpson | 509 Km ² | 1 year | 509 Km ² | 42 years | 1976-2017 | Bulgaria | Stable |
| Pilotto et al. | Abundance | 509 Km ² | 1 year | 509 Km ² | 42 years | 1976-2017 | Bulgaria | Increase |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|----------------|----------------------------|------------------------|----------------|------------------------|-----------------|-----------|----------|----------|
| Pilotto et al. | Temporal beta-diversity | 509 Km ² | 1 year | 509 Km ² | 42 years | 1976-2017 | Bulgaria | Decrease |
| Pilotto et al. | SR | 10 Km ² | 1 year | 10 Km ² | 42 years | 1976-2017 | Bulgaria | Increase |
| Pilotto et al. | Simpson | 10 Km ² | 1 year | 10 Km ² | 42 years | 1976-2017 | Bulgaria | Stable |
| Pilotto et al. | Abundance | 10 Km ² | 1 year | 10 Km ² | 42 years | 1976-2017 | Bulgaria | Stable |
| Pilotto et al. | Temporal beta-diversity | 10 Km ² | 1 year | 10 Km ² | 42 years | 1976-2017 | Bulgaria | Stable |
| Pilotto et al. | SR | 9.02 Km ² | 1 year | 9.02 Km ² | 41 years | 1977-2017 | Bulgaria | Increase |
| Pilotto et al. | Simpson | 9.02 Km ² | 1 year | 9.02 Km ² | 41 years | 1977-2017 | Bulgaria | Increase |
| Pilotto et al. | Abundance | 9.02 Km ² | 1 year | 9.02 Km ² | 41 years | 1977-2017 | Bulgaria | Stable |
| Pilotto et al. | Temporal beta-diversity | 9.02 Km ² | 1 year | 9.02 Km ² | 41 years | 1977-2017 | Bulgaria | Stable |
| Pilotto et al. | SR | 32 Km ² | 1 year | 32 Km ² | 55 years | 1961-2015 | Spain | Increase |
| Pilotto et al. | Simpson | 32 Km ² | 1 year | 32 Km ² | 55 years | 1961-2015 | Spain | Increase |
| Pilotto et al. | Abundance | 32 Km ² | 1 year | 32 Km ² | 55 years | 1961-2015 | Spain | Increase |
| Pilotto et al. | Temporal beta-diversity | 32 Km ² | 1 year | 32 Km ² | 55 years | 1961-2015 | Spain | Decrease |
| Pilotto et al. | SR | 52 000 Km ² | 1 year | 52 000 Km ² | 55 years | 1961-2015 | France | Stable |
| Pilotto et al. | Simpson | 52 000 Km ² | 1 year | 52 000 Km ² | 55 years | 1961-2015 | France | Stable |
| Pilotto et al. | Abundance | 52 000 Km ² | 1 year | 52 000 Km ² | 55 years | 1961-2015 | France | Stable |
| Pilotto et al. | Temporal beta-diversity | 52 000 Km ² | 1 year | 52 000 Km ² | 55 years | 1961-2015 | France | Stable |

Table 2.1: SR = species richness, Ab = abundance, Eve = evenness, (*continued*)

| Reference | Metric | Spatial grain | Temporal grain | Spatial extent | Temporal extent | Years | Country | Trend |
|----------------|----------------------------|-----------------------|----------------|-----------------------|-----------------|-----------|-------------|----------|
| Pilotto et al. | SR | 6 155 Km ² | 0.08 year | 6 155 Km ² | 43 years | 1975-2017 | Netherlands | Increase |
| Pilotto et al. | Simpson | 6 155 Km ² | 0.08 year | 6 155 Km ² | 43 years | 1975-2017 | Netherlands | Increase |
| Pilotto et al. | Abundance | 6 155 Km ² | 0.08 year | 6 155 Km ² | 43 years | 1975-2017 | Netherlands | Increase |
| Pilotto et al. | Temporal beta-diversity | 6 155 Km ² | 0.08 year | 6 155 Km ² | 43 years | 1975-2017 | Netherlands | Stable |
| Pilotto et al. | SR | 2 180 Km ² | 1 year | 2 180 Km ² | 45 years | 1974-2018 | Netherlands | Decrease |
| Pilotto et al. | Simpson | 2 180 Km ² | 1 year | 2 180 Km ² | 45 years | 1974-2018 | Netherlands | Increase |
| Pilotto et al. | Abundance | 2 180 Km ² | 1 year | 2 180 Km ² | 45 years | 1974-2018 | Netherlands | Stable |
| Pilotto et al. | Temporal beta-diversity | 2 180 Km ² | 1 year | 2 180 Km ² | 45 years | 1974-2018 | Netherlands | Decrease |

Supplementary materials

Table 2.2: Supplementary informations on each trend

| Reference | Trend | Note |
|---------------------|----------|--|
| Barnagaud et al. | Increase | circle of radius 400 m |
| Barnagaud et al. | Decrease | NA |
| Barnagaud et al. | Increase | NA |
| Barnagaud et al. | Increase | NA |
| Barnagaud et al. | Stable | NA |
| Barnagaud et al. | Increase | NA |
| Roels et al. | Increase | Before/after tree planting (increase 11 times) |
| Roels et al. | Increase | Before/after tree planting (increase 3 times) |
| Wretenberg et al. | Decrease | looking at the trend through different environmental policies, " local species richness (i.e. at the scale of sites) decreased significantly probably as a result of an overall reduced abundance of several species. " |
| Ram et al. | Increase | forest species, road of 8 Km with no limitations so assumed 200m, "species richness (the average number of species seen per route and year) " |
| Ram et al. | Stable | forest specialist species, road of 8 Km with no limitations so assumed 200m |
| Ram et al. | Increase | generalist species, road of 8 Km with no limitations so assumed 200m |
| Ram et al. | Increase | specialist species, road of 8 Km with no limitations so assumed 200m |
| Ram et al. | Increase | generalists species, road of 8 Km with no limitations so assumed 200m |
| Harrison et al. (b) | Increase | "Biodiversity as measured by the geometric mean of relative abundances has generally increased since 1994", Visited twice a year / Increase first half and second second half |
| Harrison et al. (b) | Stable | " The goodness-of-fit-based measure of biodiversity suggests that both rare and common species made gains through much of Britain in the first half of the time period, and losses in the second half.", Visited twice a year / Increase first half and second second half |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|---------------------|----------|---|
| Harrison et al. (b) | Stable | " The goodness-of-fit-based measure of biodiversity suggests that both rare and common species made gains through much of Britain in the first half of the time period, and losses in the second half.", Visited twice a year / Increase first half and second second half |
| Doxa et al. | Increase | Not sure for the spatial scales, HNV +6.5% |
| Doxa et al. | Stable | Not sure for the spatial scales, for HNV +1.1%, Decrease then come back to the initial value |
| Doxa et al. | Stable | Not sure for the spatial scales, national, Decrease then come back to the initial value |
| Arnold et al. | Stable | Non significant slight increase |
| Arnold et al. | Stable | Non significant slight increase |
| Arnold et al. | Stable | Non significant slight increase |
| Xu et al. | Decrease | Not sure for the spatial scales, Urbanisation of the study area |
| Jiguet et al. | Increase | Not sure for the spatial scales, Generalist Bird Indicator, +20% |
| Jiguet et al. | Increase | Not sure for the spatial scales, Woodland Bird Indicator, -12% |
| Jiguet et al. | Increase | Not sure for the spatial scales, Urban Bird Indicator, -21% |
| Jiguet et al. | Increase | Not sure for the spatial scales, Farmland Bird Indicator, -12% |
| Jiguet et al. | Increase | Not sure for the spatial scales, plus 23% |
| Jiguet et al. | Decrease | minus 75% |
| Keten | Stable | NA |
| Davey et al. | Increase | NA |
| Davey et al. | Increase | NA |
| Davey et al. | Increase | NA |
| Christian et al. | Increase | Temporal grains varies a lot,significant increase of SR |
| Dittrich et al. | Increase | Spatial grain is the mean area of the orchards, increase sr may be due to increase in sampling effort (2 months for the first period and five for the 2nd and 3rd periods) |

Table 2.2: Supplementary informations on each trend *(continued)*

| Reference | Trend | Note |
|---------------------|----------|--|
| Dittrich et al. | Increase | Spatial grain is the mean area of the orchards, increase sr may be due to increase in sampling effort (2 months for the first period and five for the 2nd and 3rd periods) |
| Dittrich et al. | Stable | Increase until April then decrease but overall stable, spatial grain is the mean area of the orchards |
| Dittrich et al. | Stable | Spatial grain is the mean area of the orchards |
| Sirami and Monadjem | Decrease | NA |
| García-Navas et al. | Decrease | sorensen score |
| McGeoch et al. | Decrease | Red list index data, very heterogeneous |
| Ellis et al. | Stable | Riparian continuous ecosystem, area = $32 \times \pi \times (402)$, spatial extent = Bear Valley of southern Grant County and Silvies Valley of northern Harney County in east-central Oregon, slight variations in sr but not exceeding se |
| Ellis et al. | Stable | Riparian discontinuous ecosystem, area = $32 \times \pi \times (402)$ |
| Ellis et al. | Increase | Riparian herbaceous ecosystem, area = $32 \times \pi \times (402)$ |
| Ellis et al. | Stable | Riparian total ecosystem, area = area = $32 \times \pi \times (402)$ |
| Ellis et al. | Increase | Riparian continuous ecosystem, area = $32 \times \pi \times (402)$, spatial extent = Bear Valley of southern Grant County and Silvies Valley of northern Harney County in east-central Oregon, slight variations in sr but not exceeding se |
| Ellis et al. | Decrease | Riparian discontinuous ecosystem, slight decrease, area = $32 \times \pi \times (402)$ |
| Ellis et al. | Increase | Riparian herbaceous ecosystem, area = $32 \times \pi \times (402)$ |
| Ellis et al. | Increase | Riparian total ecosystem, area = area = $32 \times \pi \times (402) \times 3$ |
| Ellis et al. | Increase | Riparian continuous ecosystem, area = $32 \times \pi \times (402)$, spatial extent = Bear Valley of southern Grant County and Silvies Valley of northern Harney County in east-central Oregon, slight variations in sr but not exceeding se |
| Ellis et al. | Decrease | Riparian discontinuous ecosystem, area = $32 \times \pi \times (402)$ |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|------------------------------|----------|--|
| Ellis et al. | Increase | Riparian herbaceous ecosystem, area = $32 \times \pi \times (402)$ |
| Ellis et al. | Decrease | Riparian total ecosystem, slight decrease, area = $32 \times \pi \times (402)$ |
| Sicurella et al. | Stable | NPA non protected area |
| Sicurella et al. | Stable | NR nature reserves |
| Sicurella et al. | Increase | RP regional parks |
| Nally | Increase | NA |
| Latta et al. | Decrease | 54 to 31 species, untouched forest |
| Latta et al. | Decrease | 67 to 30 species, introduced tree species in the forest |
| Scarton | Increase | 14 to 25 species |
| Scarton | Increase | 2.07 to 2.38 |
| Scarton | Increase | 2.07 to 2.38 |
| Scarton | Increase | 2.07 to 2.38 |
| Chiron et al. | Decrease | Prediction with baseline scenario |
| Chiron et al. | Decrease | Prediction with CAP greening cenario |
| Chiron et al. | Decrease | Prediction with No Pillar I scenario |
| Chiron et al. | Decrease | Prediction with biofuel scenario |
| Eglington and Pearce-Higgins | Decrease | From 1 to 0.5 |
| Harrison et al. (a) | Stable | Farmland communities |
| Harrison et al. (a) | Decrease | Farmland communities, GoF weighted towards the rare species |
| Harrison et al. (a) | Increase | Farmland communities, GoF weighted towards the common species |
| Harrison et al. (a) | Increase | Farmland communities |
| Harrison et al. (a) | Stable | Farmland communities, GoF weighted towards the rare species |
| Harrison et al. (a) | Increase | Farmland communities, GoF weighted towards the common species |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|---------------------|----------|---|
| Harrison et al. (a) | Stable | Farmland communities |
| Harrison et al. (a) | Stable | Farmland communities, GoF weighted towards the rare species |
| Harrison et al. (a) | Stable | Farmland communities, GoF weighted towards the common species |
| Harrison et al. (a) | Decrease | Farmland communities |
| Harrison et al. (a) | Decrease | Farmland communities, GoF weighted towards the rare species |
| Harrison et al. (a) | Stable | Farmland communities, GoF weighted towards the common species |
| Harrison et al. (a) | Increase | Woodland communities, supplementary material |
| Harrison et al. (a) | Stable | Woodland communities, supplementary material |
| Harrison et al. (a) | Decrease | Woodland communities, supplementary material |
| Harrison et al. (a) | Increase | Woodland communities |
| Harrison et al. (a) | Increase | Woodland communities |
| Harrison et al. (a) | Stable | not sure for the trend, Woodland communities |
| Harrison et al. (a) | Increase | Woodland communities |
| Harrison et al. (a) | Decrease | Not sure for the trend, Woodland communities |
| Harrison et al. (a) | Stable | Not sure for the trend, Woodland communities |
| Harrison et al. (a) | Stable | Not sure for the trend, Woodland communities |
| Harrison et al. (a) | Stable | Not sure for the trend, Woodland communities |
| Harrison et al. (a) | Increase | Not sure for the trend, Woodland communities |
| Juslén et al. | Decrease | NA |
| Normander et al. | Decrease | Farmland |
| Normander et al. | Stable | Forest |
| Normander et al. | Decrease | Mires |
| Schipper et al. | Increase | All |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|-----------------|----------|-----------|
| Schipper et al. | Decrease | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Stable | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Decrease | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Stable | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Decrease | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Increase | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Stable | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Increase | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Decrease | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|-----------------|----------|---|
| Schipper et al. | Decrease | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Increase | Grassland |
| Schipper et al. | Decrease | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Increase | Shrubland |
| Schipper et al. | Decrease | All |
| Schipper et al. | Increase | Grassland |
| Schipper et al. | Decrease | Woodland |
| Schipper et al. | Decrease | Wetland |
| Schipper et al. | Decrease | Shrubland |
| Schipper et al. | Increase | All |
| Schipper et al. | Increase | Grassland |
| Schipper et al. | Increase | Woodland |
| Schipper et al. | Increase | Wetland |
| Schipper et al. | Decrease | Shrubland |
| Pilotto et al. | Decrease | Dataset S004, temporal grain = 1 month |
| Pilotto et al. | Increase | Dataset S004, temporal grain = 1 month |
| Pilotto et al. | Decrease | Dataset S004, temporal grain = 1 month |
| Pilotto et al. | Stable | Dataset S004, temporal grain = 1 month, p-val = 0.8 |
| Pilotto et al. | Increase | Dataset S011 |
| Pilotto et al. | Stable | Dataset S011, p-val = 0.8 |
| Pilotto et al. | Increase | Dataset S011 |

Table 2.2: Supplementary informations on each trend (*continued*)

| Reference | Trend | Note |
|----------------|----------|----------------------------|
| Pilotto et al. | Decrease | Dataset S011 |
| Pilotto et al. | Increase | Dataset S012 |
| Pilotto et al. | Stable | Dataset S012, p-val = 0.1 |
| Pilotto et al. | Stable | Dataset S012, p-val = 0.22 |
| Pilotto et al. | Stable | Dataset S012, p-val = 0.9 |
| Pilotto et al. | Increase | Dataset S013 |
| Pilotto et al. | Increase | Dataset S013 |
| Pilotto et al. | Stable | Dataset S013 |
| Pilotto et al. | Stable | Dataset S013 |
| Pilotto et al. | Increase | Dataset S047 |
| Pilotto et al. | Increase | Dataset S047 |
| Pilotto et al. | Increase | Dataset S047 |
| Pilotto et al. | Decrease | Dataset S047 |
| Pilotto et al. | Stable | Dataset S076 |
| Pilotto et al. | Stable | Dataset S076 |
| Pilotto et al. | Stable | Dataset S076 |
| Pilotto et al. | Stable | Dataset S076 |
| Pilotto et al. | Increase | Dataset S094 |
| Pilotto et al. | Increase | Dataset S094 |
| Pilotto et al. | Increase | Dataset S094 |
| Pilotto et al. | Stable | Dataset S094 |
| Pilotto et al. | Decrease | Dataset S095 |
| Pilotto et al. | Increase | Dataset S095 |

Table 2.2: Supplementary informations on each trend *(continued)*

| Reference | Trend | Note |
|--------------------------------|----------|-----------------------------|
| Pilotto et al. | Stable | Dataset S095, p-val = 0.056 |
| Pilotto et al. | Decrease | Dataset S095 |

References

- Arnold, H., Deacon, A. E., Hulme, M. F., Sansom, A., Jaggernauth, D., and Magurran, A. E. Contrasting trends in biodiversity of birds and trees during succession following cacao agroforest abandonment. 58(6):1248–1260. _eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.13869>.
- Barnagaud, J.-Y., Gaüzère, P., Zuckerberg, B., Princé, K., and Svenning, J.-C. Temporal changes in bird functional diversity across the united states. 185(4):737–748.
- Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., and Ferrer, E. A. Has the earth's sixth mass extinction already arrived? 471(7336):51–57. Number: 7336 Publisher: Nature Publishing Group.
- Bejček, V. and Stastný. Velké ptačí mapování.
- Blowes, S. A., Supp, S. R., Antão, L. H., Bates, A., Bruelheide, H., Chase, J. M., Moyes, F., Magurran, A., McGill, B., Myers-Smith, I. H., Winter, M., Bjorkman, A. D., Bowler, D. E., Byrnes, J. E. K., Gonzalez, A., Hines, J., Isbell, F., Jones, H. P., Navarro, L. M., Thompson, P. L., Vellend, M., Waldock, C., and Dornelas, M. The geography of biodiversity change in marine and terrestrial assemblages. 366(6463):339–345. Publisher: American Association for the Advancement of Science Section: Research Article.
- Butchart, S., Stattersfield, A., Baillie, J., Bennun, L., Stuart, S., Akçakaya, H., Hilton-Taylor, C., and Mace, G. Using red list indices to measure progress towards the 2010 target and beyond. 360(1454):255–268. Publisher: Royal Society.
- Butchart, S. H. M., Akçakaya, H. R., Chanson, J., Baillie, J. E. M., Collen, B., Quader, S., Turner, W. R., Amin, R., Stuart, S. N., and Hilton-Taylor, C. Improvements to the red list index. 2(1):e140. Publisher: Public Library of Science.
- Butchart, S. H. M., Stattersfield, A. J., Bennun, L. A., Shutes, S. M., Akçakaya, H. R., Baillie, J. E. M., Stuart, S. N., Hilton-Taylor, C., and Mace, G. M. Measuring global trends in the status of biodiversity: Red list indices for birds. 2(12):e383.
- Chase, J. M., McGill, B. J., Thompson, P. L., Antão, L. H., Bates, A. E., Blowes, S. A., Dornelas, M., Gonzalez, A., Magurran, A. E., Supp, S. R., Winter, M., Bjorkman, A. D., Bruelheide, H., Byrnes, J. E. K., Cabral, J. S., Elahi, R., Gomez, C., Guzman, H. M., Isbell, F., Myers-Smith, I. H., Jones, H. P., Hines, J., Vellend, M., Waldock, C., and O'Connor, M. Species richness change across spatial scales. 128(8):1079–1091.
- Chiron, F., Princé, K., Paracchini, M. L., Bulgheroni, C., and Jiguet, F. Forecasting the potential impacts of CAP-associated land use changes on farmland birds at the national level. 176:17–23.
- Christian, K., Isabelle, L. V., Frédéric, J., and Vincent, D. More species, fewer specialists: 100 years of changes in community composition in an island biogeographical study. 15(4):641–648. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1472-4642.2009.00569.x>.
- Davey, C. M., Chamberlain, D. E., Newson, S. E., Noble, D. G., and Johnston, A. Rise of the generalists: evidence for climate driven homogenization in avian communities. 21(5):568–578. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1466-8238.2011.00693.x>.

- Dittrich, R., Giessing, B., Benito, M. M., Russ, A., Wolf, C., Foudoulakis, M., and Norman, S. Multiyear monitoring of bird communities in chlorpyrifos-treated orchards in Spain and the United Kingdom: Spatial and temporal trends in species composition, abundance, and site fidelity. 38(3):616–629. _eprint: <https://setac.onlinelibrary.wiley.com/doi/pdf/10.1002/etc.4317>.
- Dornelas, M., Magurran, A. E., Buckland, S. T., Chao, A., Chazdon, R. L., Colwell, R. K., Curtis, T., Gaston, K. J., Gotelli, N. J., Kosnik, M. A., McGill, B., McCune, J. L., Morlon, H., Mumby, P. J., Øvreås, L., Studeny, A., and Vellend, M. Quantifying temporal change in biodiversity: challenges and opportunities. 280(1750):20121931.
- Doxa, A., Bas, Y., Paracchini, M. L., Pointereau, P., Terres, J.-M., and Jiguet, F. Low-intensity agriculture increases farmland bird abundances in France. 47(6):1348–1356. _eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2010.01869.x>.
- Eglington, S. M. and Pearce-Higgins, J. W. Disentangling the relative importance of changes in climate and land-use intensity in driving recent bird population trends. 7(3):e30407.
- Ellis, M. S., Kennedy, P. L., Edge, W. D., and Sanders, T. A. Twenty-year changes in riparian bird communities of east-central Oregon. 131(1):43–61. Publisher: The Wilson Ornithological Society.
- García-Navas, V., Sattler, T., Schmid, H., and Ozgul, A. Temporal homogenization of functional and beta diversity in bird communities of the Swiss Alps. 26(8):900–911. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/ddi.13076>.
- Gregory, R. D. and Strien, A. v. Wild bird indicators: using composite population trends of birds as measures of environmental health. 9(1):3–22.
- Gregory, R. D., van Strien, A., Vorisek, P., Gmelig Meyling, A. W., Noble, D. G., Foppen, R. P., and Gibbons, D. W. Developing indicators for European birds. 360(1454):269–288. Publisher: Royal Society.
- Gregory, R. D., Vorisek, P., Strien, A. V., Meyling, A. W. G., Jiguet, F., Fornasari, L., Reif, J., Chylarecki, P., and Burfield, I. J. Population trends of widespread woodland birds in Europe. 149:78–97. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1474-919X.2007.00698.x>.
- Harrison, P. J., Buckland, S. T., Yuan, Y., Elston, D. A., Brewer, M. J., Johnston, A., and Pearce-Higgins, J. W. Assessing trends in biodiversity over space and time using the example of British breeding birds. 51(6):1650–1660. _eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.12316>.
- Harrison, P. J., Yuan, Y., Buckland, S. T., Oedekoven, C. S., Elston, D. A., Brewer, M. J., Johnston, A., and Pearce-Higgins, J. W. Quantifying turnover in biodiversity of British breeding birds. 53(2):469–478. _eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.12539>.
- Hill, M. O. Diversity and evenness: A unifying notation and its consequences. 54(2):427–432. _eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1934352>.
- Jaccard, P. The distribution of the flora in the alpine zone. 1. 11(2):37–50. _eprint: <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.1912.tb05611.x>.

- Jiguet, F., Devictor, V., Julliard, R., and Couvet, D. French citizens monitoring ordinary birds provide tools for conservation and ecological sciences. 44:58–66.
- Juslén, A., Hyvärinen, E., and Virtanen, L. K. Application of the red-list index at a national level for multiple species groups. 27(2):398–406. _eprint: <https://conbio.onlinelibrary.wiley.com/doi/pdf/10.1111/cobi.12016>.
- Keil, P., Schweiger, O., Kühn, I., Kunin, W. E., Kuussaari, M., Settele, J., Henle, K., Brotons, L., Pe'er, G., Lengyel, S., Moustakas, A., Steinicke, H., and Storch, D. Patterns of beta diversity in europe: the role of climate, land cover and distance across scales. 39(8):1473–1486. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2699.2012.02701.x>.
- Keten. Temporal patterns of wetland-associated bird assemblages in altered wetlands in turkey.
- Latta, S. C., Tinoco, B. A., Astudillo, P. X., and Graham, C. H. Patterns and magnitude of temporal change in avian communities in the ecuadorian andes. 113(1):24–40.
- McGeoch, M. A., Butchart, S. H. M., Spear, D., Marais, E., Kleynhans, E. J., Symes, A., Chanson, J., and Hoffmann, M. Global indicators of biological invasion: species numbers, biodiversity impact and policy responses. 16(1):95–108. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1472-4642.2009.00633.x>.
- Nally, R. M. Monitoring forest bird communities for impact assessment: The influence of sampling intensity and spatial scale. 82(3):355–367.
- Normander, B., Levin, G., Auvinen, A.-P., Bratli, H., Stabbetorp, O., Hedblom, M., Glimskär, A., and Gudmundsson, G. A. Indicator framework for measuring quantity and quality of biodiversity—exemplified in the nordic countries. 13(1):104–116.
- Pereira, H. M., Navarro, L. M., and Martins, I. S. Global biodiversity change: The bad, the good, and the unknown. 37(1):25–50.
- Pilotto, F., Kühn, I., Adrian, R., Alber, R., Alignier, A., Andrews, C., Bäck, J., Barbaro, L., Beaumont, D., Beenaerts, N., Benham, S., Boukal, D. S., Bretagnolle, V., Camatti, E., Canullo, R., Cardoso, P. G., Ens, B. J., Everaert, G., Evtimova, V., Feuchtmayr, H., García-González, R., Gómez García, D., Grandin, U., Gutowski, J. M., Hadar, L., Halada, L., Halassy, M., Hummel, H., Huttunen, K.-L., Jaroszewicz, B., Jensen, T. C., Kalivoda, H., Schmidt, I. K., Kröncke, I., Leinonen, R., Martinho, F., Meesenburg, H., Meyer, J., Minerbi, S., Monteith, D., Nikolov, B. P., Oro, D., Ozoliņš, D., Padedda, B. M., Pallett, D., Pansera, M., Pardal, M. n., Petriccione, B., Pipan, T., Pöyry, J., Schäfer, S. M., Schaub, M., Schneider, S. C., Skuja, A., Soetaert, K., Sprinġe, G., Stanchev, R., Stockan, J. A., Stoll, S., Sundqvist, L., Thimonier, A., Van Hoey, G., Van Ryckegem, G., Visser, M. E., Vorhauser, S., and Haase, P. Meta-analysis of multidecadal biodiversity trends in europe. 11(1):3486. Bandiera_abtest: a Cc_license_type: cc_by Cg_type: Nature Research Journals Number: 1 Primary_atype: Research Publisher: Nature Publishing Group Subject_term: Biodiversity;Climate-change ecology;Macroecology Subject_term_id: biodiversity;climate-change-ecology;macroecology.
- Ram, D., Axelsson, A.-L., Green, M., Smith, H. G., and Lindström, k. What drives current population trends in forest birds – forest quantity, quality or climate? a large-scale analysis from northern europe. 385:177–188.

- Roels, S., Hannay, M., and Lindell, C. Recovery of bird activity and species richness in an early-stage tropical forest restoration. 14(1). Publisher: The Resilience Alliance.
- Scarton, F. Long-term trend of the waterbird community breeding in a heavily man-modified coastal lagoon: the case of the important bird area "lagoon of venice". 21(1):35–45.
- Schipper, A. M., Belmaker, J., Miranda, M. D. d., Navarro, L. M., Böhning-Gaese, K., Costello, M. J., Dornelas, M., Foppen, R., Hortal, J., Huijbregts, M. A. J., Martín-López, B., Pettorelli, N., Queiroz, C., Rossberg, A. G., Santini, L., Schiffrs, K., Steinmann, Z. J. N., Visconti, P., Rondinini, C., and Pereira, H. M. Contrasting changes in the abundance and diversity of north american bird assemblages from 1971 to 2010. 22(12):3948–3959. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/gcb.13292>.
- Shannon, C. E. A mathematical theory of communication. 27(3):379–423. Conference Name: The Bell System Technical Journal.
- Sicurella, B., Orioli, V., Pinoli, G., Ambrosini, R., and Bani, L. Effectiveness of the system of protected areas of lombardy (northern italy) in preserving breeding birds. 28(3):475–492. Publisher: Cambridge University Press.
- Simpson, E. H. Measurement of diversity. 163(4148):688–688. Bandiera_abtest: a Cg_type: Nature Research Journals Number: 4148 Primary_atype: Research Publisher: Nature Publishing Group.
- Sirami, C. and Monadjem, A. Changes in bird communities in swaziland savannas between 1998 and 2008 owing to shrub encroachment. 18(4):390–400. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1472-4642.2011.00810.x>.
- Sørensen, T. J. *A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons*. I kommission hos E. Munksgaard. OCLC: 4713331.
- Studený, A. C., Buckland, S. T., Illian, J. B., Johnston, A., and Magurran, A. E. Goodness of fit measures of evenness: a new tool for exploring changes in community structure. 2(2):art15. _eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/ES10-00074.1>.
- The Convention on Biological Diversity, B. The convention on biological diversity. Publisher: Secretariat of the Convention on Biological Diversity.
- Whittaker, R. H. Evolution and measurement of species diversity. 21(2):213–251. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.2307/1218190>.
- Whittaker, R. H. Vegetation of the siskiyou mountains, oregon and california. 30(3):279–338. _eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1943563>.
- Wretenberg, J., Pärt, T., and Berg, k. Changes in local species richness of farmland birds in relation to land-use changes and landscape structure. 143(2):375–381.
- Xu, X., Xie, Y., Qi, K., Luo, Z., and Wang, X. Detecting the response of bird communities and biodiversity to habitat loss and fragmentation due to urbanization. 624:1561–1576.